



RESEARCH ARTICLE

Bacteria associated with grapevine (*Vitis vinifera* L.) rhizosphere and their efficacy in plant growth promotion

Bakhora Djalolova¹, Sokhibjon Abdusamatov¹, Vyacheslav Shurigin^{2*}, Dilfuza Jabborova³, Bakhora Turaeva⁴, Guzal Kutlieva⁴ & Kakhramon Davranov⁵

¹Department of Microbiology and Biotechnology, National University of Uzbekistan, Tashkent 100 174, Uzbekistan

²State Key Laboratory of Desert and Oasis Ecology, Key Laboratory of Ecological Safety and Sustainable Development in Arid Lands, Xinjiang Institute of Ecology and Geography, Chinese Academy of Sciences, Urumqi 830 011, China

³Laboratory of Genetics, Breeding and Selection of Pulses, Oilseeds and Medicinal Plants, Institute of Genetics and Plant Experimental Biology of the Academy of Sciences of the Republic of Uzbekistan, Tashkent 111 215, Uzbekistan

⁴Laboratory of Microbiology and Biotechnology of Probiotics, Institute of Microbiology of the Academy of Sciences of the Republic of Uzbekistan, Tashkent 100 128, Uzbekistan

⁵Laboratory of Enzymology, Institute of Microbiology of the Academy of Sciences of the Republic of Uzbekistan, Tashkent 100 128, Uzbekistan

*Email: slaventus87@hotmail.com



ARTICLE HISTORY

Received: 03 February 2024

Accepted: 03 September 2024

Available online

Version 1.0 : 29 December 2024



Additional information

Peer review: Publisher thanks Sectional Editor and the other anonymous reviewers for their contribution to the peer review of this work.

Reprints & permissions information is available at https://horizonepublishing.com/journals/index.php/PST/open_access_policy

Publisher's Note: Horizon e-Publishing Group remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Indexing: Plant Science Today, published by Horizon e-Publishing Group, is covered by Scopus, Web of Science, BIOSIS Previews, Clarivate Analytics, NAAS, UGC Care, etc See https://horizonepublishing.com/journals/index.php/PST/indexing_abstracting

Copyright: © The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited (<https://creativecommons.org/licenses/by/4.0/>)

CITE THIS ARTICLE

Djalolova B, Abdusamatov S, Shurigin V, Jabborova D, Turaeva B, Kutlieva G, Davranov K. Bacteria associated with grapevine (*Vitis vinifera* L.) rhizosphere and their efficacy in plant growth promotion. Plant Science Today (Early Access). <https://doi.org/10.14719/pst.3338>

Abstract

Plant growth, development and stress resistance depend on the presence of plant growth-promoting rhizobacteria (PGPR) in its rhizosphere. The detection of the highly active PGPR is of high importance due to their possible application as microbial inoculants for plant growth promotion (PGP) in agriculture. In this study, we report on PGPR from the rhizosphere of grapevine (*Vitis vinifera* L.) growing in Uzbekistan, as it was not studied before. Based on the screening of 37 isolates from grapevine rhizosphere for stimulation of wheat seed germination, just two isolates, BDI-1 and BDI-2 were chosen as the most active. In laboratory conditions, the isolates BDI-1 and BDI-2 increased wheat root length up to 1.48 and 1.5 times and shoot length up to 1.59 and 1.64 times, respectively, as compared to the control. Based on 16S rRNA gene analysis and comparison with the relative strains registered in GenBank of the National Center for Biotechnology Information (NCBI), the isolates BDI-1 and BDI-2 were identified as *Pantoea agglomerans* and *Priestia megaterium*, accordingly. Their 16S rRNA gene nucleotide sequences were deposited to GenBank under the accession numbers OP727725 for BDI-1 and OP782582 for BDI-2. Both isolates were phenotypically characterized and demonstrated phosphate-solubilizing and nitrogen-fixing abilities, producing indole-3-acetic acid (IAA) in a high amount; however, BDI-1 also produced siderophores and BDI-2-1-aminocyclopropane-1-carboxylate (ACC) - deaminase. Due to these features, the bacteria showed their high activity in the promotion of plant growth and seed germination. In conclusion, according to our results, *P. agglomerans* BDI-1 and *P. megaterium* BDI-2 are promising PGPR, which can be applied as microbial inoculants for plant growth improvement.

Keywords

Pantoea agglomerans; PGPR; *Priestia megaterium*; rhizosphere; *Vitis vinifera* L.; grapevine

Introduction

Vitis vinifera L., or grapevine, is a widely cultivated woody fruit crop whose annual grape production was 76,750,674 tons in 2022 (1). It was domesticated from its wild ancestor, *Vitis vinifera* subsp. *sylvestris* (Willd.) Hegi, naturally occurring from southwestern Asia and Mediterranean Europe (2). The growth and productivity of

a grapevine as well as of other plants depends on the availability of nutrients in the soil. However, the majority of nutrients in soil is in stable forms and cannot be consumed directly by plant roots. It is well known that microbial communities living in the rhizosphere, or inside of plants, play an important role in supporting plant growth and development in different soils and climatic zones using different mechanisms (3, 4). Plant growth-promoting rhizobacteria (PGPR) can make insoluble minerals containing phosphorus and potassium available to plants by solubilizing those using phosphatases and some organic acids like hydrogen cyanide (5, 6). Some bacteria can supply plants with available nitrogen using a special nitrogenase enzymatic complex, which converts atmospheric nitrogen to ammonium (NH_4^+) (7). PGPR can also produce phytohormones, such as auxin indole-3-acetic acid (IAA), improving root growth by elongation and branching, letting the plant root system cover more area and uptake more water and nutrients (8, 9). Some bacteria can contribute to plant growth by producing siderophores, which can bind iron and make it unavailable for some plant pathogenic fungi, thus protecting plants from diseases (10, 11). Other PGPR alleviates plant stress by cleaving an ethylene (stress hormone) precursor-1-aminocyclopropane-1-carboxylic acid with 1-aminocyclopropane-1-carboxylate (ACC) - deaminase (12). There were several reports about PGPR isolated from the rhizosphere of different crops such as soybean (*Glycine max*) and chilli (*Capsicum annuum*) in India (13), maize (*Zea mays*) in Africa (14), one report about grapevine (*Vitis vinifera*) in Turkey (15) and others crops. However, the microbial communities of PGPR vary in the rhizosphere of the same plant species grown in different geographical zones with specific climate conditions and soil types (16). Up to now, there were no reports about PGPR from the rhizosphere of a grapevine grown in Uzbekistan with its specific soils and climate.

Our aims were: 1) isolate the rhizosphere bacteria contributing to grapevine growth and fitness in local soils of Uzbekistan; 2) study some plant growth promoting (PGP) and phenotypic characteristics of the chosen bacteria.

Materials and Methods

Samples isolation

The light serozym soil samples in the amount of 10 were collected from the rhizosphere of a grapevine (*Vitis vinifera* L.) growing in the Samarqand region of the Kattakorgan district of Uzbekistan. The root system with the stuck soil particles was transferred into sterile plastic bags and brought to the laboratory for further analysis.

Bacteria isolation

The bacteria were isolated from the collected grapevine rhizosphere soil as follows: 10 g of the collected grapevine roots with rhizosphere soil was mixed thoroughly with 100 mL of sterile water and serially diluted up to 1×10^{-4} and 1×10^{-5} . 100 μL of final soil suspension was transferred and spread with a triangle onto the surface of Petri dishes with peptone agar. The dishes were left at 28°C for incubation and after 48 h, they were checked for the presence of bacterial colonies. The grown bacterial colonies, different in shape, size and color were isolated by transfer to individual Petri dishes with peptone agar and purified.

Screening for bacterial activity in the stimulation of seed germination

The isolated bacteria were tested in the stimulation of wheat (*Triticum aestivum* L.) seed germination. The isolated pure bacterial cultures were used as inoculants for wheat seeds to check their efficiency in stimulating seed germination. The bacteria were cultivated separately in nutrient broth at 28 °C for 72 h and the concentration of cells was adjusted to 10^7 CFU/mL. The seeds of wheat were externally sterilized by keeping in 2.5% NaClO for 3 min and rinsed in sterilized tap water (17). The outer sterilized seeds (100) were inoculated with each bacterial isolate by keeping them in bacterial suspension for 30 min and put into sterile Petri dishes with moist filter paper. As a control, the nutrient broth medium was used. The dishes were left in a dark place and the day-night temperature was 28-18 °C. The germination rate of seeds was recorded four times: after 12, 24, 36 and 48 h as a number of germinated seeds (from 100 seeds).

Screening of bacteria for plant growth-promoting properties

The bacteria were cultivated separately in nutrient broth at 28°C for 72 h and the concentration of cells was adjusted to 10^7 CFU/mL. The seeds of wheat were externally sterilized by keeping in 2.5 % NaClO for 3 min and rinsed in sterilized tap water (17). One hundred sterilized seeds were inoculated with each bacterial isolate by keeping them in bacterial suspension for 30 min and transferred into wide plastic pots with sterile light serozym soil. The pots were marked and set up in random places inside the plant growth chamber. We used pots in three replications and there were 10 seeds inoculated with the same bacterial isolate inside each pot. The plants were grown at 30 °C in the daytime and 20 °C at night and watered as required. The shoots and roots length were measured 10 days after seeds sowing.

Bacteria identification

The bacteria were identified using 16S rRNA gene analysis. To amplify the 16S rRNA region of the samples, we used a pair of primers BAK11w 5'-AGTTTGATCMTGGCTCAG-3' and BAK2 5'-GGACTACHAGGGTATCTAAT-3', which are optimal for identifying a wide range of bacterial DNA (18). The amplification reaction was carried out using an iCycler thermal cycler (Bio-Rad Laboratories, USA), programmed for the following reaction conditions: preliminary denaturation 96°C-1 min, 30 cycles, including 96°C-10 sec, 55°C-20 sec, 72°C-30 sec and final elongation 55°C-20 sec, 72°C-3 min. The volume of the reaction mixture was 25 μL , containing 0.3 mM of primers BAK11w and BAK2, 0.2 mM of each nucleotide, 2.5 μL of 10X Encyclo buffer, 0.5 μL of 50X Encyclo polymerase and 50 ng of DNA sample. The PCR products were analyzed in 1% agarose gel with the addition of ethidium bromide. The resulting PCR product was cut from the gel, purified and used in the sequencing reaction using the BigDye 3.1 Terminator Cycle Sequencing Ready Reaction Kit (Applied Biosystems, USA) according to the manufacturer's recommendations (Thermo Fisher Scientific, USA). The 16S rRNA gene sequences were compared with the identical sequences of the GenBank (NCBI) (<http://www.ncbi.nlm.nih.gov/>).

Study of the taxa evolutionary relationships

The neighbor-joining method was used to infer the evolutionary history (19). The associated taxa clustered together in the bootstrap test (500 replicates), according to Felsenstein (20). The tree was drawn to scale, with branch lengths in the same units as those of the evolutionary distances used to infer the phylogenetic tree. The Maximum Composite Likelihood method (21) was utilized to compute the evolutionary distances. The evolutionary analyses were conducted in MEGA X (22).

Study of bacteria PGP properties

The N₂-fixation of bacteria were checked by the ability to grow on malate nitrogen-deficient medium (g/L): sodium malate - 5, K₂HPO₄ - 0.5, KH₂PO₄ - 0.4, MgSO₄·7H₂O - 0.2, NaCl - 0.1, CaCl₂ - 0.02, FeCl₃ - 0.01, Na₂MoO₄·2H₂O - 0.002, agar - 15, pH 7.2-7.4, modified with yeast extract - 50 mg/L (23). Sarwar and Kremer's method (24) was used to check the production of IAA. The bacterial suspension was stabilized at 1×10^8 CFU/mL and poured into tubes containing 10 % tryptic soy agar (25) modified with L-tryptophan -5 mmol/L. After 24 h of cultivation at 28 °C, the suspension was centrifuged for 15 min at 8000×g and the supernatant was gathered. The Salkowski reagent (mixture of 7.9 mol/L H₂SO₄ and 0.5 mol/L FeCl₃) was mixed with the supernatant in a ratio of 1:1 (v/v) and stored for 30 min at room temperature in the dark. The presence of pink color indicated the production of IAA, which was measured with a spectrophotometer at 530 nm. The standard curve was drawn using different concentrations of IAA solutions. The phosphate solubilizing ability was tested by the cultivation of bacteria on NBRIP medium (%): glucose - 1, MgCl₂ - 0.5, Ca₃(PO₄)₂ - 0.5, MgSO₄·7H₂O - 0.025, KCl - 0.02, (NH₄)₂SO₄ - 0.01 and agar - 1.5. The formation of colonies after bacteria incubation at 28°C for 96 days indicated the ability to use Ca₃(PO₄)₂ as a sole phosphate source (26). The ACC - deaminase production was based on the ability of the tested bacteria to utilize 1-aminocyclopropane-1-carboxylate as a source of nitrogen. The basal medium, modified with ACC 3.0 mM was utilized for bacterial cultivation. As a positive control, we added (NH₄)₂SO₄ into media and didn't add any source of nitrogen for the negative (27). The siderophores-producing ability was tested by cultivating the bacteria on chrome

azuroil sulphonate agar (CAS) at 28 °C for 96 h. The colonies surrounded by an orange color zone were positive in siderophores production (28).

Phenotypic characterization of bacteria

The biochemical, morphological and physiological characteristics of bacteria isolates were identified based on the standardized methods recommended in Bergey's Manual of Systematic Bacteriology (29).

Statistical analysis

The analysis of variance in the Microsoft Excel 2010 package was utilized to test the data's statistical significance. The results of the plant growth experiment were subjected to analysis of variance (ANOVA) with SPSS software (v. 15) at $p \leq 0.05$. The least significant difference test ($p \leq 0.05$) was used for conducting mean comparisons. The results are presented as average means and standard error. All tests for seed germination, plant growth-promoting and phenotypic characteristics of bacteria were conducted minimum in three replications.

Results

Screening of bacteria for wheat seed germination

In total, 37 isolates different in their colonies color, size, density and shapes were isolated from the soil samples of a grapevine rhizosphere. The isolates were checked and screened for their activity in the stimulation of wheat seed germination in Petri dishes after 12, 24, 36 and 48 h (Fig. 1).

As you can see from Fig. 1, there were no germinated seeds in control, BDI-6, BDI-10, BDI-21 and BDI-30 after 12 h. However, at the same time, the germination rate after seed inoculation with bacterial isolates BDI-1, BDI-2, BDI-8, BDI-14, BDI-16, BDI-27 and BDI-37 was 7, 8, 6, 5, 5, 5 and 6 % consecutively. In 24 h, the same isolates BDI-1, BDI-2, BDI-8, BDI-14, BDI-16, BDI-27 and BDI-37 showed the highest seed germination rate: 38, 41, 34, 36, 33, 34 and 36 % consecutively in comparison with other bacterial isolates and control. The most interesting results were after 36 h when the seed germination rate increased up to 91 and 94 % in BDI-1 and BDI-2 accordingly and that was 1.9 and 2 times more than in the




Fig. 1. The influence of wheat seeds inoculation with isolated rhizosphere bacteria on germination rate (%) in 12, 24, 36 and 48 h.

* - statistically significant at $p \leq 0.05$

control. The other isolates showed results less than 90 % of seed germination rate. The isolates BDI-1 and BDI-2 showed the highest results in seed germination (98 %) after 48 h.

Screening of bacterial isolates for wheat growth promotion

The efficacy of the isolated rhizosphere bacteria in wheat growth stimulation was tested in a pot experiment (Fig. 2). The majority of the isolates improved plant growth after seed inoculation. The isolates BDI-8, BDI-14, BDI-16, BDI-27 and BDI-37 increased root length up to 1.38, 1.36, 1.36, 1.33 and 1.4 times consecutively in comparison with the control. However, the best activity demonstrated isolates BDI-1 and BDI-2, which increased root length up to 1.48 and 1.5 times and shoot length up to 1.59 and 1.64 times, respectively, as compared to the control.

Bacteria identification

The identification of the most active isolates, BDI-1 and BDI-2 was based on their 16S rRNA gene comparison with genes of relative strains registered in GenBank of NCBI (Table 1). According to 16S rRNA gene analysis, the isolate BDI-1 is related to *P. agglomerans*. It was deposited to GenBank under accession number OP727725 and called *P. agglomerans* BDI-1. The isolate BDI-2 was related to *P. megaterium* and deposited to GenBank under accession number OP782582. The phylogenetic trees of the strains *P. agglomerans* BDI-1 and *P. megaterium* BDI-2 and their closest relative strains from GenBank were constructed in MEGA X software using the neighbor-joining method (Fig. 3 and 4).

Plant growth-promoting characteristics of bacteria

The plant growth-promoting characteristics of the strains *P. agglomerans* BDI-1 and *P. megaterium* BDI-2 were studied (Table 2). Both strains showed phosphate-solubilizing and nitrogen-fixing abilities that produced IAA in high amounts. *P. megaterium* BDI-2 also produced ACC - deaminase and *P. agglomerans* BDI-1 - siderophores.




Fig. 3. The phylogenetic tree of the strain *Pantoea agglomerans* BDI-1 with the relative strains registered in GenBank (NCBI) based on 16S rRNA gene comparison.




Fig. 4. The phylogenetic tree of the strain *Priestia megaterium* BDI-2 with the relative strains registered in GenBank (NCBI) based on 16S rRNA gene comparison.




Fig. 2. The influence of wheat seeds inoculation with isolated rhizosphere bacteria on root and shoot length (10 days).

* - statistically significant at $p \leq 0.05$

Table 1. The active PGPR isolates from grapevine rhizosphere and their closest relatives from GenBank (NCBI)

Isolated bacteria deposited to GenBank			Closest match from GenBank based on 16S rRNA genes analysis		
Isolate	Length (bp)	Accession number	Reference strains	Accession number	Percent identity
BDI-1	1464	OP727725	<i>Pantoea agglomerans</i>	MT605811.1	99.73
BDI-2	1429	OP782582	<i>Priestia megaterium</i>	MT525296.1	99.72

Table 2. Plant growth-promoting and phenotypic characteristics of two active PGP bacterial strains

Characteristics	<i>Pantoea agglomerans</i> BDI-1	<i>Priestia megaterium</i> BDI-2
	Reaction	
<i>Plant growth-promoting traits</i>		
N ₂ -fixation	+	+
IAA (µg/mL) production	168.46	179.53
Phosphates solubilization	+	+
ACC-deaminase production	-	+
Siderophores production	+	-
<i>Phenotypic characteristics</i>		
Growth at 4°C	+	+
Growth at 44°C	-	+
Motility	+	+
Gelatin liquefaction	-	+
Voges-Proskauer reaction	+	-
Nitrate reduction to nitrite	+	-
Gas produced from D-glucose	-	-
Arginine dihydrolase	-	-
Phenilalanine deaminase	+	+
Catalase	+	+
Urease	-	+
<i>Acid produced from:</i>		
L-arabinose	+	-
D-ribose	+	+
D-xylose	+	-
D-fructose	+	+
D-mannose	+	-
L-rhamnose	+	-
L-sorbose	-	-
D-cellobiose	-	+
D-galactose	+	+
Lactose	-	+
Maltose	+	+
Melibiose	-	+
Sucrose	+	+
Trehalose	+	+
Melezitose	-	-
Raffinose	-	+
Glycerol	+	+
Meso-erithriol	-	-
Adonitol	-	-
Dulcitol	-	-
Inositol	-	+
D-mannitol	+	+
D-sorbitol	-	-
A- methyl-D-glucoside	-	-
Salicin	+	-
Inulin	-	+

Note: “+”- positive, “-”- negative

Phenotypic characteristics of bacteria

The phenotypic characteristics of bacteria *P. agglomerans* BDI-1 and *P. megaterium* BDI-2 are shown in Table 2. The strain *P. agglomerans* BDI-1 could grow at 4 °C but couldn't at 44 °C, was motile, couldn't liquefy gelatin, Voges-Proskauer reaction was positive, reduced nitrates to nitrites, couldn't produce gas from D-glucose, arginine dihydrolase phenylalanine negative, produced phenylalanine deaminase and catalase, but not urease. The strain produced acid from salicin, glycerol, trehalose, maltose, sucrose, D-mannitol, D-galactose, L-rhamnose, L-arabinose, D-mannose, D-fructose, D-xylose and D-ribose.

The strain *P. megaterium* BDI-2 grew at 4 °C and 44 °C, was motile, liquefied gelatin and produced phenylalanine deaminase, catalase and urease. The strain could not reduce nitrates to nitrites, did not produce arginine dihydrolase, the Voges-Proskauer reaction was negative and there was no gas formation from D-glucose. The strain produced acid from inulin, inositol, glycerol, sucrose, trehalose, raffinose, melibiose, maltose, lactose, D-mannitol, D-cellobiose, D-galactose, D-fructose and D-ribose.

Discussion

As a result of the isolation and screening of bacteria from the rhizosphere of a grapevine, just two out of 37 bacteria were chosen as active PGPR. The isolates BDI-1 and BDI-2 were highly efficient in wheat seed germination and PGP. It can be explained by the revealed properties of these isolates: nitrogen fixation, phosphate solubilization, production of IAA, siderophores and ACC-deaminase. Plants can get much benefit from free-living and symbiotic nitrogen-fixing bacteria because they can reduce atmospheric N_2 to NH_3 through the nitrogenase enzyme complex, making it accessible for plant uptake and fulfilling their nutritional requirements (30). Aasfar *et al.* (31) reported about improvement in wheat plant growth after seed bacterization with soil N_2 -fixing bacteria. Shurigin *et al.* (32) also reported the positive effect of nitrogen-fixing bacteria on the growth of cucumber. Thus, the ability of the strains *P. agglomerans* BDI-1 and *P. megaterium* BDI-2 to fix nitrogen contributes a lot to their plant growth-promoting efficacy. Phosphorus (P) is an essential element for plant development. Phosphorus plays a key role in root development and significantly contributes to plants resistance against diseases and increasing crop yield (33, 34). Phosphorus takes part in cellular division and the structure of nucleic acids (35). Most of the phosphates remain stable in the soil due to their insoluble forms and are unavailable to plants. The phosphate solubilizing ability of bacterial strains *P. agglomerans* BDI-1 and *P. megaterium* BDI-2 is a very beneficial characteristic because they can transform insoluble inorganic phosphates to soluble forms and make phosphorus available for plants (36). The strains *P. agglomerans* BDI-1 and *P. megaterium* BDI-2 produced IAA in relatively high concentrations. IAA is one of the most active phytohormones, improving plant growth (37). IAA synthesized by rhizobacteria increases root size, weight, root hairs and lateral root number (38, 39), thus increasing the area of contact with the surrounding soil. The increased availability of soil resources in larger areas and enhanced nutrient acquisition help to improve plant growth and yield (40). Iron is necessary due to its ability to sustain the life activities of a plant. The bioavailability of iron in soil is low because it mainly presents there in the form of insoluble ferric Fe (III) ions (41). Plant growth-promoting rhizobacteria improve iron uptake, using siderophores to chelate iron, providing plants with the required amount of iron (42, 43). The production of siderophores by the strains *P. agglomerans* BDI-1 and *P. megaterium* BDI-2 is one of the important factors in improving plant growth. Different kinds of stresses (drought, salinity, etc.) are known to increase ethylene production, which can inhibit root elongation, shorten vegetation time and decrease yield (44, 45). However, it was reported that ACC-deaminase-producing PGPR can improve the growth of plants in stressful conditions by reducing the level of ethylene due to the hydrolysis of its precursor - ACC, to α -ketobutyrate and ammonia (46, 47). *P. megaterium* BDI-2 producing ACC-deaminase could reduce stresses during plant growth.

Conclusion

For the first time, this study was reported about PGPR isolated from the rhizosphere of a grapevine grown in light serozym soils of Uzbekistan. Based on screening of 37 isolates for wheat seed germination stimulation and PGP, just two isolates, BDI-1 and BDI-2 were chosen as the most active. Based on 16S rRNA gene analysis, the strains BDI-1 and BDI-2 were identified as *Pantoea agglomerans* and *Priestia megaterium* accordingly. Both strains showed phosphate-solubilizing and nitrogen-fixing abilities and produced IAA; however, BDI-1 also produced siderophores and BDI-2-ACC-deaminase. The presence of these properties lets them show their high activity in plant growth and seed germination promotion. The bacteria *Pantoea agglomerans* BDI-1 and *Priestia megaterium* BDI-2 are promising PGPR, which can be applied as microbial inoculants for plant growth improvement.

Authors' contributions

BD and SA performed the experiments. VS wrote the draft of the manuscript. DJ statistically analyzed the results. BT and GK conducted the critical revision of the manuscript. KD worked out the concept and design, supervised and funded the experiments. All authors read and approved the final manuscript.

Compliance with ethical standards

Conflict of interest: Authors do not have any conflict of interest to declare.

Ethical issues: None

References

1. FAO. FAOSTAT. Crops and livestock products [Internet]. 2022 [cited 22 May 2024]. Available from: <http://www.fao.org/faostat/en/#data/QC>.
2. Di Vecchi-Staraz M, Laucou V, Bruno G, Lacombe T, Gerber S, Bourse T, et al. Low level of pollen-mediated gene flow from cultivated to wild grapevine: consequences for the evolution of the endangered subspecies *Vitis vinifera* L. subsp. *silvestris*. J Hered. 2009;100(1):66-75. <https://doi.org/10.1093/jhered/esn084>
3. Shurigin V, Alaylar B, Davranov K, Wirth S, et al. Diversity and biological activity of culturable endophytic bacteria associated with marigold (*Calendula officinalis* L.). AIMS Microbiol. 2021;7(3):336-53. <https://doi.org/10.3934/microbiol.2021021>
4. Shurigin V, Li L, Alaylar B, Egamberdieva D, et al. Plant beneficial traits of endophytic bacteria associated with fennel (*Foeniculum vulgare* Mill.). AIMS Microbiol. 2024;10(2):449-67. <https://doi.org/10.3934/microbiol.2024022>
5. Rijavec T, Lapanje A. Hydrogen cyanide in the rhizosphere: not suppressing plant pathogens, but rather regulating availability of phosphate. Front Microbiol. 2016;7. <https://doi.org/10.3389/fmicb.2016.01785>
6. Darriaut R, Lailheugue V, Masneuf-Pomarede I, Marguerit E, et al. Grapevine rootstock and soil microbiome interactions: Keys for a resilient viticulture. Horticulture Res. 2022;9. <https://doi.org/10.1093/hr/uhac019>
7. Rilling JI, Acuna JJ, Sadowsky MJ, Jorquera MA. Putative

- nitrogen-fixing bacteria associated with the rhizosphere and root endosphere of wheat plants grown in an andisol from southern Chile. *Front Microbiol.* 2018;9. <https://doi.org/10.3389/fmicb.2018.02710>
8. Baldan E, Nigris S, Romualdi C, D'Alessandro S, Clocchiatti A, et al. Beneficial bacteria isolated from grapevine inner tissues shape *Arabidopsis thaliana* roots. *Plos One.* 2015;10(10). <https://doi.org/10.1371/journal.pone.0140252>
 9. Ma H, Shurigin V, Jabborova D, dela Cruz JA, dela Cruz TE, et al. The integrated effect of microbial inoculants and biochar types on soil biological properties and plant growth of lettuce (*Lactuca sativa* L.). *Plants.* 2022;11(3):423. <https://doi.org/10.3390/plants11030423>
 10. Mardanov AV, Chizhevskaya EP, Lazarev AM, Rakitin AL, Beletsky AV, et al. Complete genome sequences of endophytic bacilli isolated from grapevine plants. *Microbiol Resour Announc.* 2019;8(48). <https://doi.org/10.1128/MRA.01265-19>
 11. Shurigin V, Alimov J, Davranov K, Gulyamova T, Egamberdieva D. The diversity of bacterial endophytes from *Iris pseudacorus* L. and their plant beneficial traits. *Curr Res Microb Sci.* 2022;3. <https://doi.org/10.1016/j.crmicr.2022.100133>
 12. Marasco R, Rolli E, Fusi M, Michoud G, Daffonchio D. Grapevine rootstocks shape underground bacterial microbiome and networking but not potential functionality. *Microbiome.* 2018;6:3. <https://doi.org/10.1186/s40168-017-0391-2>
 13. Gavande SS, Maurya A, Sharma S. Isolation and characterization of plant growth promoting rhizobacteria (PGPR) from rhizosphere of major crops grown in Marathwada region of Maharashtra, India. *Vegetos.* 2024;37:637-48. <https://doi.org/10.1007/s42535-023-00779-y>
 14. Agunbiade VF, Fadiji AE, Agbodjato NA, Babalola OO. Isolation and characterization of plant-growth-promoting, drought-tolerant rhizobacteria for improved maize productivity. *Plants.* 2024;13(10). <https://doi.org/10.3390/plants13101298>
 15. Karagoz K, Ates F, Karagoz H, Kotan R, Cakmakci R. Characterization of plant growth-promoting traits of bacteria isolated from the rhizosphere of grapevine grown in alkaline and acidic soils. *Eur J Soil Biol.* 2012;50:144-50. <https://doi.org/10.1016/j.ejsobi.2012.01.007>
 16. Labouyrie M, Ballabio C, Romero F, et al. Patterns in soil microbial diversity across Europe. *Nat Commun.* 2023;14:3311. <https://doi.org/10.1038/s41467-023-37937-4>
 17. Fouda A, Eid AM, Elsaied A, El-Belely EF, Barghoth MG, Azab E, et al. Plant growth promoting endophytic bacterial community inhabiting the leaves of *Pulicaria incisa* (Lam.) DC inherent to arid regions. *Plants.* 2021;10:76. <https://doi.org/10.3390/plants10010076>
 18. Zucol F, Ammann RA, Berger C, Aebi C, Altwegg M, et al. Real-time quantitative broad-range PCR assay for detection of the 16S rRNA gene followed by sequencing for species identification. *J Clin Microbiol.* 2006;44(8):2750-59. <https://doi.org/10.1128/JCM.00112-06>
 19. Saitou N, Nei M. The neighbor-joining method: A new method for reconstructing phylogenetic trees. *Mol Biol Evol.* 1987;4(4):406-25. <https://doi.org/10.1093/oxfordjournals.molbev.a040454>
 20. Felsenstein J. Confidence limits on phylogenies: An approach using the bootstrap. *Evolution.* 1985;39(4):783-91. <https://doi.org/10.2307/2408678>
 21. Tamura K, Nei M, Kumar S. Prospects for inferring very large phylogenies by using the neighbor-joining method. *Proc Natl Acad Sci USA.* 2004;101(30):11030-35. <https://doi.org/10.1073/pnas.0404206101>
 22. Kumar S, Stecher G, Li M, Knyaz C, Tamura K. MEGA X: Molecular evolutionary genetics analysis across computing platforms. *Mol Biol Evol.* 2018;35(6):1547-49. <https://doi.org/10.1093/molbev/msy096>
 23. Bashan Y, Holguin G, Lifshitz R. Isolation and characterization of plant growth-promoting rhizobacteria. In: Glick BR, Thompson JE, editors. *Methods in Plant Molecular Biology and Biotechnology.* USA, FL, Boca Raton: CRC Press; 1993:331-45.
 24. Sarwar M, Kremer RJ. Determination of bacterially derived auxins using a microplate method. *Lett Appl Microbiol.* 1995;20:282-85. <https://doi.org/10.1111/j.1472-765X.1995.tb00446.x>
 25. Kuklinsky-Sobral J, Araújo WL, Mendes R, Geraldi IO, et al. Isolation and characterization of soybean-associated bacteria and their potential for plant growth promotion. *Environ Microbiol.* 2004;6:1244-51. <https://doi.org/10.1111/j.1462-2920.2004.00658.x>
 26. Mehta S, Nautiyal CS. An efficient method for qualitative screening of phosphate-solubilizing bacteria. *Curr Microbiol.* 2001;43:51-56. <https://doi.org/10.1007/s002840010259>
 27. Egamberdieva D, Kucharova Z, Davranov K, Berg G, Makarova N, et al. Bacteria able to control foot and root rot and to promote growth of cucumber in salinated soils. *Biol Fertil Soils.* 2011;47:197-205. <https://doi.org/10.1007/s00374-010-0523-3>
 28. Schwyn B, Neilands JB. Universal chemical assay for the detection and determination of siderophores. *Anal Biochem.* 1987;160:45-46. [https://doi.org/10.1016/0003-2697\(87\)90612-9](https://doi.org/10.1016/0003-2697(87)90612-9)
 29. Logan NA, De Vos P. Genus I. *Bacillus* Cohn 1872, 174AL. In: De Vos P, Garrity GM, Jones D, Krieg NR, Ludwig W, Rainey FA, Schleifer KH, Whitman WB, editors. *Bergey's Manual of Systematic Bacteriology*, 2nd Edition. New York: Springer; 2009;21-128.
 30. Egamberdieva D, Shurigin V, Gopalakrishnan S, Ram S. Microbial strategies for the improvement of legume production in hostile environments. In: Azooz MM, Ahmad P, editors. *Legumes Under Environmental Stress: Yield, Improvement and Adaptations.* UK: John Wiley and Sons Ltd; 2015:133-44. <https://doi.org/10.1002/9781118917091.ch9>
 31. Aasfar A, Meftah Kadmiri I, Azaroual SE, Lemriss S, et al. Agronomic advantage of bacterial biological nitrogen fixation on wheat plant growth under contrasting nitrogen and phosphorus regimes. *Front Plant Sci.* 2024;15. <https://doi.org/10.3389/fpls.2024.1388775>
 32. Shurigin V, Alikulov B, Davranov K, Ismailov Z. Bacterial endophytes from halophyte black saxaul (*Haloxylon aphyllum* Minkw.) and their plant growth-promoting properties. *J Appl Biol Biotech.* 2022;10(1):45-53. <https://doi.org/10.7324/JABB.2021.100106>
 33. Kondracka A, Rychter AM. The role of P_i recycling processes during photosynthesis in phosphate-deficient bean plants. *J Exp Bot.* 1997;48:1461-68. <https://doi.org/10.1093/jxb/48.7.1461>
 34. Ma Z, Bielenberg DG, Brown KM, Lynch JP. Regulation of root hair density by phosphorus availability in *Arabidopsis thaliana*. *Plant Cell Environ.* 2001;24:459-67. <https://doi.org/10.1111/pce.12059>
 35. Vance CP, Uhde-Stone C, Allan DL. Phosphorus acquisition and use: critical adaptations by plants for securing a nonrenewable resource. *New Phytol.* 2003;157(3):423-47. <https://doi.org/10.1046/j.1469-8137.2003.00695.x>
 36. Egamberdieva D, Ma H, Shurigin V, Alimov J, et al. Biochar additions alter the abundance of p-cycling-related bacteria in the rhizosphere soil of *Portulaca oleracea* L. under salt stress. *Soil Syst.* 2022;6:64. <https://doi.org/10.3390/soilsystems6030064>
 37. Chandra S, Askari K, Kumari M. Optimization of indole acetic acid production by isolated bacteria from *Stevia rebaudiana* rhizosphere and its effects on plant growth. *J Genet Eng Biotechnol.* 2018;16(2):581-86. <https://doi.org/10.1016/j.jgeb.2018.09.001>

38. Fierro-Coronado RA, Quiroz-Figueroa FR, Garcia-Perez LM, Ramirez-Chavez E, et al. IAA-producing rhizobacteria from chickpea (*Cicer arietinum* L.) induce changes in root architecture and increase root biomass. *Can J Microbiol*. 2014;60(10):639-48. <https://doi.org/10.1139/cjm-2014-0399>
39. Jaiswal A, Das K, Koli DK, Pabbi S. Characterization of cyanobacteria for IAA and siderophore production and their effect on rice seed germination. *Int J Curr Microbiol App Sci*. 2018;5:212-22.
40. Ramos Solano B, Maicas JB, Pereyra De La Iglesia MT, Domenech J, Gutierrez Manero FJ. Systemic disease protection elicited by plant growth promoting rhizobacteria strains: relationship between metabolic responses, systemic disease protection and biotic elicitors. *Am Phytopath Soc*. 2008;98(4):451-57. <https://doi.org/10.1094/PHYTO-98-4-0451>
41. Gu S, Wei Z, Shao Z, Friman VP, Cao K, Yang T, et al. Competition for iron drives phytopathogen control by natural rhizosphere microbiomes. *Nat Microbiol*. 2020;5:1002-10. <https://doi.org/10.1038/s41564-020-0719-8>
42. Khan A, Singh P, Srivastava A. Synthesis, nature and utility of universal iron chelator - Siderophore: a review. *Microbiol Res*. 2018;212-213:103-11. <https://doi.org/10.1016/j.micres.2017.10.012>
43. Kramer J, Ozkaya O, Kummerli R. Bacterial siderophores in community and host interactions. *Nat Rev Microbiol*. 2020;18:152-63. <https://doi.org/10.1038/s41579-019-0284-4>
44. Pan YJ, Liu L, Lin YC, Zu YG, et al. Ethylene antagonizes salt-induced growth retardation and cell death process via transcriptional controlling of ethylene-, BAG- and senescence-associated genes in *Arabidopsis*. *Front Plant Sci*. 2016;7:696. <https://doi.org/10.3389/fpls.2016.00696>
45. Subhan D, Muhammad Z, Fauzia M, Mubshar H. ACC-deaminase producing plant growth promoting rhizobacteria and biochar mitigate adverse effects of drought stress on maize growth. *Plos One*. 2020;16. <https://doi.org/10.1371/journal.pone.0250286>
46. Singh RP, Shelke GM, Kumar A, Jha PN. Biochemistry and genetics of ACC deaminase: a weapon to “stress ethylene” produced in plants. *Front Microbiol*. 2015;6:937. <https://doi.org/10.3389/fmicb.2015.00937>
47. Tiwari G, Duraivadivel P, Sharma S, Hariprasad P. 1-Aminocyclopropane-1-carboxylic acid deaminase producing beneficial rhizobacteria ameliorate the biomass characters of *Panicum maximum* Jacq. by mitigating drought and salt stress. *Sci Rep*. 2018;8. <https://doi.org/10.1038/s41598-018-35565-3>