



REVIEW ARTICLE

Molecular characterization for identifying the evolutionary lineage of Indonesia strawberry (*Fragaria* spp.) using DNA barcode markers: A review and current research on breeding strategies

Ganies Riza Aristya¹, Muhammad Ariq Alfito¹, Muhammad Fatih¹, Muhammad Sena Budiman¹, Rina Sri Kasiamdari² & Ani Widiastuti^{3*}

¹Laboratory of Genetics and Breeding, Faculty of Biology, Universitas Gadjah Mada 55281, Indonesia

²Laboratory of Plant Systematics, Faculty of Biology, Universitas Gadjah Mada 55281, Indonesia

³Laboratory of Pests and Plant Diseases, Department of Plant Protection, Faculty of Agriculture, Universitas Gadjah Mada 55281, Indonesia

*Correspondence email - ganies_riza@ugm.ac.id

Received: 05 February 2024; Accepted: 28 December 2024; Available online: Version 1.0: 19 July 2025

Cite this article: Ganies RA, Muhammad AA, Muhammad F, Muhammad SB, Rina SK, Ani W. Molecular characterization for identifying the evolutionary lineage of Indonesia strawberry (*Fragaria* spp.) using DNA barcode markers: A review and current research on breeding strategies. Plant Science Today (Early Access). <https://doi.org/10.14719/pst.3345>

Abstract

Fragaria spp. (Strawberry) is one of the most popular plant species, with a high economic and nutritional value. It also plays a key role in evolutionary biology, particularly due to its widespread cultivation in Indonesia, where diverse cultivars are imported from abroad. Understanding the evolutionary paths of these cultivars is essential for grasping their genetic diversity. This review aims to explore the evolutionary dynamics of strawberry species found in Indonesia using a DNA barcoding approach. DNA barcoding is a powerful tool for uncovering the complex evolutionary histories of species and revealing their phylogenetic relationships. In this comprehensive review, we discuss various cultivated strawberry species present in Indonesia and examine the evolutionary forces that have driven their diversification. By analyzing genetic evidence, we seek a deeper understanding of the evolutionary processes behind these cultivars. However, challenges remain, particularly for *Fragaria* species, which face increasingly threats from the introduction and spread of “alien species”. Addressing these challenges requires a holistic approach, that combines evolutionary biology principles with conservation strategies to protect native biodiversity while maximizing agricultural productivity. By enhancing our understanding of the evolutionary patterns and genetic evidence associated with cultivated strawberries in Indonesia, this review not only advances scientific knowledge but also provides a comparative phylogenetic tree of strawberry species in Indonesia. This serves as a foundation for informed conservation and breeding efforts in the constantly changing agricultural landscape.

Keywords: DNA barcoding; *Fragaria* spp.; molecular genetics; strawberry

Introduction

Strawberries (*Fragaria* spp.) are plants that produce fruits with high economic value and nutritional content (1). According to data from the Central Statistics Agency (2023), strawberry production in Indonesia reached 28895 t in 2022, marking a significant increase compared to the 9860 t produced in 2021 (2). The successful cultivation of strawberries in Indonesia is the result of cultivation from several countries, with 23 species and cultivars currently grown in the highlands of Indonesia (3).

Indonesia, despite being a subtropical country, has a high potential for the development of fruit commodities, including strawberries. Strawberry cultivation in Indonesia is primarily limited to highland areas, following the natural habitat of strawberries, which thrive in cold regions (1, 4). The required temperature for strawberries to grow ranges from 20 -25 °C. However, recent climate changes have increased air temperatures, necessitating strawberry adaptation (5). The

adaptation of strawberries in Indonesia enables them to thrive at higher temperatures than those of their native habitat (6).

The cultivated strawberries in Indonesia are cultivars from abroad, but there is currently no research discussing the evolutionary forces that naturally occur in strawberries in Indonesia. Cultivated strawberries tend to undergo evolution due to human breeding. Previous study illustrates that a crossbreeding programme between *Fragaria* × *ananassa* cultivars from California and Hoybride, resulting in a new cultivar exhibiting resistance to fruit rot, high productivity and a sweet fruit flavour (7).

Generally, the relationship between genotype and phenotype reflects how genetic information is inherited from DNA and chromosomes and how this information produces observable physical characteristics in an individual (8). The interaction between genotype and phenotype in strawberries

can affect heritability and the relationship between the quality and quantity of its traits (8). Therefore, the outcomes of hybrid strawberries and new strawberry species can have different genotypes, resulting in phenotypic variations.

This research aims to understand and identify the evolutionary relationships among various strawberry species worldwide, especially in Indonesia, using DNA barcoding techniques.

Biodiversity of strawberry in Indonesia

The genus *Fragaria*, comprising approximately 23 species worldwide (Table 1), exhibits the same basic life history: they are insect-pollinated, low-growing, herbaceous perennials capable of clonal propagation and produce animal-dispersed fleshy accessory fruits (9). However it can also have fascinating diversity in both ploidy levels and morphological traits. Most commonly, species such as wild strawberry *Fragaria vesca* are diploid ($2n = 14$), while others, like *Fragaria* \times *ananassa* (garden strawberry), are octoploid ($2n = 56$) (10). This variation in ploidy significantly influences their morphological characteristics, including leaf shape, flower size and fruit characteristics. Typically, diploid species bear smaller, more delicate flowers and fruits, while polyploidy species tends to produce larger, more robust fruits with increased sweetness and varying colors (9).

The floral and vegetative morphology of *Fragaria* species is relatively uniform. Leaves are usually evergreen (with *F. iinumae* being an exception as it is deciduous) and trifoliate. While 5 leaflets are common among some Chinese species, the

occurrence of 4-5 leaflets is rare and has been observed in *F. virginiana* and *F. cascadiensis* (11). The flowers are always actinomorphic, predominantly white (sometimes tinged with pink) and usually possess 5 petals (9).

Indonesia boasts high biodiversity, also hosts a variety of strawberry cultivars. The country currently lacks a National Biodiversity Index (IBI) to monitor biodiversity trends comprehensively. As such, there is a pressing need to prioritize conservation, identification and systematic documentation, particularly in relation to their potential inclusion in the Indonesian Biodiversity Index (12). This effort should encompass cultivars, hybrids and native species. There are several compelling reasons to support the conservation and identification of strawberry species in Indonesia, including the preservation of biodiversity, improved agricultural productivity, enhanced fruit quality and increased economic value of strawberries at the national level (13). Taking proactive steps to conserve and identify strawberries in accordance with the IBI will contribute both to biodiversity conservation and to the economic and agricultural development of the country (12, 14).

Based on Table 2, the majority of identified strawberries in Indonesia belong to the hybrid species *F* \times *ananassa*, which has been imported for food production purposes (3). These hybrid species have been distributed in several locations, namely West Java, Central Java, East Java, North Sumatra and Bali (3, 7, 16 -18). The distribution of these cultivars is closely linked to environmental suitability, as strawberries generally thrive in highland areas at elevations of around 1000 m above sea level (3).

Table 1. *Fragaria* species and their ploidy level, mating system and geographic distribution (9)

Taxon	Ploidy	Mating system	Geographic distribution
Vesca clade			
<i>F. \times ananassa</i> Duchesne subsp. <i>ananassa</i>	8x	Subdioecious (modern cultivars: hermaphroditic)	Cultivated around the world
<i>F. \times ananassa</i> Duchesne subsp. <i>cuneifolia</i> (Nutt. ex Howell) Staudt	8x	Subdioecious	NW N. America
<i>F. bucharica</i> Losinsk.	2x	Hermaphrodite SI	W Himalayas
<i>F. cascadiensis</i> Hummer	10x	Subdioecious	Oregon, USA
<i>F. chiloensis</i> (L.) Duchesne	8x	Subdioecious	Alaska-California; Hawaii; Chile, Argentina
<i>F. iturupensis</i> Staudt	8x, 10x	Subdioecious	Iturup Island
<i>F. mandshurica</i> Staudt	2x	Hermaphrodite SI	NF Asua
<i>F. moschata</i> Duchesne	6x	Dioecious	W Eurasia
<i>F. orientalis</i> Losinsk.	4x	Dioecious	NE Asia
<i>F. vesca</i> L. subsp. <i>americana</i> (Porter) Staudt	2x	Hermaphrodite SC	NE N. America
<i>F. vesca</i> L. subsp. <i>bracteata</i> (A. Heller) Staudt	2x	Gynodioecious or Hermaphrodite SC	W N. America
<i>F. vesca</i> L. subsp. <i>californica</i> (Cham. & Schldl.) Staudt	2x	Gynodioecious or Hermaphrodite SC	SW N. America
<i>F. vesca</i> L. subsp. <i>vesca</i> L.	2x	Hermaphrodite SC	W Eurasia
<i>F. virginiana</i> Duchesne	8x	Subdioecious	N. America
China clade			
<i>F. chinensis</i> Losinsk.	2x	Hermaphrodite SI	China
<i>F. corymbosa</i> Losinsk.	4x	Dioecious	China
<i>F. daltoniana</i> J. Gay	2x	Hermaphrodite SC	Nepal, adjacent China
<i>F. gracilis</i> Losinsk.	4x	Dioecious	China
<i>F. moupinensis</i> (Franch.) Cardot	4x	Dioecious	China
<i>F. nipponica</i> Makino	2x	Hermaphrodite SI	Japan
<i>F. nubicola</i> Lindl.	2x	Hermaphrodite SI	Himalayas
<i>F. pentaphylla</i> Losinsk	2x	Hermaphrodite SI	China
<i>F. tibetica</i> Staudt & Dickore	4x	Dioecious	China
Unresolved phylogenetic position			
<i>F. hayatai</i> Makino	2x	Hermaphrodite SC?	Taiwan
<i>F. iinumae</i> Makino	2x	Hermaphrodite SC	Japan
<i>F. nilgerrensis</i> Schldtl. ex J. Gay	2x	Hermaphrodite SC	SE Asia
<i>F. viridis</i> Duchesne	2x	Hermaphrodite SI	W Eurasia

Table 2. Strawberry species cultivated in Indonesia

No	Cultivars	Locations	Ref.
1	<i>Fragaria</i> × <i>ananassa</i> cv. Camarosa	-	
2	<i>Fragaria</i> × <i>ananassa</i> cv. Sunset	-	
3	<i>Fragaria</i> × <i>ananassa</i> cv. Selva	Purbalingga, Karanganyar, Magelang (Center Java)	
4	<i>Fragaria</i> × <i>ananassa</i> cv. Carmine	-	
5	<i>Fragaria</i> × <i>ananassa</i> cv. Pajero	-	
6	<i>Fragaria</i> × <i>ananassa</i> cv. Sweetcharlie	Ciwidey, Garut, Lembang (Center Java) & Batu, Pasuruan, Bondowoso, Magetan (East Java) & Bedugul (Bali) & Kecamatan Berastagi (North Sumatera)	
7	<i>Fragaria</i> × <i>ananassa</i> cv. Festival	-	
8	<i>Fragaria</i> × <i>ananassa</i> cv. Earlybrite	Ciwidey, Garut, Lembang (West Java)	
9	<i>Fragaria</i> × <i>ananassa</i> cv. Cal Giant	-	
10	<i>Fragaria</i> × <i>ananassa</i> cv. Aromas	-	
11	<i>Fragaria</i> × <i>ananassa</i> cv. Diamante	-	
12	<i>Fragaria</i> × <i>ananassa</i> cv. Giavonta	-	
13	<i>Fragaria</i> × <i>ananassa</i> cv. Terry	-	
14	<i>Fragaria</i> × <i>ananassa</i> cv. Adina	-	
15	<i>Fragaria</i> × <i>ananassa</i> cv. Collma	-	(3)
16	<i>Fragaria</i> × <i>ananassa</i> cv. Whitney	-	
17	<i>Fragaria</i> × <i>ananassa</i> cv. Chandler	Pasuruan, Bondowoso, Magetan (East Java)	
18	<i>Fragaria</i> × <i>ananassa</i> cv. Ozogrande	Purbalingga, Karanganyar, Magelang (Center Java) & Kecamatan Berastagi (North Sumatera)	
19	<i>Fragaria</i> × <i>ananassa</i> cv. Rosalinda	Batu (East Java) & Bedugul (Bali)	
20	<i>Fragaria</i> × <i>ananassa</i> cv. Kanaka Peak 794-15	-	
21	<i>Fragaria</i> × <i>ananassa</i> cv. Kanaka Peak 770-506	-	
22	<i>Fragaria</i> × <i>ananassa</i> cv. Winter down	-	
23	<i>Fragaria</i> × <i>ananassa</i> cv. Icigo	-	
24	<i>Fragaria</i> × <i>ananassa</i> cv. California	Ciwidey, Garut, Lembang (West Java)	
25	<i>Fragaria</i> × <i>ananassa</i> cv. Holland	Ciwidey, Garut, Lembang (West Java) & Pasuruan, Bondowoso, Magetan (East Java)	
26	<i>Fragaria</i> × <i>ananassa</i> cv. Anna	Purbalingga, Karanganyar, Magelang (Center Java)	
27	<i>Fragaria</i> × <i>ananassa</i> cv. Tristar	Purbalingga, Karanganyar, Magelang (Center Java)	
28	<i>Fragaria</i> × <i>ananassa</i> cv. Quantum	Purbalingga, Karanganyar, Magelang (Center Java)	
29	<i>Fragaria</i> × <i>ananassa</i> cv. Berastagi	Berastagi (North Sumatera)	
30	<i>Fragaria</i> × <i>ananassa</i> cv. Dorit	Berastagi (North Sumatera)	
31	<i>Fragaria</i> × <i>ananassa</i> cv. Hoybride	Batu (East Java)	(7)
32	<i>Fragaria</i> × <i>ananassa</i> cv. Santung	Malang (East Java)	(15)
33	<i>Fragaria</i> × <i>ananassa</i> cv. Aerut	Malang (East Java)	
34	<i>Fragaria</i> sp.	Pekanbaru (Riau)	(16)
35	<i>Fragaria</i> sp.	Buleleng (Bali)	(17)




Fig. 1. Types of growth of nine cultivars of strawberries. (A) California, (B) Rosa Linda, (C) Berastagi, (D) Festival, (E) Santung, (F) Holland, (G) Sweet Charlie, (H) Aerut, (I) Earlibrite (Upright: A and H; Intermediate: B-G and I) (15).

Fragaria × *ananassa* originated from a natural hybridization event between *Fragaria chiloensis* and *Fragaria virginiana*, which occurred in Europe during the 18th century. This hybridization resulted in the mixing of genetic material from both species, yielding new offspring with unique morphological characteristics and genetic sequences (9). Over time, various cultivars of *F. × ananassa* have been developed through selective breeding to enhance desired traits such as fruit size, flavor, yield, disease resistance and adaptability to different growth conditions. The breeding process involves crossing various cultivars and selecting individuals with favourable traits for further propagation (18). These hybrid cultivars often exhibit morphological and physiological adaptations that contribute to increase disease resistance and faster growth (19).

Several cultivars of strawberries found in Indonesia, namely California, Rosa Linda, Berastagi, Festival, Santung, Holland, Sweet Charlie, Aerut and Earlibrite, exhibit differences in morphological characteristics. Among these cultivars, Sweet Charlie and Aerut have an upright growth type, whereas California, Rosa Linda, Berastagi, Festival, Santung, Holland and Earlibrite have an intermediate growth type. Furthermore, concerning fruit shape, 4 dominant types

are observed: ovoid (Californica, Santung and Holland); cylindrical (Sweet Charlie); conical (Rosa Linda, Berastagi and Festival) and cordate (Aerut and Early Brite) (Fig. 2)(15).

The evolutionary history of the wild genus *Fragaria* is distinguished by complex interactions with genetic diversity, hybridization, varying ploidy levels and wide geographic distribution. Based on Fig. 3, the genus *Fragaria* comprises a number of species exhibiting diverse ploidy levels, including diploid (2x), tetraploid (4x), hexaploid (6x) and octaploid (8x). The cultivated strawberry, *F. × ananassa*, is the result of a hybridization event between the wild species *F. virginiana*, native to North America and

Frchiloensis, from South America. This hybridization led to the development of a novel species that has become the dominant strawberry crop globally. Notably, the diploid species *F. vesca* served as a primary root in the establishment of this evolutionary lineage. Recent genomic studies have further elucidated the evolutionary dynamics within *Fragaria*, revealing multiple diploid progenitor species and highlighting the influence of environmental factors in shaping genetic diversity (20, 21).

Based on Fig. 4, *F.* species selected for analysis were collected from diverse regions, including India, China, Japan, East Asia, Southeast Asia, America, Europe and the Kuril Islands. Phylogenetic analysis of *Fragaria* cultivars and species worldwide indicates that *F. × ananassa* is most closely related to *F. virginiana*. This relationship exists because *F. ananassa* is a hybrid of *F. virginiana* and *F. chiloensis* (22, 23). This phylogenetic analysis is further supported in Fig. 5, which examines the phylogeny of *Fragaria* found in Pakistan and highlights the crucial role of gene inheritance in *Fragaria* evolution.

The incorporation of outgroups in phylogenetic analysis is crucial, as it establishes a baseline for comparison, clarifies evolutionary relationships, minimizes ambiguity and enhances analytical precision. In this study, *Potentilla fruticosa* was selected as an outgroup based on its morphological and ecological similarities to *Fragaria* species (25). Moreover, the distribution of these 2 plants overlaps, making them suitable for evolutionary comparison. This is supported by a previous study, which revealed *Fragaria*'s close relationship with the *Potentilla* clade, one of which includes *P. fruticosa* (22).

Essentially, species are defined as individual entities or groups sharing consistent characteristics. Meanwhile, cultivars are varieties of plant species that have been selected or bred for specific characteristics and cultivated to exhibit specific traits (26). In other words, species are groups of organisms with similar characteristics, while cultivars are varieties resulting from selective breeding to obtain specific traits.




Fig. 2. Fruit shapes of nine cultivars of strawberries: ovoid (A, E, F), cylindrical (G), conical (B, C, D) and cordate (H, I). The cultivars are (A) Californica, (B) Rosa Linda, (C) Berastagi, (D) Festival, (E) Santung, (F) Holland, (G) Sweet Charlie, (H) Aerut and (I) Earlibrite (15).




Fig. 3. Phylogenetic analyses of wild type strawberry based on number of chromosomes (20).




Fig. 4. Phylogenetic analysis of *Fragaria* based on maximum parsimony worldwide (22).




Fig. 5. Phylogenetic analysis of *Fragaria* based on ITS2 gene by maximum Likelihood in Pakistan (27).

However, due to the government's limited attention to the biodiversity of strawberry plants in Indonesia, there is an urgent need for a conservation approach to protect, document and collect them using DNA barcoding. DNA barcoding is an essential tool in the conservation context, as it enables rapid and accurate species identification, aiding in the recognition of various species, whether invasive or endangered and supporting their management and protection. Additionally, DNA barcoding enables the measurement of phylogenetic diversity, offering a more detailed and descriptive picture of biodiversity compared to alternative indices such as species richness and abundance (28).

Genetic conservation aims to preserve the genetic material of organisms to minimize the risk of extinction and the threat of population decline that may occur through genetic and evolutionary processes (29, 30). Thus, DNA barcoding is crucial in estimating species richness and establishing conservation priorities, especially in areas with minimal information about their biodiversity. Moreover, this method can help identify small areas that are important for conservation efforts, considering locations that have served as natural refuges for certain species in the past and areas that have recently become home to new lineages (28). Conservation efforts through genetic approaches have been undertaken in several plants, especially strawberries, such as *Fragaria* spp. (31) and wild *Fragaria* (32).

The analysis of the phylogenetic tree of *Fragaria* species was conducted using three genes, including ITS2, *rbcL* and *matK*, employing both Maximum Likelihood (ML) and Bayesian Inference (BI) methods (Fig. 6-8). In general, the results from both methods exhibited high similarity; therefore, only the phylogenetic trees constructed using the ML method are presented, supported by statistical values from both ML and BI analyses.

Based on Fig. 6, the results of the phylogenetic analysis of *Fragaria* species concerning the ITS2 gene can be observed using ML and BI methods. These results show similarities to previous studies (22) (Fig. 4). Overall, the closest relationship is between *F. virginiana*, *F. × ananassa* and *F. chiloensis*. Additionally, among the three outgroups used, the results suggest that *Fragaria* species are more closely related to *Spyridium buxifolium* and *Stenanthemum centrale* than *Spiraea humilis*.

Fig. 7 presents the results of the phylogenetic analysis based on the *rbcL* gene utilizing both ML and BI

methods. The results indicate that the most *Fragaria* species share a similar evolutionary relationship. However, *F. nilgerrensis* is situated in a distinct lineage compared to other *Fragaria* species. Additionally, among the 3 outgroups, *Rosa multiflora* and *Potentilla angelica* exhibit a closer relationship than *Geum henryi*.

Similarly, Fig. 8 illustrates the results of the phylogenetic analysis of *Fragaria* species using the *matK* gene with ML and BI methods. These results exhibit similarities to the phylogenetic analysis results of *Fragaria* species with the *rbcL* gene (Fig. 7). The analysis demonstrates that various *Fragaria* species, such as *F. virginiana*, *Fragaria × ananassa*, *Fragaria vesca*, *Fragaria orientalis*, *F. chiloensis* and *F. sp.*, are in the same lineage. However, *F. nilgerrensis* once again forms a distinct branch from the other *F. species*. Additionally, among the 3 outgroups utilized, the results suggest that *Fragaria* species have a closer relationship to *Potentilla pulchella* and *Rubus idaeus* than *Drymocallis arguta*.




Fig. 6. Phylogenetic analysis *Fragaria* using ITS2 gene based on maximum-likelihood (ML) and Bayesian Inference (BI).




Fig. 7. Phylogenetic analysis *Fragaria* using *rbcl* gene based on Maximum-Likelihood (ML) and Bayesian Inference (BI).




Fig. 8. Phylogenetic analysis *Fragaria* using *matK* gene based on Maximum-Likelihood (ML) and Bayesian Inference (BI).

Correlation between molecular genetics analysis and molecular systematics

Molecular genetics is one of the latest technologies that can be used to reveal the identity of a new species by leveraging the genetic information of an organism and comparing its DNA or RNA sequences with those of known organisms (33). Strawberries, which belong to the Rosaceae family, have undergone significant variations in taxonomic classification over the last few decades, with various classification methods

proposed (34). The complexity of taxonomy and the limitations of morphological characteristics within this family pose difficulties in identifying species based solely on morphological taxonomy. Therefore, Rosaceae is an ideal subject for identification using DNA barcoding methods, as this family presents taxonomic barriers that complicate the separation of closely related species within the family using conventional approaches (35).

DNA barcoding is a highly useful method for taxonomic identification, utilizing short, universally conserved DNA segments that have sufficient sequence variations to distinguish species (36, 37). The primary goal of the DNA barcoding approach is to authenticate known species by matching their DNA sequences with references in the barcode sequence library and to facilitate the recognition of new species (36, 38). For example, the mitochondrial cytochrome c oxidase 1 (cox1) gene sequence has been used as a universal barcode in identifying several groups of animals (39). Meanwhile, the combination of the *matK* + *rbcL* genes has been proposed as the nuclear DNA barcode for plant identification by the Consortium for the Barcode of Life (CBOL) Plant Working Group (PWG), which also suggests using ITS2 (internal transcribed spacer 2) as an additional identifier (35). The study DNA barcodes such as ITS, ITS2, *matK*, *rbcL*, *rbcLA*, *rbcLC*, *Ycf3* and *trnV* were used to characterize strawberries from Mansehra, Pakistan and determine their relationships and phylogenetic tree (33).

The % of similarity between species, derived from genetic analysis, provides insights into their evolutionary history by revealing how closely related they are. A high % of similarity indicates that 2 species share a more recent common ancestor, reflecting a closer evolutionary relationship. Conversely, a lower % suggests a more distant evolutionary connection (40). Based on Table 3 the % of similarity between the *Fragaria* species indicates a high % similarity suggesting that these species share a more recent common ancestor, indicating closer evolutionary ties. Based on Table 3, the percentage of similarity among *Fragaria* species indicates a high % of similarity, suggesting that these species share a recent common ancestor, indicating closer evolutionary ties. Specifically, *F. chiloensis*, *F. virginiana* and *F. × ananassa* in particular exhibit 99.9% to 100% similarity. This finding is consistent with their evolutionary history, as *F. × ananassa* is a cross between *F. chiloensis* and *F. virginiana* (22).

Before molecular genetics was applied as a species identification technique, several other identification techniques existed, such as morphology, anatomy, physiology and biochemistry-based identification. Each of these identification methods has its advantages and disadvantages. The advantage of molecular taxonomy is its ability to identify species that are difficult to identify by other means, such as species with similar morphologies or species that are challenging to differentiate physiologically or biochemically (33). Another major advantage lies in the fact that DNA sequences are directly inherited and are not subject to developmental or environmental influences that may obscure morphological traits. Consequently, molecular taxonomy provides a more stable and objective basis for species identification. Moreover, it allows for the identification of future evolutionary lineages based on their genetic sequences (33).

However, molecular taxonomy is not without its limitations. Its application still relies on species definitions provided by morphological taxonomy, making the 2 methods interdependent. Additionally, the accuracy of molecular identification may be compromised by potential contamination, which can distort the results of genetic analyses (41).

Table 3. Percent similarity of *Fragaria* species using DNA barcoding

Taxon	Percent Identity (%)	Accession
<i>Fragaria × ananassa</i>	100	OP723290
<i>Fragaria virginiana</i>	99.99	NC_019602
<i>Fragaria moupinensis</i>	99.41	MZ702805
<i>Fragaria tibetica</i>	99.41	MZ702810
<i>Fragaria gracilis</i>	99.41	NC_062837
<i>Fragaria chiloensis</i>	100	MW537844
<i>Fragaria tibetica</i>	99.41	NC_062835
<i>Fragaria vesca</i> subsp. <i>vesca</i>	99.77	MZ851769
<i>Fragaria mandshurica</i>	99.80	MZ851758
<i>Fragaria pentaphylla</i>	99.51	NC_034347
<i>Fragaria chinensis</i>	99.51	MZ702808
<i>Fragaria nubicola</i>	99.49	MW537841
<i>Fragaria corymbosa</i>	99.48	MZ851751
<i>Fragaria nilgerrensis</i>	99.49	MZ702804
<i>Fragaria nipponica</i>	99.47	MW537843
<i>Fragaria daltoniana</i>	99.39	MZ851755
<i>Fragaria iinumae</i>	99.39	MW537854
<i>Fragaria orientalis</i>	99.72	MW537853


The development of DNA barcoding as an accurate identification tool

DNA barcoding is used because it can identify species quickly and inexpensively (42). It is estimated that morphological analysis incurs higher costs and takes several months of fieldwork compared to species sequencing (43). However, there are challenges in applying DNA barcoding to plants because the genes used as phylogenetic markers have very limited variations, making them insufficient to differentiate species. Therefore, using techniques based on a chloroplast region is highly recommended (44).

Several genes commonly used for DNA barcoding include *rbcL*, *matK*, *cox1* and 16S (36, 45). The selection of appropriate genes is based on several factors: the marker should exhibit sufficient genetic variability and divergence at the species level, possess conserved regions suitable for the development of universal PCR primers for taxonomic applications and have a relatively short sequence length to facilitate efficient DNA extraction and amplification (46). Nonetheless, gene selection must be contextual, as certain genes are more suitable for specific taxonomic groups. For instance, the Internal Transcribed Spacer (ITS) region is commonly used for fungi because it is one of the most frequently sequenced markers and is routinely applied in systematics, phylogenetics and species identification (47). Therefore, gene selection should be guided by the research objectives and the biological characteristics of the species under investigation. The development of DNA barcoding for the identification of strawberry species involves the use of specific DNA sequences of genes, such as chloroplast DNA (Table. 4). Chloroplast partial (cp) has a genome that contains information for taxonomic and phylogenetic purposes, making it often used in plant identification research, including strawberries (31). In Fig. 9, *Potentilla fruticosa* and *Drymocalis saviczii* are used as outgroups with

Table 4. Function and characterization DNA barcode (8)

Genes	Genome	Organism	Functions	Characterization
<i>matK</i>	Chloroplast	Plant	Intra and Interspecies Taxonomy	Small genome in size, middle in variability and middle in mutation
<i>rbcl</i>			Interspecies	Small genome in size, the mutation is lower than <i>matK</i> , <i>ITS</i> and <i>rbcl</i>
<i>ITS1-rDNA</i>	Chromosomal		Intra and Interspecies Taxonomy	Degree of variation of genome is higher than <i>18S</i> , <i>matK</i> and <i>rbcl</i>
<i>ITS2-rDNA</i>				
<i>18S-rDNA</i>	Mitochondria	Animals	Intraspecies	Conserved gene
<i>COI-barcode</i>				conserved, stable, fast evolutions

**Fig. 9.** Phylogenetic analysis of *Fragaria* based on partial chloroplast sequences (31).

partial chloroplast genome sequences from 34 *Fragaria* accessions. Additionally, there is research focused on molecular phylogenetic analysis in the *Fragaria* genus based on whole chloroplast genome sequences (32). In Fig. 10 the outgroups used include *Rosa multiflora*, *Potentilla stolonifera* and *Malus hupehensis*, with whole chloroplast genomes from 20 *Fragaria* samples.

Opportunities and challenges

Difficulties in identifying new *Fragaria* species in Indonesia are often encountered, particularly due to the presence of numerous alien species (48). These alien species may be introduced either intentionally or unintentionally (48, 49). In Indonesia and other agronomically focused countries, such introductions often occur for cultivation experiments or through botanical garden collections (48). As a result, the presence of alien species reduces the genetic variation within *Fragaria*, frequently leading to the dominance of these introduced species (50). This situation is further exacerbated by limitations in DNA barcoding analysis, which complicates accurate identification (51). DNA may be degraded or lost during the analysis process, leading to partial or fragmented genetic data (52). Additionally, genetic sequences available in the National Center for Biotechnology Information (NCBI) database are often unpublished or lack sufficient detail, with

some studies failing to specify the genes associated with particular species (52). As a result, *Fragaria* identification becomes inaccurate or unsuccessful, as seen in a study, which failed to identify *Fragaria* species due to limited psbA-trnH genes and the use of other genes with limited chloroplast DNA (53).

The cultivation of alien species in close proximity to natural habitats also increases the potential for gene flow (54). In Central Europe, *F. x ananassa* has emerged in areas near abandoned strawberry gardens. However, this species has not yet been recorded in the NCBI GenBank database (55). While such occurrences may pose a risk to native species, the impact remains limited due to the rarity of hybridization events and the presence of reproductive barriers that restrict gene flow between cultivated and wild species (49, 55). It is still unclear whether pollinators significantly contribute to natural hybridization (55). Nonetheless, the possibility of hybridization affecting *Fragaria* diversity in Indonesia highlights the need for further research into species identification.

Beyond the challenge of distinguishing alien from native species, the genetic complexity of *F.* also presents difficulties due to its monogenic and polygenic characteristics (57, 58). In polygenic species, DNA barcoding is a time-intensive process, requiring the discrimination of multiple barcode genes (61). Depending on the species in question, certain genes, such as




Fig. 10. Phylogenetic analysis of *Fragaria* based on full-length chloroplast sequences (47).

matK, may provide a more accurate reflection of plant evolutionary history than ITS2 (58). However, DNA barcoding is limited in its ability to account for environmental influences on polygenic traits (58, 59). Similarly, challenges exist for monogenic species, where successful identification depends on the availability of the exact target gene for sequencing. Unfortunately, the limited reference data for *Fragaria* in NCBI makes it difficult to obtain precise identification (Table 5)(52, 60).

Despite these limitations, DNA barcoding remains an effective method for species identification and for elucidating evolutionary histories (Table 6) (62). A previous study demonstrated that DNA barcoding analysis of *Fragaria* species can produce a phylogenetic tree with a bootstrap value exceeding 99% (31). Additionally, vegetation evolutionary analyses conducted on Barro Colorado Island and Ailao Mountain using *rbcl*, *matK* and *trnH-psbA* genes have shown high reliability (62). These findings underscore the important role of DNA analysis in understanding species evolution and highlight the urgent need to conserve genetic diversity (63).

Acknowledgements

Authors would like to thank Dean and the Head of the Laboratory of Genetics and Breeding, Faculty of Biology, Universitas Gadjah Mada for the facilities and the support provided. The author also would like to express our sincere gratitude to all staff members Faculty of Biology for their valuable assistance and attention.

Authors' contributions

GRA carried out validation data, supervision, drafted and finalization the manuscript. MAA carried out molecular genetic studies, taking data and articulated in the sequence alignment and drafted the manuscript. MF carried out molecular genetic studies molecular genetic studies, taking data and articulated in the sequence alignment and drafted the manuscript. MSB carried out molecular genetic studies molecular genetic studies, taking data and articulated in the sequence alignment and drafted the manuscript. RSK participated in the design of the study and performed the statistical analysis. AW conceived of the study and participated in its design and coordination. All authors read and approved the final manuscript.

Table 5. Characterization of genetic marker in plant DNA barcoding (56)

Marker	Type	Basa size	Number of accessions in genebank	Number of species in genebank
nrITS2	Nuclear	157–670	111370	57579
matK	Plastid	862–910	34647	22701
rbcl	Plastid	654–654	27725	20374

Table 6. Challenges and opportunities of DNA barcoding

Challenges	Opportunities
Alien species	An increasing number of species for identification Identification of hybrid species
Monogenic and Polygenic	Research involving various other Barcoding genes
Limited information	Collaborative research with research institutions Opportunities for additional research (Research Gap)

Compliance with ethical standards

Conflict of interest: Authors do not have any conflict of interests to declare.

Ethical issues: None

References

- Azizah UDL, Yulianti F, Adiredjo AL, Sitawati S. Genetic relationship analysis of strawberry (*Fragaria* sp.) germplasm based on morphology character and random amplified polymorphic DNA (RAPD). *Plantropica J Agric Sci*. 2019;4(1):77–85. Available from: <https://jpt.ub.ac.id/index.php/jpt/article/view/152>
- Pramudhita DA, Azzahra F, Arfat K, Magdalena R, Saidah S. Strawberry plant diseases classification using CNN based on MobileNetV3-Large and EfficientNet-B0 Architecture. *J Ilm Tek Elektro Komput dan Inform* 2023;9(3):522–34.
- Hanif Z, Ashari H. Sebaran Stroberi (*Fragaria × ananassa*) di Indonesia. *Semin Nas Pekan Inov Teknol Hortik Nas Penerapan Inov Teknol Hortik dalam Mendukung Pembang Hortik yang Berdaya Saing dan Berbas Sumber Daya Genet Lokal Teknol Hortik Nas Penerapan I*. 2013;(May):87–95.
- Dwiastuti ME, Soesanto L, Aji TG, Devy NF, Hardiyanto. Biological control strategy for postharvest diseases of citrus, apples, grapes and strawberries fruits and application in Indonesia. *Egypt J Biol Pest Control*. 2021;31(1):141. <https://doi.org/10.1186/s41938-021-00488-1>.
- Balasooriya HN, Dassanayake KB, Seneweera S, Ajlouni S. Interaction of elevated carbon dioxide and temperature on strawberry (*Fragaria × ananassa*) growth and fruit yield. *International Journal of Agricultural and Biosystems Engineering*. 2018;12(9):279–87.
- Suminarti NE, Sebayang HT, Maghfoer MD, Bulan B. Effect of para-net shade level on plant microenvironment, growth and yield of three strawberry varieties. *Biodiversitas J Biol Divers*. 2023;24(4):2149–55. Available from: <https://smujo.id/biodiv/article/view/13267>
- Haqiqi I, Damanhuri, Kendarini N, Agisimanto D. Studi Keberhasilan Persilangan Stroberi (*Fragaria × ananassa* Duch). *J Produksi Tanam*. 2015;3(2):107–12.
- Zareei E, Karami F, Aryal R, Saed-Moucheshi A. Genotypic by phenotypic interaction affects the heritability and relationship among quantity and quality traits of strawberry (*Fragaria × ananassa*). *New Zeal J Crop Horti Sci*. 2022 Feb 20;1–20. Available from: <https://doi.org/10.1080/01140671.2022.2039725>
- Liston A, Cronn R, Ashman T. *Fragaria*: A genus with deep historical roots and ripe for evolutionary and ecological insights. *Am J Bot*. 2014;101(10):1686–99. Available from: <https://bsapubs.onlinelibrary.wiley.com/doi/10.3732/ajb.1400140>
- Hummer KE, Hancock J. *Plant Genetics and Genomics: Crops and Models*. New York (United States of America): Springer. 2009. Available from: <https://doi.org/10.1007/978-0-387-77491-6>
- Hummer KE, Nathewet P, Yanagi T. Decaploidy in *Fragaria iturupensis* (Rosaceae). *American Journal of Botany*. 2012; 96:713–16.
- Rahayu DA, Jannah M. DNA Barcode Hewan dan Tumbuhan Indonesia. 2019: 9–25.
- Kim H, Margie L. Global conservation strategy for *Fragaria* (Strawberry). *Scr Horti*. 2008;(March):83. ISBN 978 90 6605 129 4.
- Kurniasih MD. Menumbuhkan Karakter Konservasi Biodiversitas Melalui Penerapan Species Identification and Response Software. *Edu Sains J Pendidik Sains Mat*. 2018;6(2):30.
- Kasiamdari RS, Aristya GR, Inayati E. Phylogenetic relationships of nine cultivars of strawberries (*Fragaria* spp.) based on anatomical and morphological characters. *Planta Trop J Agro Sci*. 2017;5(2):116–26.
- Oktarina DO. Pertumbuhan Dan Produksi Stroberi (*Fragaria* sp.) dengan Pemberian Berbagai Konsentrasi Pupuk Organik Cair (POC) Secara Hidroponik Substrat. *JOM F*. 2017;4(1):1–12.
- Sudiarta IP, Ngurah G, Susanta A, Gede D, Selangga W, Getas M, et al. Detection of strawberry vein banding virus (SVBV) and identification of viruliferous insects associated with strawberry plants (*Fragaria* sp.) in Bali. *J Perlindungan Tanam Indonesia*. 2021;25(2):121–26.
- Sánchez-Sevilla JF, Vallarino JG, Osorio S, Bombarely A, Posé D, Merchante C, et al. Gene expression atlas of fruit ripening and transcriptome assembly from RNA-seq data in octoploid strawberry (*Fragaria × ananassa*). *Sci Rep*. 2017;7(1):1–13.
- Sargent DJ, Fernández-Fernández F, Ruiz-Roja JJ, Sutherland BG, Passey A, Whitehouse AB, et al. A genetic linkage map of the cultivated strawberry (*Fragaria × ananassa*) and its comparison to the diploid *Fragaria* reference map. *Mol Breed*. 2009;24(3):293–303. Available from: <http://link.springer.com/10.1007/s11032-009-9292-9>
- Staudt G. Strawberry Biogeography, Genetics and Systematics. *Proc. VIth Internat. Strawberry Symposium*. 2009:71–84.
- Song Y, Li C, Liu L, Hu P, Li G, Zhao X, Zhou H. The population genomic analyses of chloroplast genomes shed new insight on the complicated ploidy and evolutionary history in *Fragaria*. *Front. Plant Sci*. 2023;13:1065218. Available from: <https://doi.org/10.3389/fpls.2022.1065218>
- Potter D, Luby JJ, Harrison RE. Phylogenetic relationships among species of *Fragaria* (Rosaceae) inferred from Non-Coding Nuclear and Chloroplast DNA sequences. *Syst Bot*. 2000;25(2):337. Available from: <https://www.jstor.org/stable/2666646?origin=crossref>
- Saroinsong D, Panelewen VVJ, Laoh OEH, Pakasi CBD. Agribisnis Tanaman Stroberi Di Desa Rurukan Kecamatan Tomohon Timur. *Eugenia*. 2012;18(3). Available from: <https://ejournal.unsrat.ac.id/index.php/eugenia/article/view/4099>
- Post NW, Gilbert, CC, Pugh KD, Mogle CS. Implication of outgroup selection in The phylogenetic inference of humanoids and fossil hominins. *Journal of Human Evolution*. 2023;184(103437).
- Buti M, Moretto M, Barghini E, Mascagni F, Natali L, Brilli M, et al. The genome sequence and transcriptome of *Potentilla micrantha* and their comparison to *Fragaria vesca* (the woodland strawberry). *Gigascience*. 2018;7(4):1–14. Available from: <https://academic.oup.com/gigascience/article/doi/10.1093/gigascience/giy010/4860432>
- Wolfgang A, Zachow C, Müller H, Grand A, Temme N, Tilcher R, et al. Understanding the Impact of Cultivar, seed origin and substrate on bacterial diversity of the sugar beet rhizosphere and suppression of soil-borne pathogens. *Front Plant Sci*. 2020;11 (September):1–15. Available from: <https://www.frontiersin.org/article/10.3389/fpls.2020.560869/full>
- Qarni A, Muhammad K, Wahab A, Ali A, Khizar C, Ullah I, et al. Molecular Characterization of wild and cultivated strawberry (*Fragaria × ananassa*) through DNA barcode markers. *Hesham AEL, editor. Genet Res (Camb)*. 2022;2022:1–14. Available from: <https://www.hindawi.com/journals/gr/2022/9249561/>
- Gostel MR, Kress WJ. The Expanding Role of DNA barcodes: indispensable tools for ecology, evolution and conservation. *Diversity*. 2022;14(213):1–23.
- Kramer A, Havens K. Plant conservation genetics in a changing world. *Trends Plant Sci*. 2009;14;11:599–607.
- Turhadi T, Hakim L. Evaluasi Locus Kloroplas untuk DNA barcoding pada Marga *Stelechocarpus* (Annonaceae) Secara insilico. *Agro Bali Agric J*. 2023;6(1):56–64. Available from: <https://ejournal.unipas.ac.id/index.php/Agro/article/view/1105>
- Li C, Cai C, Tao Y, Sun Z, Jiang M, Chen L, et al. Variation and evolution of the whole chloroplast genomes of *Fragaria* spp. (Rosaceae). *Front Plant Sci*. 2021;12(October).

32. Hollingsworth PM, Forrest LL, Spouge JL, Hajibabaei M, Ratnasingham S, van der Bank M, et al. A DNA barcode for land plants. *Proc Natl Acad Sci U S A*. 2009;106(31):12794–97.
33. Pfenninger M, Cordellier M, Streit B. Comparing the efficacy of morphologic and DNA-based taxonomy in the freshwater gastropod genus *Radix* (Basommatophora, Pulmonata). *BMC Evol Biol*. 2006;6:1–14.
34. Potter D, Eriksson T, Evans RC, Oh S, Smedmark JEE, Morgan DR, et al. Phylogeny and classification of Rosaceae. *Plant Syst Evol*. 2007;266(1-2):5–43. Available from: <http://link.springer.com/10.1007/s00606-007-0539-9>
35. Pang X, Song J, Zhu Y, Xu H, Huang L, Chen S. Applying plant DNA barcodes for Rosaceae species identification. *Cladistics*. 2011;27(2):165–70. Available from: <https://onlinelibrary.wiley.com/doi/10.1111/j.1096-0031.2010.00328.x>
36. Hebert PDN, Cywinska A, Ball SL, DeWaard JR. Biological identifications through DNA barcodes. *Proc R Soc London Ser B Biol Sci*. 2003;270(1512):313–21. Available from: <https://royalsocietypublishing.org/doi/10.1098/rspb.2002.2218>
37. Savolainen V, Cowan RS, Vogler AP, Roderick GK, Lane R. Towards writing the encyclopaedia of life: An introduction to DNA barcoding. *Philos Trans R Soc B Biol Sci*. 2005;360(1462):1805–11.
38. Stoeckle M. The Barcode of Life. *Bioscience*. 2003;53(9):2–3.
39. Hebert PDN, Stoeckle MY, Zemlak TS, Francis CM. Identification of birds through DNA barcodes. *PLoS Biol*. 2004;2(10):1657–63.
40. Gallal-Khallal A, Abdelbaset-Donya A, Hamza W, Mohammed-Geba K. Molecular tools for assuring human health and environment-friendly frozen shellfish products in the United Arab Emirates markets. *Food Chemistry: Molecular Sciences*. 2021;3:1–9. Available from: <https://doi.org/10.1016/j.fochms.2021.100028>
41. Friedheim S. Comparison of Species Identification methods: DNA barcoding versus morphological taxonomy. *Manoa Horizons*. 2016;1(1):74–86.
42. Frézal L, Leblois R. Four years of DNA barcoding: Current advances and prospects. *Infect Genet Evol*. 2008;8(5):727–36. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S1567134808001238>
43. Farooq Q, Shakir M, Ejaz F, Zafar T, Durrani K, Ullah A. Role of DNA barcoding in plant biodiversity conservation. *Sch Int J Biochem*. 2020;03(03):48–52. Available from: https://saudijournals.com/media/articles/SIJB_33_48-52.pdf
44. Hollingsworth PM. Refining the DNA barcode for land plants. *Proc Natl Acad Sci U S A*. 2011;108(49):19451–2.
45. Kress WJ, Erickson DL. DNA barcodes: Genes, genomics and bioinformatics. *Proc Natl Acad Sci*. 2008;105(8):2761–62. Available from: <https://pnas.org/doi/full/10.1073/pnas.0800476105>
46. Badotti F, de Oliveira FS, Garcia CF, Vaz ABM, Fonseca PLC, Nahum LA, et al. Effectiveness of ITS and sub-regions as DNA barcode markers for the identification of Basidiomycota (Fungi). *BMC Microbiol*. 2017;17(1):42. Available from: <http://bmcmicrobiol.biomedcentral.com/articles/10.1186/s12866-017-0958-x>
47. Sun J, Sun R, Liu H, Chang L, Li S, Zhao M, et al. Complete chloroplast genome sequencing of ten wild *Fragaria* species in China provides evidence for phylogenetic evolution of *Fragaria*. *Genomics*. 2021;113(3):1170–9. Available from: <https://doi.org/10.1016/j.ygeno.2021.01.027>
48. Lestari R. Identification and assessment of invasive plant species at Bogor Botanic Gardens, Indonesia. *IOP Conf Ser Earth Environ Sci*. 2021;800(1).
49. Langmaier M, Lapin K. A Systematic review of the impact of invasive alien plants on forest regeneration in European Temperate Forests. *Front Plant Sci*. 2020;11(September):1–15.
50. Davis MA, Thompson K. Eight Ways to Be a Colonizer; Two ways to be an invader: A proposed nomenclature scheme for invasion ecology. *Bulletin of the Ecological Society of America*. 2000;81;3: 226–30. <https://www.jstor.org/stable/20168448?origin=crossref>
51. Thielecke L, Aranyosy T, Dahl A, Tiwari R, Roeder I, Geiger H, et al. Limitations and challenges of genetic barcode quantification. *Sci Rep*. 2017;7(1):43249. Available from: <https://www.nature.com/articles/srep43249>
52. Raclariu AC, Heinrich M, Ichim MC, de Boer H. Benefits and limitations of DNA barcoding and metabarcoding in herbal product authentication. *Phytochem Anal*. 2018;29(2):123–8.
53. Njuguna W. DNA Barcoding: Unsuccessful for species identification in *Fragaria* L. *Plant Genet Resour*. 2011;349–56.
54. Smith AL, Hodgkinson TR, Vilellas J, Catford JA, Csörgő AM, Blomberg SP, et al. Global gene flow releases invasive plants from environmental constraints on genetic diversity. *Proc Natl Acad Sci*. 2020;117(8):4218–27. Available from: <https://pnas.org/doi/full/10.1073/pnas.1915848117>
55. Schulze J, Stoll P, Widmer A, Erhardt A. Searching for gene flow from cultivated to wild strawberries in Central Europe. *Ann Bot*. 2011;107(4):699–707.
56. Hollingsworth PM, Graham SW, Little DP. Choosing and using a plant DNA barcode. *PLoS One*. 2011;6(5).
57. Agrios G. Plant pathology. 5th Edition. Amsterdam (Netherlands): Elsevier Academic Press;2005.
58. Devi MP, Dasgupta M, Mohanty S, Sharma SK, Hegde V, Roy SS, et al. DNA barcoding and ITS2 secondary structure predictions in taro (*Colocasia esculenta* L. Schott) from the Northeastern hill region of India. *Genes (Basel)*. 2022;13(12):1–13. Available from: https://www.mdpi.com/2073-4425/13/12/2294?type=check_update&version=1#
59. Cowan RS, Fay MF. Challenges in the DNA Barcoding of Plant Material. In: Sucher N, Hennell J, Carles M, editors. *Methods in Molecular Biology*. Humana Press; 2012 [cited 2023 Des 9];23–33. https://doi.org/10.1007/978-1-61779-609-8_3
60. Li Y, Pi M, Gao Q, Liu Z. Updated annotation of the wild strawberry *Fragaria vesca* V4 genome. *Hortic Res*. 2019;6(61):1–9. Available from: <http://dx.doi.org/10.1038/s41438-019-0142-6>
61. Ma Y, Ding X, Qanbari S, Weigend S, Zhang Q, Simianer H. Properties of different selection signature statistics and a new strategy for combining them. *Heredity (Edinb)*. 2015;115(5):426–36.
62. Kang Y, Deng Z, Zang R, Long W. DNA barcoding analysis and phylogenetic relationships of tree species in tropical cloud forests. *Sci Rep*. 2017;7(September):1–9.
63. Gostel MR, Kress WJ. The expanding role of DNA barcodes: Indispensable tools for ecology, evolution and conservation. *Diversity*. 2022;14(213):1–23.

Additional information

Peer review: Publisher thanks Sectional Editor and the other anonymous reviewers for their contribution to the peer review of this work.

Reprints & permissions information is available at https://horizonpublishing.com/journals/index.php/PST/open_access_policy

Publisher's Note: Horizon e-Publishing Group remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Indexing: Plant Science Today, published by Horizon e-Publishing Group, is covered by Scopus, Web of Science, BIOSIS Previews, Clarivate Analytics, NAAS, UGC Care, etc
See https://horizonpublishing.com/journals/index.php/PST/indexing_abstracting

Copyright: © The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited (<https://creativecommons.org/licenses/by/4.0/>)

Publisher information: Plant Science Today is published by HORIZON e-Publishing Group with support from Empirion Publishers Private Limited, Thiruvananthapuram, India.