



REVIEW ARTICLE

Organic farming: A strategy for a sustainable and secure food system

Manisha Arora Pandit¹, Saloni Gulati², Mallika Kalia³, Charu Dogra Rawat⁴ & Jasleen Kaur^{2*}

¹Department of Zoology, Kalindi College, University of Delhi, New Delhi 110 008, India

²Department of Botany, Dyal Singh College, University of Delhi, New Delhi 110 003, India

³Research and Business Development Analyst, Biokriti, Noida 201 301, Uttar Pradesh, India

⁴Department of Zoology, Ramjas College, University of Delhi, New Delhi 110 007, India

*Correspondence email - jasleen@dsc.du.ac.in

Received: 05 January 2025; Accepted: 05 May 2025; Available online: Version 1.0: 17 June 2025; Version 2.0: 01 July 2025

Cite this article: Manisha AP, Saloni G, Mallika K, Charu DR, Jasleen K. Organic farming: A strategy for a sustainable and secure food system. Plant Science Today. 2025; 12(3): 1-17. <https://doi.org/10.14719/pst.7076>

Abstract

The growing human population presents a global challenge in providing adequate food, shelter and a healthy lifestyle. Sustainable food production and environmental stewardship are essential for addressing food security and environmental preservation. Organic farming is gaining popularity due to its perceived safety, health and ecological benefits. However, there are differing viewpoints on its role in sustainability. Despite advances in agricultural productivity and the use of GMOs, hunger persists in the poorest countries. This review aims to answer whether organic farming is the future of sustainable agriculture to address the world's growing food needs. The Green Revolution has led to increased agricultural output and income, but it has also led to lower-quality food and environmental degradation due to chemical use. Health concerns, such as hormone imbalances and cancers, have been raised due to these practices. Organic farming, which avoids artificial inputs and uses biological control methods, has gained momentum in developed nations. It promotes ecological restoration, plant, animal and soil health and improves biodiversity. Sustainable agriculture should support both the economy and the environment.

Keywords: biological control methods; environmental stewardship; food security; global challenge; organic farming; sustainable agriculture

Introduction

The exponentially growing human population presents a global challenge in terms of the future provision of adequate food, shelter and a healthy lifestyle. The struggle is compounded when one considers the imperative need to preserve the environment and the Earth's natural resources for future generations. Sustainable food production and environmental stewardship must be aimed to address food security and preservation. Addressing this will require a collaborative One Health effort, focusing on the fitness of people, animals and the environment since they are inseparably linked and cannot be considered individually. The quality of available food is a significant cause for concern for the growing human population. Therefore, the gravitation towards organic foods worldwide can be easily understood. The demand is mainly fuelled by consumers in developed nations, whose unease stems from indiscriminate and large-scale use of chemical fertilizers, pesticides, etc., in traditional agricultural practices. Generally, the public in these countries is noticeably choosing organic foods since it is perceived as a safer and healthier alternative (1). There is also an insight that organic farming is a more sustainable agricultural practice, is better suited for environmental health and aids in climate guard and overall

animal well-being (2, 3). It is also thought that a large-scale move towards organic agriculture will aid in reducing worldwide hunger (4). Surveys have also shown that consumers feel food security is threatened by excessive use of chemicals and genetically modified organisms (GMOs). In contrast to the consumers of developed nations, most consumers in developing nations are still ignorant about the benefits of organic agriculture. Nevertheless, there has been a shift towards organic foods by the well-off consumers in these countries, possibly due to European influence (5, 6). Incongruous viewpoints exist for organic agriculture and its role in sustainability, where some regard it as ineffective while others visualize considerable benefits (7–11).

Apart from trying to fulfil the growing nutrition needs of an increasing human population, plant products are also used as energy sources. The mounting demand for improved crop yields has led to an augmented use of chemical fertilizers and better plant varieties, including GMOs. These steps have improved agricultural productivity and marked the fulfilment of the green revolution (12, 13). Despite these advances, hunger persists in most of the poorest countries in Asia and Africa (14, 15). Food requisitions will only increase in the coming years based on projections of an escalating world

population and incomes. To satisfy these demands, agricultural output needs to improve by a minimum of 60 % and a possible 100 % by the year 2050 regardless of the scarcity of crucial natural resources such as land and water (16, 17). Attention also needs to be paid to avoiding further environmental degradation while addressing the growing food security concerns of the world (18). Prevention of further ecosystem damage should be of paramount importance and the adoption of methodologies to improve food security should be cognizant of these factors (19–22). These issues raise the question of whether organic farming is the future of sustainable agriculture that will address the ever-increasing food needs of the world. This review is, therefore, an attempt to answer this question by analyzing the available literature on the topic and focusing on understanding the different modalities of certified organic farming, along with its impact on the economy, environment and human health.

The legacy of the Green Revolution: achievements and challenges

Green revolution technologies enabled several countries to rise from food deficit situations to increased agricultural output, self-sufficiency, improved food exports and an increase in agriculture-based income (23). However, to attain these goals, most of these countries ended up producing lower-quality food and suffered environmental degradation predominantly due to pollution stemming from the indiscriminate use of chemicals necessary for high agricultural productivity.

Many health concerns have also been raised due to the use of green farming methodologies. These include hormone imbalances, cancers, respiratory issues, birth defects, etc. Chemical fertilizers augment nitrate levels in groundwater, as shown by a study of wells in Kalpitiya where only 56 % of the wells, as per WHO standards had safe drinking water (24). High nitrate levels were also seen in some vegetables mostly due to contamination of soil from shallow nitrate-polluted groundwater sources. These pollutants ultimately destroy and damage natural ecosystems. It is, therefore, time to digress from conventional, chemical-intensive farming methods and shift to practices aimed at restoring declining soil fertility and ensuring food security and availability of resources for future generations (25).

Organic farming systems

The shift towards organic farming practices was first seen in some European nations, mostly in response to the overuse of chemical fertilizers in agriculture. It was encouraged by agriculturists who advocated traditional farming practices based on biological control methods (26). Even though the movement was small for a few decades, it slowly gained momentum in the 1970s, largely due to increasing public awareness and health concerns about the overuse of chemicals in conventional agriculture (27). Eventually, some developed nations started subsidies for organic farming along with certifying organic produce. The United States and EU eventually formulated regulations, policies, standards, labelling, research and sponsoring of organic produce. Organic farmers were praised for their environmental consciousness and offered compensation for increased production costs associated with organic farming (28). Recently, developing

nations have also joined the organic farming revolution by adopting policies that support organic farmers.

As per the United States Department of Agriculture (USDA), organic farming is "a system which avoids and largely excludes the use of artificial inputs". It primarily banks on the use of biological control methods, manures from animals and organic farm waste and rotation of crops to maintain soil fertility and nutrient input (2). These practices favour plant, animal and soil health and aid in maintaining more robust ecosystems (29). The subsequently increased agroecosystem resilience helps in combating the detrimental consequences of climate change thereby promoting ecological restoration. Organic methodologies benefit the health of their consumers by providing plant and animal products that are free from chemicals such as pesticides, drugs and antibiotics (30–32). These have a positive impact not only on the end users but also on slowing the propagation of antibiotic-resistant bacteria. Organic practices also contribute to improved biodiversity as farmers depend on pollinators and biocontrol agents to manage crop production (33).

Sustainable agriculture should support both the economy and the environment. As a farming practice, it should provide solutions for the concerns being raised by conventional farming techniques. In this respect, organic agricultural processes fulfil most of the demands required for attaining sustainable food production (27, 34, 35) (Fig. 1).

Global adoption of organic farming

Organic farming is a rapidly growing agricultural category that contributes to about 1 % of the world's total agricultural area. In its present form, organic farming originated in the early 1970s even though it is an age-old form of agriculture that was practised quite commonly in Asian countries like Sri Lanka, which have been known to produce staples such as rice using organic farming methods (36). Today, both Governmental and private standards exist for organic agricultural practices (37). These are built on the standards formulated by the International Federation of Organic Agriculture Movements (IFOAM) (<https://www.ifoam.bio/news/global-organic-area-grows-more-ever>). These standards are quite comparable worldwide but may have regional differences. They cover all aspects of organic farming related to crops, livestock, aquaculture etc. They also state the actions that are favoured as well as those that are restricted in organic farming (Fig. 2). Some examples of necessary actions include rotation of crops with legumes and mixed crop farming to balance and recycle nutrients and applying only organic fertilizers. The use of chemicals in any form is strictly prohibited. Similarly, in the case of livestock, only organic fodder must be used to feed the animals possibly from the farm where they are being raised and have access to open spaces.

Accredited certification is required to ensure compliance with organic standards and farm inspections should be done on an annual basis (37). It is therefore imperative to keep records of all farming activities. Certification, in developed countries, is mostly done for singular farms while it is carried out in groups in developing nations. This helps reduce the cost for farmers and certifiers (38, 39).

Even though the global area under certified organic




Fig. 1. Cultivating a sustainable future: exploring the practices and benefits of organic farming.




Fig. 2. Prohibited and required activities in organic farming.

agriculture has grown immensely over the past couple of decades, from 15 million ha in 2000 to 96 million ha in 2022 (Fig. 3), it still accounts for just 2 % of the total agricultural land area worldwide. Organic farming lands are expanding worldwide, with a presence on all continents. Oceania boasts the largest share of organic farmland, covering an impressive 53.2 million ha, which is over half of the total global land area. Europe comes in second with 18.5 million ha under organic management. Latin America follows closely behind with a significant area of 9.5 million ha dedicated to organic farming (Fig. 3). The number of organic producers skyrocketed in 2022, reaching nearly 4.5 million globally, reflecting a significant increase of 25.6 % compared to 2021. Asia, Africa and Europe dominated the organic producer landscape, accounting for almost 93 % of all producers. India emerged as the leader boasting an impressive 2.5 million producers in 2022, followed by Uganda and Thailand. While growth was observed in Oceania, Europe and Asia, the numbers declined in Latin America, North America and Africa in 2022 (40).

The organic market globally, reached around 135 billion euros in 2022, reflecting growing consumer interest in organic products and aligned with the increased global area of its production as per the FiBL survey. The United States remained the world's leading organic market, with sales reaching 56.6 billion euros. While some European countries saw a decline in sales, Germany (15.3 billion euros) remained a strong market and China (12.4 billion euros) emerged as the new number 3. Certain countries such as Canada, Japan and Estonia showed positive growth trends. Canada and the United States experienced positive growth in retail sales, indicating continued consumer demand for organic options in these regions. Most European countries spent a larger portion of their incomes on organic food and certain European countries such as Denmark, Austria and Switzerland improved their organic food share to around 11.2 - 12 % of the total food market (40). Switzerland held the title for the highest per capita spending on organic food, averaging 437 euros per person

while Denmark retained a large 12 % share of organic products, in its total food market. On the other hand, the demand for organic food and animal produce remains contained to a small high-income section of society in the developing nations due to its significantly higher price (37, 41).

Medicinal and aromatic plants showed the highest percentage increase in organic area (81.5 %) between 2021 and 2022 while berries and olives exhibited the most significant decrease in organic area in 2022 compared to 2021. Nut production has steadily increased in organic area over the past decade (181.2 %). Overall, the total permanent organic cropland area has increased slightly (0.8 %) between 2021 and 2022. However, compared to 2013, there's a significant increase of 96.0 % (Fig. 4) (40). Government subsidies, ability to withstand higher production risk, access to correct information and marketplaces with consumers willing to pay the higher price associated with organic products are some key factors motivating farmers to shift to organic farming (42–46). Organic farming also comes with its own constraints such as prohibiting the use of chemical fertilizers and pesticides and policy risks which further reduce its wide-scale adoption (44, 47). These issues can be mitigated by providing proper training, support and subsidies to farmers, particularly in developing countries, to ensure its wider acceptance (48).

Shifts in farming practices

Modern farming practices can significantly negatively impact our environment and health due to increased greenhouse gases, soil, water pollution, etc. Organic agriculture on the other hand offers a more sustainable alternative since it helps to reduce carbon emissions, improve soil health, cleanses air and water and supports natural ecosystems. Importantly, organic farming avoids using harmful pesticide residues. The key difference lies in how each method deals with pests, weeds and nutrients. Conventional farming relies heavily on chemical solutions - synthetic pesticides, herbicides and fertilizers. On the other hand, organic farming prioritizes natural methods




Fig. 3. Organic farmland and percentage of global area (40).




Fig. 4. Comparison of organic permanent cropland between 2013, 2021 and 2022 (Source: FiBL survey 2024, based on information from the private sector, certifiers and governments (40).

like biodiversity and composting to achieve healthy and abundant harvests. Organic farming is more than just saying no to chemicals. It's about reviving ancient practices like crop rotation and using composted manure and green manure crops. Importantly, these methods are adapted for economic viability in today's world. The core of organic production isn't simply replacing synthetic inputs with natural ones. It's about fostering a healthy and balanced ecosystem within the farm itself. Organic farmers employ a variety of techniques to achieve this, promoting biodiversity while ensuring long-term soil fertility (49).

Farming in harmony with nature

Harmonious agricultural practices aim to meet the current population's food needs without compromising on future generations' needs (50). Agriculture greatly impacts the environment and conventional practices can cause widespread deforestation, pollution and soil degradation (51). The advantages of organic farming are manifold and include a reduction in the production and release of greenhouse gases, better food quality due to enhanced polyphenol production in organically grown plants, improved environment and reduced pollution due to the non-use of synthetic fertilizers, soil enrichment due to addition of nutrient-rich composts and crop rotation among others. Apart from these advantages, organic farming also generates rural employment, reduces procurement costs for chemical fertilizers and preserves groundwater resources (52–55).

Organic farming focuses on the rotation of crops, multiple cropping systems, permaculture, integrated farming systems, agroforestry, biological control, etc., all crucial to maintaining sustainability (19). In the long term, the conception of sustainable agricultural mechanisms will help meet the needs of the increasing population, prevent further degradation of the environment and reduce the impact of farming on climate change. In other words, organic farming

can be the way forward to move in harmony with nature and equalize society, economics and the environment.

Organic farming: A key ingredient for sustainable global agriculture

Agricultural practices must evolve to align with the United Nations Sustainable Development Goals by 2030 (Table 1). While debating between incremental improvements in agriculture versus transformative changes there is always a focus on different strategies for achieving long-term sustainability goals in the agriculture sector. Incremental steps can provide short-term solutions and build momentum, while transformative changes are crucial for addressing underlying systemic issues and creating a resilient and sustainable food system for the future. Achieving the United Nations Sustainable Development Goals (SDGs) by 2030 will require combining both approaches with the challenge to find a balance between the two so that there are maximum benefits across environmental, social and economic dimensions while ensuring food security and livelihoods for all (56).

The Research Institute of Organic Agriculture FiBL's statistics indicate that the Oceania region consists of Australia, New Zealand and the Pacific Island states. Despite facing numerous challenges such as natural disasters, Australia is notably successful in organic farming and sets an example for other countries. Australia manages most of the organic land in the Oceania region, totalling 35.7 million ha, primarily used for extensive grazing. This accounts for a significant portion of the region's organic land and agricultural lands. It has created a supportive regulatory environment by developing legislation specifically for organic agriculture.

Additionally, several other countries in Oceania have national standards for organic production, although they lack specific legislation. Supportive policies, resilience in the face of challenges and the potential for organic agriculture have contributed a lot to the sustainable development of Australia. Other countries can enhance their organic farming sectors and

Table 1. Organic farming and the sustainable development goals (SDGs)

SDG	Themes	How organic farming contributes	Reference
SDG 2: Zero hunger	Aims to end hunger, malnutrition and provide universal access to safe, nutritious and sufficient food throughout the year.	Improves soil health and fertility, leading to long-term food security. Promotes biodiversity, which increases resilience of crops to pests and diseases. Encourages production of diverse and nutritious crops.	(15, 123-126)
SDG 3: Good health and well-being	Ensure healthy lives and promote well-being for all at all ages	Reduces exposure to harmful synthetic pesticides and fertilizers, promoting public health. Encourages consumption of fresher, more nutritious food	
SDG 6: Clean water and sanitation	It aims to provide clean water, hygiene and sanitation for all	Reduces water pollution from agricultural runoff of chemicals. Promotes water conservation practices	
SDG 8: Decent work and economic growth	Promoting sustainable, inclusive economic growth and full productive employment for all	Creates new job opportunities in organic farming and certification. Supports fair trade practices for farmers	
SDG 12: Responsible consumption and production	Ensure sustainable consumption and production patterns	Reduces the environmental impact of food production. Promotes local food systems and reduces food waste	
SDG 13: Climate action	Take urgent action to combat climate change and its impacts	Reduces greenhouse gas emissions associated with conventional agriculture. Builds soil health, which helps to sequester carbon	
SDG 15: Life on land	Promotes the sustainable use of terrestrial ecosystems. Withstand desertification and reverse land degradation. Protect, restore and sustainably manage forests and biodiversity	Promotes biodiversity in agricultural ecosystems. Reduces soil erosion and promotes soil health	

achieve similar successes in promoting sustainable agriculture and food systems by following Australia's approach.

Cultivating a sustainable future: The synergy between organic farming and sustainable development

Agriculture is widely acknowledged as crucial for achieving sustainable development over the long haul. Integrating small-scale farmers in developing nations into the transformation of agrifood systems is essential for achieving the Sustainable Development Goals by 2030, as emphasised by FAO (56). Goal 2 of the 2030 sustainability agenda, aimed at ending hunger, underscores the central role of agriculture. Additionally, agriculture contributes, to varying degrees, to all other Sustainable Development Goals. The initial Green Revolution was prompted by a rising number of undernourished individuals, leading to the global spread of agricultural technologies such as pesticides and fertilizers to developing nations (57). Despite the initial adoption of sustainable farming practices during this period, their implementation was inconsistent. While chemical inputs contributed to increased yields worldwide, concerns about the environmental and human health impacts of intensive chemical farming spurred the emergence of modern organic farming. Organic agriculture gained momentum in the 1970s in response to growing awareness of the harmful environmental effects of industrial pesticides and fertilizers.

Numerous organisations and member states have endorsed the "Zero Hunger Challenge" in alignment with United Nations Sustainable Development policies (18). This initiative aims to achieve zero undersized children under the age of two, 100 % year-round adequate food, sustainable food systems, a 100 % increase in income and smallholder productivity and zero food waste. Promoting sustainable development through agriculture has predominantly been associated with organic farming. However, despite its global popularity, sustainable organic agriculture faces challenges and limitations.

The greening wave: emerging trends in sustainable organic farming

Organic agriculture, a leading and rapidly expanding movement in sustainability, is also referred to as regenerative farming. It lies at the core of many emerging food and beverage sustainability initiatives. This approach aims to phase out synthetic fertilizers, enhance agricultural productivity and yields, promote biodiversity, conserve water and sequester carbon in the soil. Aligned with organic farming practices, the merits of organic farming encompass integrated pest management (IPM) to reduce pesticide usage, crop rotation, planting nitrogen-fixing cover crops and implementing mulching to enhance soil health (58). Additionally, adopting no-till or reduced-tillage methods reduces soil compaction. Precision farming, guided by soil analysis, optimizes the application of organic and synthetic fertilizers, while innovative irrigation systems maximize yields and minimize water wastage, erosion and salinization (52, 53, 59–61). These practices safeguard soil health and biodiversity and mitigate the environmental impacts of agriculture. Organic farming is increasingly valued for its health benefits and environmental stewardship. Consumer preferences for organic products, driven by health concerns and ecological consciousness, have led to greater willingness to pay premium prices for organic produce, further supporting sustainability efforts.

The seeds of success

Uncaptured yield potential

An important aspect in evaluating the potential of organic agriculture to contribute to sustainable development is its yields, compared to conventional farming. Estimating the yield influences of organic measures is challenging due to the need to account for various confounding factors. For example, if

organic farmers achieve lower yields, it could be attributed to their farming methods, but it is also plausible that these farmers are less skilled or work in less favourable environments. In such cases, organic farming might appear to have a wider yield gap even if both groups used identical technology. Conversely, it is conceivable that organic farmers are consistently more skilled than their peers, leading to an underestimation of the yield gap when directly comparing observed organic and conventional yields. Remarkably, there has been relatively little research attempting to address selection biases in the observational data when estimating the yield impacts of organic farming.

However, many studies have endeavoured to evaluate the yield impacts of organic farming based on the data derived from trials conducted at experimental sites. Experimental data are advantageous for minimising bias from confounding factors, but can pose challenges in generalizability (as discussed later). The outcomes of these studies exhibit considerable variability depending on specific conditions. In some instances, organic yields have been documented to exceed those of conventional farming, whereas in others, they were significantly lower. Several review papers have recently sought to synthesise the available evidence on this topic. Initially, research indicates that a pioneering study compiled data from various sources. They concluded that globally, organic agriculture showed an average crop yield 33 % higher than conventional methods. Organic yields in the developed countries were reportedly 9 % lower than traditional yields, whereas in the developing countries, organic practices potentially increased yields by 74 % (10). Nonetheless, this study received significant criticism on multiple fronts (62–64). Critics pointed out that many studies included in Badgley et al.'s review did not comply with the basic scientific standards of experimental design and that other appropriate studies were overlooked (62, 64). Specifically, regarding developing countries, research shows that crops receiving substantial organic nutrients with those receiving minimal or no

conventional fertilizer.

As a result, despite its widespread citation, the findings of a study are viewed sceptically and may not be considered reliable or robust in their implications (10). In recent years, three meta-analyses have provided more robust scientific comparisons of organic and conventional crop yields (65–67). These meta-analyses (Fig. 5), reveal that organic agriculture generally exhibits yield gaps ranging from 19 - 25 % across various crops. Variations are notable among different types of crops: fruits and legumes show lesser yield gaps when compared to cereals, tubers and root crops. Some indication is there that the yield gap widens with the increase in the conventional yields (65). Optimal management of the 2 systems does not significantly alter the yield gaps between the developed and developing countries (66). However, it is crucial to note that observations from developing countries are notably underrepresented in all three meta-analyses, requiring caution when interpreting statements about geographic variations in yield gaps (2).

Recent initiatives have begun addressing, through longer-term research efforts, the productivity impacts of organic farming in developing countries (68). A key problem in comparing yield levels between conventional and organic agriculture is the time frame covered by the original studies. There is a widespread assumption that yields may initially decline after transitioning to organic practices, but due to improved soil conditions, they could recover over time. The evidence supporting this assumption is, however, limited. Some studies indicate increasing organic yields over time; others report no change or even declines during long-term assessments (65, 69).

Bridging the harvest gap

Besides sunlight and optimal temperatures, plants necessitate a diverse array of nutrients for robust growth, like, phosphorus, nitrogen, potassium and various micronutrients. Soil texture, composition, water availability and challenges related to pests




Fig. 5. Mean organic crop yields in comparison to conventional yields (results from global meta-analyses).

and diseases are also pivotal factors. These factors can vary between organic and conventional farming, leading to differences in yield. Organic farming standards, for instance, ban synthetic fertilizers. While organic fertilizers can theoretically provide all necessary nutrients, managing these nutrients in organic systems can be more complex (70). Organic farming often struggles with the availability of nitrogen and phosphorus (71, 72). Nitrogen from organic sources is released gradually, which can fall short of meeting the crop's needs during critical growth phases (67). Likewise, phosphorus in organic setups may be insufficient to replace what is lost during harvest (72). Overall, achieving the optimal nutrient balance for plant growth is more challenging in organic systems because of the limited ability to modify the nutrient content of organic inputs (2).

Nutrient limitations are key in explaining the yield differences seen in organic farming. This helps clarify why yield gaps are smaller for fruits and legumes. Legumes can fix atmospheric nitrogen, reducing their dependence on external nitrogen inputs. Similarly, fruit trees, with their longer seasons of growth and elaborate root systems, are better able to take up nutrients in alignment with their growth needs (67). Regarding usage and availability of water, organic farming often holds an advantage due to the improved water-holding capacity and increased water infiltration rates observed in soils managed with organic practices. This contributes to the perception that organic systems are more robust and exhibit greater yield stability, especially during drought (72, 73). However, organic systems can be more vulnerable to pest outbreaks, leading to yield losses and increased variability (2). The prohibition of GMOs and synthetic pesticides in organic agriculture limits farmers' options for managing insect pests, weeds and plant diseases. Therefore, in regions with high incidence of pests, biological control methods struggle against persistent pests and diseases and the yield gaps in organic farming tend to be larger.

Real-world relevance of estimated yield gaps

Most comparisons of crop yields between organic and conventional agriculture are drawn from experimental trials conducted at research sites. These trials often show higher yields than seen in actual farming as farmers may not always implement the suggested management practices exactly. The comparisons would be fair if the yield discrepancies between research stations and real-world farms were similar for both organic and conventional systems. However, evidence indicates that the yield gap between research trials and farmers' fields is greater in organic farming than in conventional farming (74). This is primarily due to the increased complexity and knowledge requirements of organic farming, where yields are more dependent on timely interventions (67, 75). Thus, although the reported differences between organic and conventional farming yields may be accurate under experimental conditions, they likely undervalue the actual yield gaps seen in real-world farming scenarios.

A study aimed to validate findings from experimental stations by comparing data from multiple commercial farms across the United States (76). Their study revealed that organic farms generally achieved cereal yields about 20 % lower than

conventional farms, consistent with existing meta-analyses. However, they noted yield differences of over 50 % for certain vegetables, which exceeded previous meta-analytical estimates. In practical European contexts, a study reported even larger yield gaps, with organic cereal yields sometimes dropping by 50 % compared to conventional methods (77). Nevertheless, it is essential to approach these findings cautiously due to the limited availability of comprehensive and impartial evidence on organic yield impacts in actual fields (74, 78). Moreover, considering the higher expertise required for the success of organic farming, there is a possibility that yield gaps could widen with organic practices becoming more widespread among farmers.

Another exciting aspect is the way the yield disparities between organic and routine agriculture might evolve over the long term, especially considering technological progress. Current research does not definitively answer this question (2). It is recognized that yield gaps tend to widen as conventional yields increase. Looking forward, these gaps could widen further due to slower advancements in plant genetics. Organic standards prohibit the usage of genome editing techniques and GMOs, which are known for their ability to boost crop yields and resilience significantly (13). Conversely, organic farming has historically received less research attention and increasing research efforts could potentially help reduce these knowledge gaps over time (72).

Organic food: A source of dietary advantages?

Consumers frequently believe that organic foods offer superior nutritional value and health benefits than conventional foods (37). This perception may be influenced by the assumption that organic products have lower harmful contaminants or higher amounts of beneficial nutrients. Numerous studies have investigated whether significant variations exist in the chemical composition between organic and conventional foods. Multiple systematic reviews have found that lower levels of residues of synthetic pesticides are generally found in organic foods (79–81). The effect of this difference on human health depends on the types and quantities of pesticides used in conventional farming. The risk of exceeding maximum allowable limits is typically low in developed countries with stringent pesticide regulations (82, 83). Additionally, organic foods often contain lower concentrations of nitrate and cadmium than their conventional counterparts (80, 81). Studies typically do not reveal significant differences in microbial-bacterial or fungal contamination, however, some indicate microbial load in certain organic fruits (84).

Systematic reviews have indicated that slightly higher levels of nutritionally beneficial components, phenolics are often found in organic plant products (81, 83, 85). Findings regarding carotenoids and vitamin C varied across studies (81, 83). Organic milk and chicken are reported to have elevated omega-3 fatty acid levels (80, 81, 83). Conversely, organic foods generally exhibit slightly lower concentrations of proteins and amino acids (81). Nevertheless, the clinical relevance of these variances in nutritionally beneficial components between organic and non-organic foods remains uncertain (79, 81, 83, 86). Variances in the organic and non-organic foods compositions is expected due to the influence of farming

practices on plant chemistry (85). Organic plants typically exhibit lower levels of cadmium and nitrate, which is attributed to the absence of synthetic fertilizers in organic farming (81). The use of nitrogen fertilizers in conventional farming by focusing on proteins and carbohydrates enhances vegetative growth and limits generative growth, which involves the production of secondary metabolites (80). In animal products, enhanced levels of omega-3 fatty acids are often linked to outside grazing and greater biodiversity in the organic farms' pastures.

The chemistry of the plants is influenced by the production system and factors such as soil type, weather conditions, ripening stage at harvest, postharvest conditions and genotype (variety) also have an impact on it (80, 85). Cadmium levels, for example, vary significantly depending on soil type, potentially resulting in elevated levels in organic products (81). Moreover, management practices differ widely (80, 83). For example, organic cattle may not always graze on biodiverse pastures and conventional cattle are not necessarily confined and fed silage. Thus, there can be greater variation in food composition within organic and conventional systems than between the two (85). Numerous studies have explored the potential health effects of organic diets. Some studies have revealed that in infants, a lower risk of eczema and allergies is associated with consuming organic foods (87, 88). Additionally, a cohort study conducted in France showed a reduced incidence of obesity due to regular organic food consumption (89). A systematic review, however, concluded that there are no significant differences in health outcomes related to organic food consumption (79).

Establishing causality with observational data is difficult because organic consumers often make distinct, typically healthier, lifestyle and food choices potentially introducing selection bias in impact assessments (80, 89). Due to these complexities and the limited evidence available, definitive conclusions about the effect of organic food consumption on health cannot be universally established (79, 81, 86).

Growing green: how organic farming impacts the environment

Agricultural production is known as a significant contributor to many environmental problems like water pollution, soil degradation, biodiversity loss and climate change (18). There is an accepted belief that organic agriculture produces less adverse impact on environment when compared to conventional agriculture. This belief is a key rationale for governmental subsidies allocated to support the organic sector. Here, we look at the evidence of effects of organic farming on various environmental factors.

Optimizing land use

Roughly two-fifths of the ice-free land, globally, is currently under agricultural production (18). Ongoing changes in land usage, such as deforestation, are linked to many environmental challenges, including biodiversity loss and the release of soil carbon into the atmosphere. Therefore, achieving a balance between food production and environmental conservation will depend on more efficient land use and utilization of natural resources. The assessment of

environmental impacts in both organic and conventional production systems crucially relies on understanding their respective land-use requirements. Thus, impacts on the environment are evaluated per unit of land and per unit of output. The output measure reflects the lower land-use efficiency observed in organic systems. As global food demand rises, evaluating impacts per unit of output becomes increasingly relevant on a global scale. However, this approach may not fully capture the environmental impacts of widespread adoption of organic agriculture.

Currently, a small portion of global agricultural land is certified organic. Expanding organic agriculture would likely necessitate transitioning natural habitats into agricultural use. This shift in land-use would carry environmental costs that are often overlooked when evaluating the impacts based solely on output per unit (78). Thus, while measuring evaluating impacts per unit of output provides a useful metric, it may overlook broader ecological implications associated with large-scale expansion of organic farming.

Sustainable practices for energy and emissions

Food production contributes to approximately 25 % of human-caused greenhouse gas (GHG) emissions (90). These emissions, which include nitrous oxide (N_2O), carbon dioxide (CO_2) and methane (CH_4), arise from several sources such as fossil fuel combustion (used for mineral fertilizer production and agronomic machinery), animal digestion and soil chemical processes (73). Assessments of energy consumption and GHG emissions in agriculture generally encompass the production process of a product until it leaves the farm (91, 92).

Research indicates that organic farming generally requires less energy/unit of land and per unit of output than conventional farming (Fig. 6). This difference is mainly because organic systems do not use synthetic fertilizers and pesticides. Fuel consumption for agricultural activities is generally comparable in both systems. However, organic farming, particularly for crops like vegetables, may require more fuel due to the frequent use of mechanical or thermal methods for weed control (91, 93, 94). In certain situations, this can increase total energy consumption in organic systems (95). Regarding greenhouse gas (GHG) emissions, most of the studies suggest that organic farming generally has a lower impact when stated per unit of land, though not necessarily when assessed per unit of output (Fig. 6). Organic systems typically use less nitrogen, which reduces the potential for nitrous oxide (N_2O) emissions. However, achieving a balance between nutrient availability and plant requirement is often more complex in organic systems. Excess nitrogen from organic fertilizers can cause substantial N_2O emissions and insufficient nitrogen supply can lead to reduced yields (94, 96, 97).

In crop production, organic systems have been found to exhibit significantly higher soil carbon stocks and sequestration rates compared to conventional ones (98, 99). Conversely, in livestock production, less intensive organic animal husbandry practices generate more manure per unit of meat, resulting in increased methane (CH_4) and nitrous oxide (N_2O) emissions (100). Overall, the evidence does not strongly support the common belief that organic agriculture is inherently more climate-friendly than conventional farming

(101). The comparison of GHG emissions per unit land versus per unit output underscores the complexities involved in assessing the environmental impacts of different farming systems.

Minimizing nutrient loss: strategies for cleaner water

Nutrient leaching poses a significant environmental challenge, particularly in intensively farmed regions, as it contributes to water body eutrophication and marine ecosystem degradation (102). Under organic management, nitrate (NO_3^-) leaching has been observed to be lower per unit of land and not per unit of output (Fig. 6). Interestingly, organic systems tend to have higher eutrophication potential per unit of output, measured in phosphate (PO_4) equivalents, as well as more significant acidification potential, measured in sulphur dioxide (SO_2) equivalents, compared to conventional systems (97).

The absence of synthetic fertilizers in organic farming typically causes decreased leaching potential (70, 99). However, the challenge of matching nutrient supply with plant requirements in organic systems can lead to increased loss of nutrients (73, 102, 103). Regarding pesticides, prohibiting chemical pesticides in organic farming reduces the adverse effects of pollution in water systems (11). Nevertheless, certain non-chemical pesticides in organic agriculture, such as copper-based solutions in horticultural production, can also adversely affect aquatic life (70).

Soil health: the cornerstone of sustainable agriculture

Millions of hectares of land that were once fertile have become unsuitable for agriculture due to soil degradation, many times exacerbated by poor management practices (102). Organic farming practices, like adding organic matter (animal or plant

manure) and implementing extended, more varied crop rotations including cover and catch crops, are crucial in mitigating soil erosion and preventing fertility loss (70, 99). Long-term field trials' findings and meta-analyses consistently demonstrate that organically maintained fields have higher levels of organic matter and support more diverse and active soil microbial communities (Fig. 6), the key indicators of soil health.

Nature's web of life

The widening and standardization of agricultural scenarios have markedly caused the decrease in biodiversity (102, 104). A widespread consensus is that organic farming promotes greater biodiversity attributed to less pesticide use, extensive crop rotations and semi-natural landscape elements like hedges (69, 70, 77, 105–107). Meta-analyses indicate that organic farms consistently exhibit higher species richness and greater species evenness than conventional farms (refer to Fig. 6). Significant differences are particularly evident when compared with highly intensive conventional systems (99, 104). However, the biodiversity advantages associated with organic farming diminish as its intensity increases (108). Ongoing discussion exists regarding whether these benefits also decline with farm scale (99, 104, 106).

As previously discussed, expanding large-scale organic agriculture would probably result in reduced yields and further depletion of natural habitats. There is a consensus that while organic farming can enhance biodiversity, these gains are unlikely to compensate for the biodiversity losses resulting from additional changes in land use (106, 108–110). The debate between "land sharing" and "land sparing" is multifaceted and demands tailored solutions for specific locations, making

SUSTAINABILITY METRICS						
	Land use	Energy use	Greenhouse gas (GHG) emissions	Nutrient leaching/eutrophication potential	Soil quality	Biodiversity
Per unit of land	Not applicable	Energy use 10-70% lower	Nitrous oxide emissions 14-31% lower [a,c] GHG emissions 39% lower	Nitrate leaching 30-31% lower [a,c] Ammonia emissions 18% lower Phosphorus losses 1% lower [a]	Soil organic matter 6-7% higher [a,c,g] Larger and more active soil microbial communities	Species richness 30-34% higher [h,i] Organism abundance 50% higher [h] Species evenness higher [k]
Per unit of output	Land use 20-110% higher [a,b,c] Land use for organic diets 40% higher [d]	Energy use 15-21% lower [a,b]	Nitrous oxide emissions 8% higher GHG emissions 0-10% lower [a,b,c] GHG emissions of organic diet equal to conventional diet [d]	Nitrate leaching 5% lower globally and 49% higher in the EU Ammonia emissions 11% higher Acidification potential (sulphur dioxide equivalents) 13% higher [b]	Not applicable	Biodiversity loss through indirect land-use change not evaluated in available studies

Fig. 6. The environmental footprint of food: organic vs. conventional ^a104 ^b98, ^c111, ^d101, ^e141, ^f100, ^g70, ^h105, ⁱ108, ^k142. Adapted from earlier report (21).

broad global recommendations unsuitable (107, 111).

Organic farming's environmental footprint: a closer look

In general, the environmental benefits of organic agriculture are not as straightforward as commonly assumed. While organic systems often show better environmental performance per unit of land compared to conventional systems, this advantage diminishes when measured per unit of output (Fig. 6). Consequently, organic farming is more appropriate for mitigating local environmental problems such as soil degradation, than addressing global challenges like alterations in land usage and climate change (110). Some findings on environmental impacts remain inconclusive. Limited data from developing countries may restrict the global representativeness of existing results (95-98). Additionally, many studies comparing organic and conventional farming do not sufficiently control for confounding factors, which means observed differences may not necessarily indicate causal effects of organic standards. Also, it is important to recognize that significant variability in environmental effects is there within both conventional and organic systems (92, 96, 99, 100, 102, 105). Organic standards promote environmentally friendly farming techniques, such as effective nutrient management, extended crop rotations and integration of semi-natural landscape elements. However, it should be noted that these practices are also adopted by many conventional farmers (77, 94).

Science meets sustainability: technologies for optimized organic farming

Organic farming is increasingly recognized as crucial for steering towards a sustainable future, mitigating the harmful impacts of crop burning and global warming. Traditional agricultural methods heavily rely on chemical fertilizers and pesticides to address agricultural demands. However, the rise of modern organic farming arose in response to the environmental damage caused by these chemicals, offering manifold ecological advantages. Integrating organic fertilizers with cutting-edge technologies is pivotal in overcoming the constraints associated with organic farming.

Sludge: waste-to-resource solution

Sludge is a semi-solid blend generated from wastewater

treatments, industrial water, or on-site sanitation measures. It includes beneficial organic matter and key nutrients, like phosphorus and nitrogen, making it well-suited for soil amendment or organic fertilizer application. Table 2 details studies investigating sludge utilisation from wastewater treatment plants as an organic fertilizer.

The dynamic duo: biochar and organic fertilizer for thriving soil

Biochar originates from carbon-rich biomass sources, such as animal, food, industrial wastes, wood chips, sludge and agricultural and forestry residues, each exhibiting varied chemical and physical attributes. Biochar production involves several technologies, such as pyrolysis (slow, fast and intermediate), torrefaction, gasification, hydrothermal and flash carbonization. Among these, pyrolysis is the most widely used method for converting biomass into biochar (112).

Applying organic fertilizer and biochar amendments in agriculture is intended to enhance soil properties and increase crop yields. Research highlight the crucial role plant roots play in overall plant functions, noting that the addition of organic fertilizers and biochar affects root morphology and physiology, although the precise responses are not yet fully understood (113).

A previous study suggest that compost made from anaerobically digested rice straw and urea coated with rice husk biochar substantially improve soil nutrient content (114). Moreover, combinations of biochar with fertilizers, whether chemical or organic, demonstrate enhanced performance compared to using fertilizers alone, improving crop yield and enhancing plant nutrition. The use of biochar-coated chemical fertilizers also contributes to mitigating water pollution from inorganic fertilizers (114–116).

Biofertilizer: natural soil enhancer

Biofertilizers are formulations comprising of one or many types of microorganisms that enhance the availability of essential nutrients through biological procedures. These processes include soil and compost biodegradation, phosphate solubilization, nitrogen fixation and the release of cellulose or plant growth-promoting constituents into the environment. Typically, biofertilizers consist of living microbial bioinoculants, such as bacteria (e.g., free-living, nitrogen-fixing bacteria like

Table 2. Transforming waste: valuable applications of sludge

The many uses of sludge	Benefits and applications	Reference
Enhances fertilizer effectiveness	Boosts plant growth and reduces overall fertilizer needs	127
Fertilizer substitute	Comparable crop yields to conventional fertilizers	128, 129
Improves nutrient uptake in crops	Micronutrients and nitrogen supplementation for plants	130, 131
Improves soil productivity and crop yields	Enhances soil health and increases agricultural output	132
Nutrient recycling solution	Reduces waste and recovers valuable nutrients for agriculture	133
Organic fertilizer	Sustainable source of nutrients and reduces reliance on non-renewable resources	134
Potential replacement for manure	Alternative nutrient source for organic farming	135
Safe and effective fertilizer	No adverse impacts on plant growth or soil health	136
Source of phosphorus	Reduces reliance on commercial fertilizers	137
Sustainable nutrient source	Replaces commercial fertilizers and improves soil health	138
Sustainable phosphorus source	Reduces the need for mining and offers similar plant uptake efficiency as commercial fertilizers	139, 140




Fig. 7. Supercharging soil: The benefits of combining organic manure and biochar.

Azotobacter and *Rhodospirillum*), algae and fungi, either used individually or together, to naturally improve nutrient availability for plants.

Combining decomposed organic substances with plant growth-promoting microbes promotes eco-friendly practices and reduces reliance on synthetic fertilizers for crop production (Fig. 7). In an earlier study, it was reported that the usage of bio-organic fertilizer not only increased soil organic carbon by 6 - 13 % but also enhanced the population of plant growth-promoting bacteria compared to using chemical fertilizers alone (117). Furthermore, bio-organic fertilizer serves as a sustainable technology, reducing the need for Chemical N by 30 % and eliminating the requirement for TSP (triple superphosphate) in rice cultivation. Biofertilizers play a pivotal role in agriculture by lowering the cost of synthetic fertilizers and minimizing their negative impact on soil health (117). A key element of the sustainable organic farming is mycorrhizae, a symbiotic association between fungi and plants (118).

Vermicomposting harnesses earthworms to transform biodegradable waste into organic fertilizer without requiring pile turning, emitting odours and facilitates rapid compost production. The cultivation of earthworms occurs in a mixture of soil, animal wastes, decomposed leaf litter or agricultural residues. This mixture transforms into vermicompost, suitable for enhancing soil fertility for various plant types. Various approaches involving earthworms demonstrate versatile uses, particularly in aiding composting alongside microbes (119). Their study highlights effective control and reduction of microbes, pathogenic genes and greenhouse gas emissions. Additionally, soil nutrient levels are conserved and enriched. Despite these advantages, composting and vermicomposting technologies still encounter challenges in certain scenarios, prompting the need for continued investigation and discussion among upcoming researchers.

Organo-mineral nutrient supplements

Organo-mineral fertilizers are composed of both organic and

mineral elements, formulated in different ratios of nitrogen (N), phosphorus (P) and potassium (K) to suit the specific requirements of crops. The final product can be in granular, pelletized, or powdered form. These fertilizers contribute to soil fertility by enhancing its physical and chemical properties, improving nutrient availability and water retention and aiding in phosphate solubilization. Moreover, organo-mineral fertilizers promote plant agro-physiological characteristics such as higher biomass and production, improved nutrient uptake and enhanced physiological metabolism in plants (120, 121).

Web-based solutions

Digital tools and technology provide organic farmers with avenues to monitor and optimise crop and livestock health, improve efficiency in resource utilization, reduce environmental footprint and boost farm profitability. These technologies can be applied across various aspects of organic farming, like soil preparation, fertilization, irrigation, postharvest handling, storage and transportation of agricultural products. While digital advancements in agriculture are predominantly utilized by large-scale farmers and contribute significantly to sustainable practices, they also hold potential to alleviate poverty and food insecurity among smallholder farmers worldwide. Yet, digital services designed for small-scale organic farmers frequently lack long-term sustainability. Enhancements in these technologies are essential for fostering economic and environmental sustainability in organic farming (122).

Key findings

Meeting the United Nations Sustainable Development Goals (SDGs) by 2030 requires transforming agricultural practices, a topic of passionate debate. Research and discussions emphasise the need for sustainable agriculture practices and food systems that ensure adequate and wholesome food for all, minimize adverse environmental impacts and enable farmers to achieve profitability. There is consensus that agriculture and food systems need rapid adaptation to meet

multiple SDGs within ecological limits. Yet, there is intense debate over the optimal approach, centred on two main strategies: one advocating incremental improvement in conventional agriculture to enhance efficiency and reduce negative externalities and the other promoting radical restructuring towards agroecologically based organic farming systems. Integrating organic farming with improved techniques such as sludge and biochar applications, biofertilizers, organo-minerals and digital technology is crucial for overcoming the constraints of organic farming. This innovative and sustainable approach not only boosts agricultural productivity but also improves the quality of life for many farmers in an environmentally conscious manner.

Acknowledgements

The authors are thankful to their institution for its facilities and support.

Authors' contributions

MAP performed material preparation, data collection and analysis and contributed to writing the first draft of the manuscript. SG performed material preparation, data collection and analysis. MK performed material preparation, data collection and analysis. CDR performed material preparation, data collection and analysis. JK contributed to the study's conception and design, performed material preparation, data collection and analysis, wrote the first draft of the manuscript and commented on previous versions. All authors commented on previous versions of the manuscript and approved the final version.

Compliance with ethical standards

Conflict of interest: The authors do not have any conflict of interest to declare.

Ethical issues: None

References

- Funk C, Kennedy B. The new food fights: US public divides over food science. Pew Research Center. 2016;1.
- Seufert V, Ramankutty N. Many shades of gray-The context-dependent performance of organic agriculture. *Sci Adv*. 2017;3(3):e1602638. <https://doi.org/10.1126/sciadv.1602638>
- Mercati V. Organic agriculture as a paradigm of sustainability: Italian food and its progression in the global market. *Agric Sci Procedia*. 2016;8:798–802. <https://doi.org/10.1016/j.aaspro.2016.02.071>
- Klumper W, Kathage J, Qaim M. Perception of global food security issues in the German public. *Berichte Über Landwirtschaft*. 2013;91(1).
- Curtis M. Fostering economic resilience: The financial benefits of ecological farming in Kenya and Malawi. Greenpeace Africa, Johannesburg, South Africa. 2015
- Probst L, Houedjofonon E, Ayerakwa HM, Haas R. Will they buy it? The potential for marketing organic vegetables in the food vending sector to strengthen vegetable safety: A choice experiment study in three West African cities. *Food Policy*. 2012;37(3):296–308. <https://doi.org/10.1016/j.foodpol.2012.02.014>
- Connor DJ, Minguez MI. Evolution not revolution of farming systems will best feed and green the world. *Global Food Sec*. 2012;1(2):106–13. <https://doi.org/10.1016/j.gfs.2012.10.004>
- Lotter D. Facing food insecurity in Africa: Why, after 30 years of work in organic agriculture, I am promoting the use of synthetic fertilizers and herbicides in small-scale staple crop production. *Agric Human Values*. 2015;32:111–18. <https://doi.org/10.1007/s10460-014-9547-x>
- Trewavas A. Urban myths of organic farming. *Nat*. 2001;410(6827):409–10. <https://doi.org/10.1038/35068639>
- Badgley C, Moghtader J, Quintero E, Zakem E, Chappell MJ, Aviles-Vazquez K, et al. Organic agriculture and the global food supply. *Renew Agr Food Syst*. 2007;22(2):86–108. <https://doi.org/10.1017/S1742170507001640>
- Reganold JP, Wachter JM. Organic agriculture in the twenty-first century. *Nat Plants*. 2016;2(2):1–8. <https://doi.org/10.1038/nplants.2015.221>
- Evenson RE, Gollin D. Assessing the impact of the Green revolution, 1960 to 2000. *Sci*. 2003;300(5620):758–62. <https://doi.org/10.1126/science.1078710>
- Qaim M. Globalisation of agrifood systems and sustainable nutrition. *Proc Nutr Soc*. 2017;76(1):12–21. <https://doi.org/10.1017/S0029665116000598>
- UNICEF. The state of food security and nutrition in the world 2018. FAO; 2019. Available from: <https://doi.org/10.18356/c94f150c-en>
- World Health Organization. Guidelines for drinking-water quality: incorporating the first and second addenda. World Health Organization; 2022.
- Godfray HC, Beddington JR, Crute IR, Haddad L, Lawrence D, Muir JF, et al. Food security: The challenge of feeding 9 billion people. *Sci*. 2010;327(5967):812–18. <https://doi.org/10.1126/science.1185383>
- Hertel TW. The challenges of sustainably feeding a growing planet. *Food Secur*. 2015;7(2):185–98. <https://doi.org/10.1007/s12571-015-0440-2>
- Foley JA, Ramankutty N, Brauman KA, Cassidy ES, Gerber JS, Johnston M, et al. Solutions for a cultivated planet. *Nat*. 2011;478(7369):337–42. <https://doi.org/10.1038/nature10452>
- Gamage A, Gangahagedara R, Gamage J, Jayasinghe N, Kodikara N, Suraweera P, et al. Role of organic farming for achieving sustainability in agriculture. *Farming Syst*. 2023;1(1):100005. <https://doi.org/10.1016/j.farsys.2023.100005>
- Garcia SN, Osburn BI, Jay-Russell MT. One health for food safety, food security and sustainable food production. *Front Sustain Food Syst*. 2020;4:1. <https://doi.org/10.3389/fsufs.2020.00001>
- Meemken EM, Qaim M. Organic agriculture, food security and the environment. *Annu Rev Resour Econ*. 2018;10(1):39–63. <https://doi.org/10.1146/annurev-resource-100517-023252>
- Muller A, Schader C, El-Hage Scialabba N, Bruggemann J, Isensee A, Erb KH, et al. Strategies for feeding the world more sustainably with organic agriculture. *Nat Commun*. 2017;8(1):1–3. <https://doi.org/10.1038/s41467-017-01410-w>
- Kansanga M, Andersen P, Kpienbaareh D, Mason-Renton S, Atuoye K, Sano Y, et al. Traditional agriculture in transition: examining the impacts of agricultural modernization on smallholder farming in Ghana under the new Green Revolution. *Int J Sustain Dev World Ecol*. 2019;26(1):11–24. <https://doi.org/10.1080/13504509.2018.1491429>
- Liyanage CE, Thabrew MI, Kuruppuarachchi DS. Nitrate pollution in ground water of Kalpitiya: An evaluation of the content of nitrates in the water and food items cultivated in the area. *J Natn Sci Found Sri Lanka*. 2000;28(2):101–12. <https://doi.org/10.4038/jnsfsv.28i2.2679>

25. Geissen V, Silva V, Lwanga EH, Beriot N, Oostindie K, Bin Z, et al. Cocktails of pesticide residues in conventional and organic farming systems in Europe-Legacy of the past and turning point for the future. *Environ Pollut*. 2021;278:116827. <https://doi.org/10.1016/j.envpol.2021.116827>
26. Vogt G. The origins of organic farming. In: Lockeretz W, editors. *Organic farming: An international history*; Wallingford UK:CABI; 2007. p. 9–29 <https://doi.org/10.1079/9780851998336.0009>
27. Lockeretz W. What explains the rise of organic farming?. In: Lockeretz W, editors. *Organic farming: An international history*; Wallingford UK:CABI; 2007. p. 1–8. <https://doi.org/10.1079/9780851998336.0001>
28. Stolze M, Lampkin N. Policy for organic farming: rationale and concepts. *Food Policy*. 2009;34(3):237–44. <https://doi.org/10.1016/j.foodpol.2009.03.005>
29. Clark S. Organic farming and climate change: The need for innovation. *Sustain*. 2020;12(17):7012. <https://doi.org/10.3390/su12177012>
30. Hammed TB, Oloruntoba EO, Ana GR. Enhancing growth and yield of crops with nutrient-enriched organic fertilizer at wet and dry seasons in ensuring climate-smart agriculture. *Int J Recycl Org Waste Agric*. 2019;8:81–92. <https://doi.org/10.1007/s40093-019-0274-6>
31. Pandiselvi T, Jeyajothi R, Kandeshwari M. Organic nutrient management a way to improve soil fertility and sustainable agriculture- A review. *Int J Adv Life Sci*. 2017;10(2):175–81. <https://doi.org/10.26627/ijals/2017/10.02.0005>
32. Ebitu L, Avery H, Mourad KA, Enyetu J. Citizen science for sustainable agriculture- A systematic literature review. *Land use policy*. 2021;103:105326. <https://doi.org/10.1016/j.landusepol.2021.105326>
33. Merrigan K, Giraud EG, Scialabba NE, Brook L, Johnson A, Aird NS. Grow organic: The climate, health and economic case for expanding organic agriculture. The Natural Resources Defence Council. 2022.
34. Muller A. Benefits of organic agriculture as a climate change adaptation and mitigation strategy in developing countries. rapport nr.: Working Papers in Economics 343. 2009. <https://doi.org/10.1088/1755-1307/6/7/372032>
35. Pandit MA, Kumar J, Gulati S, Bhandari N, Mehta P, Katyal R, et al. Major biological control strategies for plant pathogens. *Pathog*. 2022;11(2):273. <https://doi.org/10.3390/pathogens11020273>
36. Seneviratne G, Kulasooriya SA, Rosswall T. Sustainment of soil fertility in the traditional rice farming, dry zone, Sri Lanka. *Soil Biol Biochem*. 1994;26(6):681–88. [https://doi.org/10.1016/0038-0717\(94\)90259-3](https://doi.org/10.1016/0038-0717(94)90259-3)
37. Seufert V, Ramankutty N, Mayerhofer T. What is this thing called organic? - How organic farming is codified in regulations. *Food Policy*. 2017;68:10–20. <https://doi.org/10.1016/j.foodpol.2016.12.009>
38. Jena PR, Chichaibelu BB, Stellmacher T, Grote U. The impact of coffee certification on small-scale producers' livelihoods: A case study from the Jimma Zone, Ethiopia. *Agric Econ*. 2012;43(4):429–40. <https://doi.org/10.1111/j.1574-0862.2012.00594.x>
39. Meemken EM, Veettil PC, Qaim M. Toward improving the design of sustainability standards- A gendered analysis of farmers' preferences. *World Dev*. 2017;99:285–98. <https://doi.org/10.1016/j.worlddev.2017.05.021>
40. Willer H, Travnicek J, Schlatter S. The world of organic agriculture. Statistics and emerging trends 2024. <https://doi.org/10.1017/s0014479709007807>
41. Carlson A, Jaenicke E. Changes in retail organic price premiums from 2004 to 2010. USDA; 2016. Available from: <https://www.ers.usda.gov/publications/pub-details?pubid=45550>
42. Brenes-Munoz T, Lakner S, Bruummer B. What influences the growth of organic farms? Evidence from a panel of organic farms in Germany. *German J Agric Econ*. 2016;65(1):1–5. <https://doi.org/10.52825/gjae.v65i1.2010>
43. Serra T, Zilberman D, Gil JM. Differential uncertainties and risk attitudes between conventional and organic producers: The case of Spanish arable crop farmers. *Agric Econ*. 2008;39(2):219–29. <https://doi.org/10.1111/j.1574-0862.2008.00329.x>
44. Kallas Z, Serra T, Gil JM. Farmers' objectives as determinants of organic farming adoption: The case of Catalanian vineyard production. *Agric Econ*. 2010;41(5):409–23. <https://doi.org/10.1111/j.1574-0862.2010.00454.x>
45. Lapple D. Adoption and abandonment of organic farming: An empirical investigation of the Irish drystock sector. *J Agric Econ*. 2010;61(3):697–714. <https://doi.org/10.1111/j.1477-9552.2010.00260.x>
46. Wollni M, Andersson C. Spatial patterns of organic agriculture adoption: evidence from Honduras. *Ecol Econ*. 2014;97:120–28. <https://doi.org/10.1016/j.ecolecon.2013.11.010>
47. Kuminoff NV, Wossink A. Why isn't more US farmland organic?. *J Agric Econ*. 2010;61(2):240–58. <https://doi.org/10.1111/j.1477-9552.2009.00235.x>
48. Bolwig S, Gibbon P, Jones S. The economics of smallholder organic contract farming in tropical Africa. *World Dev*. 2009;37(6):1094–104. <https://doi.org/10.1016/j.worlddev.2008.09.012>
49. Walmsley A, Sklenicka P. Various effects of land tenure on soil biochemical parameters under organic and conventional farming - implications for soil quality restoration. *Ecol Eng*. 2017;107:137–43. <https://doi.org/10.1016/j.ecoleng.2017.07.006>
50. Malik DS, Sharma AK, Sharma AK, Thakur R, Sharma M. A review on impact of water pollution on freshwater fish species and their aquatic environment. *Adv Environ Pollut Manag*. 2020;1:10–28. <https://doi.org/10.26832/aesa-2020-aepm-02>
51. Mateo-Sagasta J, Zadeh SM, Turrall H, Burke J. Water pollution from agriculture: A global review. Executive Summary. Rome: FAO; 2017.
52. Chaichi W, Djazouli Z, Zebib B, Merah O. Effect of vermicompost tea on faba bean growth and yield. *Compost Sci Util*. 2018;26(4):279–85. <https://doi.org/10.1080/1065657x.2018.1528908>
53. Merah O, Djazouli ZE, Zebib B. Aqueous extract of algerian nettle (*Urtica dioica* L.) as possible alternative pathway to control some plant diseases. *Iran J Sci Technol Trans A: Sci*. 2021;45:463–68. <https://doi.org/10.1007/s40095-021-01061-z>
54. Mohammedi A, Zebib B, Atika S, Merah O, Djazouli ZE. Effect of *Ulva rigida* as crude and formulated extract on vineyard growth at an early stage. *Ukr J Ecol*. 2022;12(6):15–25.
55. Zhou X, Ding D. Factors influencing farmers' willingness and behaviors in organic agriculture development: An empirical analysis based on survey data of farmers in Anhui Province. *Sustain*. 2022;14(22):14945. <https://doi.org/10.3390/su142214945>
56. Food and Agriculture Organization of the United Nations. Transforming food and agriculture to achieve the Sustainable Development Goals (SDGs): Good practices from FAO-GEF projects around the world. Rome, FAO. 2020. p. 64. <https://doi.org/10.4060/ca8768en>
57. Pingali PL. Green revolution: impacts, limits and the path ahead. *Proc Natl Acad Sci*. 2012;109(31):12302–08. <https://doi.org/10.1073/pnas.0912953109>
58. Pretty J, Benton TG, Bharucha ZP, Dicks LV, Flora CB, Godfray HC, et al. Global assessment of agricultural system redesign for sustainable intensification. *Nat Sustain*. 2018;1(8):441–46. <https://doi.org/10.1038/s41893-018-0114-0>
59. Baker BP, Cooley D, Futrell S, Garling L, Gershuny G, Green TA, et al. Organic agriculture and integrated pest management:

- synergistic partnership needed to Improve the Sustainability of Agriculture and Food Systems. In: Brian PB, Thomas AG, editors. National Organic and IPM working group. 2015.
60. Merah O, Djazouli Z. Toxicity Evaluation of *Dittrichia viscosa* L's aqueous extracts in combination with bio-adjuvant *Silene fuscata* on *Chaitophorus leucomelas* Koch. (Hom., Aphididae) and on biocenotic resumption of functional groups. Jordan J Agric Sci. 2016;12(3):797–814. <https://doi.org/10.12816/0033386>
 61. Salim D, De Caro P, Merah O, Chbani A. Control of postharvest citrus green mold using *Ulva lactuca* extracts as a source of active substances. Int J Bio-resource Stress Manag. 2020;11(3):287–96. <https://doi.org/10.23910/1.2020.2107>
 62. Cassman KG. Editorial response by Kenneth Cassman: can organic agriculture feed the world-science to the rescue?. Renew Agric Food Syst. 2007;22(2):83–84.
 63. Connor DJ. Organic agriculture cannot feed the world. Field Crops Res. 2008;106(2):187–90. <https://doi.org/10.1016/j.fcr.2007.11.010>
 64. Goulding KW, Trewavas AJ, Giller KE. Can organic farming feed the world? A contribution to the debate on the ability of organic farming systems to provide sustainable supplies of food. In: Proceedings of the International Fertiliser Society 663. Paris: IFS; 2009.
 65. De Ponti T, Rijk B, Van Ittersum MK. The crop yield gap between organic and conventional agriculture. Agric Syst. 2012;108:1–9. <https://doi.org/10.1016/j.agsy.2011.12.004>
 66. Ponisio LC, M'Gonigle LK, Mace KC, Palomino J, De Valpine P, Kremen C. Diversification practices reduce organic to conventional yield gap. Proc R Soc B: Biol Sci. 2015;282(1799):20141396. <https://doi.org/10.1098/rspb.2014.1396>
 67. Seufert V, Ramankutty N, Foley JA. Comparing the yields of organic and conventional agriculture. Nat. 2012;485(7397):229–32. <https://doi.org/10.1038/nature11069>
 68. Forster D, Andres C, Verma R, Zundel C, Messmer MM, Mader P. Yield and economic performance of organic and conventional cotton-based farming systems—results from a field trial in India. PloS One. 2013;8(12):e81039. <https://doi.org/10.1371/journal.pone.0081039>
 69. Mader P, Fliessbach A, Dubois D, Gunst L, Fried P, Niggli U. Soil fertility and biodiversity in organic farming. Sci. 2002;296(5573):1694–97. <https://doi.org/10.1126/science.1071148>
 70. Niggli U. Sustainability of organic food production: challenges and innovations. Proc Nutr Soc. 2015;74(1):83–88. <https://doi.org/10.1017/s0029665114001438>
 71. Berry PM, Sylvester-Bradley R, Philipps L, Hatch DJ, Cuttle SP, Rayns FW, et al. Is the productivity of organic farms restricted by the supply of available nitrogen?. Soil Use Manag. 2002;18(1):248–55. <https://doi.org/10.1079/sum2002129>
 72. Oehl F, Oberson A, Tagmann HU, Besson JM, Dubois D, Mader P, et al. Phosphorus budget and phosphorus availability in soils under organic and conventional farming. Nutr Cycling Agroecosyst. 2002;62:25–35. <https://doi.org/10.1023/a:1015195023724>
 73. Gomiero T, Pimentel D, Paoletti MG. Environmental impact of different agricultural management practices: conventional vs. organic agriculture. Crit Rev Plant Sci. 2011;30(1-2):95–124. <https://doi.org/10.1080/07352689.2011.554355>
 74. Kravchenko AN, Snapp SS, Robertson GP. Field-scale experiments reveal persistent yield gaps in low-input and organic cropping systems. Proc Natl Acad Sci. 2017;114(5):926–31. <https://doi.org/10.1073/pnas.1612311114>
 75. Taheri F, Azadi H, D'Haese M. A world without hunger: organic or GM crops?. Sustain. 2017;9(4):580. <https://doi.org/10.3390/su9040580>
 76. Kniss AR, Savage SD, Jabbour R. Commercial crop yields reveal strengths and weaknesses for organic agriculture in the United States. PloS One. 2016;11(8):e0161673. <https://doi.org/10.1371/journal.pone.0161673>
 77. Pimentel D, Burgess M. An environmental, energetic and economic comparison of organic and conventional farming systems. Integrated pest management: pesticide problems. 2014;3. p. 141–66. https://doi.org/10.1007/978-94-007-7796-5_6
 78. Leifeld J. Current approaches neglect possible agricultural cutback under large-scale organic farming. A comment to Ponisio et al. Proc R Soc B: Biol Sci. 2016;283(1824):20151623. <https://doi.org/10.1098/rspb.2015.1623>
 79. Dangour AD, Lock K, Hayter A, Aikenhead A, Allen E, Uauy R. Nutrition-related health effects of organic foods: A systematic review. Am J Clin Nutr. 2010;92(1):203–10. <https://doi.org/10.3945/ajcn.2010.29269>
 80. Huber M, Knottnerus JA, Green L, Van Der Horst H, Jadad AR, Kromhout D, et al. How should we define health?. Br Med J. 2011;343:d4163. <https://doi.org/10.1136/bmj.d4163>
 81. Baranski M, Srednicka-Tober D, Volakakis N, Seal C, Sanderson R, Stewart GB, et al. Higher antioxidant and lower cadmium concentrations and lower incidence of pesticide residues in organically grown crops: A systematic literature review and meta-analyses. Br J Nutr. 2014;112(5):794–811. <https://doi.org/10.1017/s0007114514001366>
 82. Magkos F, Arvaniti F, Zampelas A. Organic food: buying more safety or just peace of mind? A critical review of the literature. Crit Rev Food Sci Nutr. 2006;46(1):23–56. <https://doi.org/10.1080/10408690490911846>
 83. Smith-Spangler C, Brandeau ML, Hunter GE, Bavinger JC, Pearson M, Eschbach PJ, et al. Are organic foods safer or healthier than conventional alternatives? A systematic review. Ann Intern Med. 2012;157(5):348–66. <https://doi.org/10.7326/0003-4819-157-5-201209040-00007>
 84. Mditshwa A, Magwaza LS, Tesfay SZ, Mbili N. Postharvest quality and composition of organically and conventionally produced fruits: A review. Sci Hortic. 2017;216:148–59. <https://doi.org/10.1016/j.scienta.2016.12.033>
 85. Brandt K, Leifert C, Sanderson R, Seal CJ. Agroecosystem management and nutritional quality of plant foods: The case of organic fruits and vegetables. Crit Rev Plant Sci. 2011;30(1-2):177–97. <https://doi.org/10.1080/07352689.2011.554417>
 86. Forman J, Silverstein J. Organic Foods: Health and Environmental Advantages and Disadvantages. Pediatric Clinical Practice Guidelines & Policies [Internet]. 2013;130(5):651–62. Available from: http://doi.org/10.1542/9781581108224-organic_sub01
 87. Alfven T, Braun-Fahrlander C, Brunekreef BV, Von Mutius E, Riedler J, Scheynius A, et al. Allergic diseases and atopic sensitization in children related to farming and anthroposophic lifestyle—the PARSIFAL study. Allergy. 2006;61(4):414–21. <https://doi.org/10.1111/j.1398-9995.2005.00939.x>
 88. Kummeling I, Thijs C, Huber M, van de Vijver LP, Snijders BE, Penders J, et al. Consumption of organic foods and risk of atopic disease during the first 2 years of life in the Netherlands. Br J Nutr. 2008;99(3):598–605. <https://doi.org/10.1017/s0007114507815844>
 89. Kesse-Guyot E, Peneau S, Mejean C, Szabo de Edelenyi F, Galan P, Hercberg S, et al. Profiles of organic food consumers in a large sample of French adults: results from the Nutrinet-Sante cohort study. PloS One. 2013;8(10):e76998. <https://doi.org/10.1371/journal.pone.0076998>
 90. Edenhofer O, Pichs-Madruga R, Sokona Y, Matshoss P, Seyboth K. Renewable energy sources and climate change mitigation. UK: Cambridge University Press; 2011. <https://doi.org/10.1017/cbo9781139151153>
 91. Smith LG, Williams AG, Pearce BD. The energy efficiency of organic agriculture: A review. Renew Agr Food Syst. 2015;30(3):280–301.

- <https://doi.org/10.1017/s1742170513000471>
92. Meier MS, Stoessel F, Jungbluth N, Juraske R, Schader C, Stolze M. Environmental impacts of organic and conventional agricultural products– are the differences captured by life cycle assessment?. *J Environ Manag.* 2015;149:193–208. <https://doi.org/10.1016/j.jenvman.2014.10.006>
 93. Scialabba NE, Muller-Lindenlauf M. Organic agriculture and climate change. *Renew Agr Food Syst.* 2010;25(2):158–69. <https://doi.org/10.1017/s1742170510000116>
 94. Lynch DH, MacRae R, Martin RC. The carbon and global warming potential impacts of organic farming: does it have a significant role in an energy constrained world?. *Sustain.* 2011;3(2):322–62. <https://doi.org/10.3390/su3020322>
 95. Lee KS, Choe YC, Park SH. Measuring the environmental effects of organic farming: A meta-analysis of structural variables in empirical research. *J Environ Manag.* 2015;162:263–74. <https://doi.org/10.1016/j.jenvman.2015.07.021>
 96. Skinner C, Gattinger A, Muller A, Mader P, Fließbach A, Stolze M, et al. Greenhouse gas fluxes from agricultural soils under organic and non-organic management-A global meta-analysis. *Sci Total Environ.* 2014;468:553–63. <https://doi.org/10.1016/j.scitotenv.2013.08.098>
 97. Clark M, Tilman D. Comparative analysis of environmental impacts of agricultural production systems, agricultural input efficiency and food choice. *Environ Res Lett.* 2017;12(6):064016. <https://doi.org/10.1088/1748-9326/aa6cd5>
 98. Gattinger A, Muller A, Haeni M, Skinner C, Fließbach A, Buchmann N, et al. Enhanced top soil carbon stocks under organic farming. *Proc Natl Acad Sci.* 2012;109(44):18226–31. <https://doi.org/10.1073/pnas.1209429109>
 99. Lori M, Symnaccik S, Mader P, De Deyn G, Gattinger A. Organic farming enhances soil microbial abundance and activity- A meta-analysis and meta-regression. *PLoS One.* 2017;12(7):e0180442. <https://doi.org/10.1371/journal.pone.0180442>
 100. Treu H, Nordborg M, Cederberg C, Heuer T, Claupein E, Hoffmann H, et al. Carbon footprints and land use of conventional and organic diets in Germany. *J Clean Prod.* 2017;161:127–42. <https://doi.org/10.1016/j.jclepro.2017.05.041>
 101. Beirat WB. Klimaschutz in der land-und forstwirtschaft sowie den nachgelagerten bereichen ernahrung und holzverwendung. *Berichte über Landwirtschaft-Zeitschrift für Agrarpolitik und Landwirtschaft.* 2016.
 102. Halberg N. Assessment of the environmental sustainability of organic farming: definitions, indicators and the major challenges. *Can J Plant Sci.* 2012;92(6):981–96. <https://doi.org/10.4141/cjps2012-035>
 103. Tuomisto HL, Hodge ID, Riordan P, Macdonald DW. Does organic farming reduce environmental impacts?– A meta-analysis of European research. *J Environ Manag.* 2012;112:309–20. <https://doi.org/10.1016/j.jenvman.2012.08.018>
 104. Bengtsson J, Ahnstrom J, Weibull AC. The effects of organic agriculture on biodiversity and abundance: A meta-analysis. *J Appl Ecol.* 2005;42(2):261–69. <https://doi.org/10.1111/j.1365-2664.2005.01005.x>
 105. Hole DG, Perkins AJ, Wilson JD, Alexander IH, Grice PV, Evans AD. Does organic farming benefit biodiversity?. *Biol Conserv.* 2005;122(1):113–30. <https://doi.org/10.1016/j.biocon.2004.07.018>
 106. Schneider MK, Luscher G, Jeanneret P, Arndorfer M, Ammari Y, Bailey D, et al. Gains to species diversity in organically farmed fields are not propagated at the farm level. *Nat Commun.* 2014;5(1):4151. <https://doi.org/10.1038/ncomms5151>
 107. Tuck SL, Winqvist C, Mota F, Ahnstrom J, Turnbull LA, Bengtsson J. Land-use intensity and the effects of organic farming on biodiversity: A hierarchical meta-analysis. *J Appl Ecol.* 2014;51(3):746–55. <https://doi.org/10.1111/1365-2664.12219>
 108. Gabriel D, Sait SM, Kunin WE, Benton TG. Food production vs. biodiversity: comparing organic and conventional agriculture. *J Appl Ecol.* 2013;50(2):355–64. <https://doi.org/10.1111/1365-2664.12035>
 109. Green RE, Cornell SJ, Scharlemann JP, Balmford A. Farming and the fate of wild nature. *Sci.* 2005;307(5709):550–55. <https://doi.org/10.1126/science.1106049>
 110. Mondelaers K, Aertsens J, Van Huylenbroeck G. A meta-analysis of the differences in environmental impacts between organic and conventional farming. *Brit Food J.* 2009;111(10):1098–119. <https://doi.org/10.1108/00070700910992925>
 111. Phalan B, Onial M, Balmford A, Green RE. Reconciling food production and biodiversity conservation: land sharing and land sparing compared. *Sci.* 2011;333(6047):1289–91. <https://doi.org/10.1126/science.1208742>
 112. Kukreti A, Kurmanchali N, Rawat. Organic farming and biochar. In: Uniyal A, Sharma I, editors. *Trends in agriculture: Traditional and modern approaches.* Jodhpur Agrobios; 2021. p. 141–153.
 113. Zhang Z, Dong X, Wang S, Pu X. Benefits of organic manure combined with biochar amendments to cotton root growth and yield under continuous cropping systems in Xinjiang, China. *Sci Rep.* 2020;10(1):4718. <https://doi.org/10.1038/s41598-020-61118-8>
 114. Gamage A, Basnayake B, De Costa J, Merah O. Effects of rice husk biochar coated urea and anaerobically digested rice straw compost on the soil fertility and cyclic effect of phosphorus. *Plants.* 2021;11(1):75. <https://doi.org/10.3390/plants11010075>
 115. Agu OS, Tabil LG, Mupondwa E, Emadi B, Dumonceaux T. Impact of biochar addition in microwave torrefaction of camelina straw and switchgrass for biofuel production. *Fuels.* 2022;3(4):588–606. <https://doi.org/10.3390/fuels3040036>
 116. Glaser B, Wiedner K, Seelig S, Schmidt HP, Gerber H. Biochar organic fertilizers from natural resources as substitute for mineral fertilizers. *Agron Sustain Dev.* 2015;35:667–78. <https://doi.org/10.1007/s13593-014-0251-4>
 117. Kumar R, Kumawat N, Sahu YK. Role of biofertilizers in agriculture. *Popular Khedi.* 2017;5(4):63–66.
 118. Kunal, Sharma S, Gosal SK, Choudhary R, Singh R, Adholeya A. Optical sensing and arbuscular mycorrhizal fungi for improving fertilizer nitrogen and phosphorus use efficiencies in maize. *J Soil Sci Plant Nutr.* 2020;20:2087–98. <https://doi.org/10.1007/s42729-020-00277-z>
 119. Zhou Y, Liu H, Chen H, Awasthi SK, Sindhu R, Binod P, et al. Introduction: trends in composting and vermicomposting technologies. *Curr Dev Biotechnol Bioeng.* 2023:1–28. <https://doi.org/10.1016/b978-0-323-91874-9.00004-8>
 120. Atere CT, Olayinka A. Effect of organo-mineral fertilizer on soil chemical properties, growth and yield of soybean. *Afr J Agric Res.* 2012;7(37):5208–16. <https://doi.org/10.5897/ajar11.1378>
 121. Crusciol CA, Campos MD, Martello JM, Alves CJ, Nascimento CA, Pereira JC, et al. Organomineral fertilizer as source of P and K for sugarcane. *Sci Rep.* 2020;10(1):5398. <https://doi.org/10.1038/s41598-020-62315-1>
 122. Mushi GE, Di Marzo Serugendo G, Burgi PY. Digital technology and services for sustainable agriculture in Tanzania: A literature review. *Sustain.* 2022;14(4):2415. <https://doi.org/10.3390/su14042415>
 123. World Health Organization. The state of food security and nutrition in the world 2022: repurposing food and agricultural policies to make healthy diets more affordable. *Food Agri Org.* 2022. <https://doi.org/10.4060/cc0639en>
 124. Seremesic S, Dolijanovic Z, Tomas Simin M, Vojnov B, Glavas Trbic D. The future we want: sustainable development goals accomplishment with organic agriculture. *Probl Ekorozw.* 2021;16(2):171–80. <https://doi.org/10.35784/pe.2021.2.18>

125. Setboonsarng S, Gregorio EE. Achieving sustainable development goals through organic agriculture: empowering poor women to build the future. Manila: Asian Development Bank; 2017. <https://doi.org/10.22617/wps179123-2>
126. Markandya A, Setboonsarng S, editors. Organic agriculture and post-2015 development goals: Building on the comparative advantage of poor farmers. Manila: Asian Development Bank; 2015.
127. Bouhia Y, Hafidi M, Ouhdouch Y, Zeroual Y, Lyamlouli K. Organo-mineral fertilization based on olive waste sludge compost and various phosphate sources improves phosphorus agronomic efficiency, *Zea mays* agro-physiological traits and water availability. *Agron*. 2023;13(1):249. <https://doi.org/10.3390/agronomy13010249>
128. Rathod PH, Patel JC, Shah MR, Jhala AJ. Recycling gamma irradiated sewage sludge as fertilizer: A case study using onion (*Alium cepa*). *Appl Soil Ecol*. 2009;41(2):223–33. <https://doi.org/10.1016/j.apsoil.2008.11.001>
129. Kominko H, Gorazda K, Wzorek Z. The possibility of organo-mineral fertilizer production from sewage sludge. *Waste Biomass Valori*. 2017;8:1781–91. <https://doi.org/10.1007/s12649-016-9805-9>
130. Koutroubas SD, Antoniadis V, Damalas CA, Fotiadis S. Municipal sewage sludge effects on maize yield, nitrogen use efficiency and soil properties. *J Soil Sci Plant Nutr*. 2023;23(1):1209–21. <https://doi.org/10.1007/s42729-022-01115-0>
131. Correa RS, Barbosa JZ, Poggere GC, Magri E, de Oliveira SA. Grain and foliar nutritional responses of corn (*Zea mays* L.) to sewage sludge soil application. *Waste Biomass Valori*. 2023;14(8):2629–41. <https://doi.org/10.1007/s12649-023-02037-3>
132. Jamil M, Qasim M, Umar M. Utilization of sewage sludge as organic fertilizer in sustainable agriculture. *J Appl Sci*. 2006;6(3):531–35. <https://doi.org/10.3923/jas.2006.531.535>
133. Kominko H, Gorazda K, Wzorek Z. Potentiality of sewage sludge-based organo-mineral fertilizer production in Poland considering nutrient value, heavy metal content and phytotoxicity for rapeseed crops. *J Environ Manag*. 2019;248:109283. <https://doi.org/10.1016/j.jenvman.2019.109283>
134. Chojnacka K, Skrzypczak D, Szopa D, Izydorczyk G, Moustakas K, Witek-Krowiak A. Management of biological sewage sludge: fertilizer nitrogen recovery as the solution to fertilizer crisis. *J Environ Manag*. 2023;326:116602. <https://doi.org/10.1016/j.jenvman.2022.116602>
135. Kumar A, Kumar A, Kumar Yadav A, Thakur S. Impact of organic farming on eco-friendly sustainable agriculture: A review. *Pharma Innov J*. 2022;11(2):709–15.
136. Rathod PH, Patel JC, Jhala AJ. Potential of gamma irradiated sewage sludge as fertilizer in radish: evaluating heavy-metal accumulation in sandy loam soil. *Commun Soil Sci Plant Anal*. 2011;42(3):263–82. <https://doi.org/10.1080/00103624.2011.538880>
137. Sichler TC, Adam C, Montag D, Barjenbruch M. Future nutrient recovery from sewage sludge regarding three different scenarios-German case study. *J Clean Prod*. 2022;333:130130. <https://doi.org/10.1016/j.jclepro.2021.130130>
138. Banerjee DS, Balkrishna A, Arya V, Ghosh S, Singh SK. Reuse of sewage sludge as organic agricultural products: An efficient technology-based initiative. *Authorea Preprints*. 2022. <https://doi.org/10.22541/au.164899210.06660033/v1>
139. Franz M. Phosphate fertilizer from sewage sludge ash (SSA). *Waste Manag*. 2008;28(10):1809–18. <https://doi.org/10.1016/j.wasman.2007.08.011>
140. Herzel H, Kruger O, Hermann L, Adam C. Sewage sludge ash-A promising secondary phosphorus source for fertilizer production. *Sci Total Environ*. 2016;542:1136–43. <https://doi.org/10.1016/j.scitotenv.2015.08.059>
141. Gomiero T, Paoletti MG, Pimentel D. Energy and environmental issues in organic and conventional agriculture. *Crit Rev Plant Sci*. 2008;27(4):239–54. <https://doi.org/10.1080/0735268080225456>
142. Crowder DW, Northfield TD, Strand MR, Snyder WE. Organic agriculture promotes evenness and natural pest control. *Nat*. 2010;466(7302):109–12. <https://doi.org/10.1038/nature09183>

Additional information

Peer review: Publisher thanks Sectional Editor and the other anonymous reviewers for their contribution to the peer review of this work.

Reprints & permissions information is available at https://horizonpublishing.com/journals/index.php/PST/open_access_policy

Publisher's Note: Horizon e-Publishing Group remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Indexing: Plant Science Today, published by Horizon e-Publishing Group, is covered by Scopus, Web of Science, BIOSIS Previews, Clarivate Analytics, NAAS, UGC Care, etc
See https://horizonpublishing.com/journals/index.php/PST/indexing_abstracting

Copyright: © The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited (<https://creativecommons.org/licenses/by/4.0/>)

Publisher information: Plant Science Today is published by HORIZON e-Publishing Group with support from Empirion Publishers Private Limited, Thiruvananthapuram, India.