

RESEARCH ARTICLE

Climate change linked spatio-temporal drought prediction over Tamil Nadu (India)

Vengateswari Mahalingam¹, Bhuvaneswari Kulanthaivel², Geethalakshmi Vellingiri^{2*}, Arul Prasad Sivaprakasam², Sathyapriya Eswaran³, Guna Mani⁴, Dharani Chelladurai⁵, Balaji Thiruvalliparam⁶ & Vigneswaran Selvaraj⁷

¹Department of Agricultural Engineering, Mohamed Sathak Engineering College, Ramanathapuram 623 806, India

²Agro Climate Research Centre, Tamil Nadu Agricultural University, Coimbatore 641 003, India

³Agricultural Extension, Amrita School of Agricultural Sciences, Amrita Vishwa Vidyapeetham University, Coimbatore 642 109, India

⁴North Karnataka Agrometeorological Forecasting and Research Centre, India Meteorological Department, Dharwad 580 005, India

⁵Department of Agronomy, SRM College of Agricultural Sciences, Chengalpattu 603 201, India

⁶Soil Science and Agricultural Chemistry, Agricultural College and Research Institute, Vazhavachanur 606 753, India

⁷Division of Genetics and Tree Breeding, Institute of Forest Genetics and Tree Breeding, Coimbatore 641 002, India

*Email: geetha@tnau.ac.in



ARTICLE HISTORY

Received: 07 December 2024

Accepted: 12 February 2025

Available online

Version 1.0 : 11 March 2025

Version 2.0 : 01 April 2025



Additional information

Peer review: Publisher thanks Sectional Editor and the other anonymous reviewers for their contribution to the peer review of this work.

Reprints & permissions information is available at https://horizonepublishing.com/journals/index.php/PST/open_access_policy

Publisher's Note: Horizon e-Publishing Group remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Indexing: Plant Science Today, published by Horizon e-Publishing Group, is covered by Scopus, Web of Science, BIOSIS Previews, Clarivate Analytics, NAAS, UGC Care, etc See https://horizonepublishing.com/journals/index.php/PST/indexing_abstracting

Copyright: © The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited (<https://creativecommons.org/licenses/by/4.0/>)

CITE THIS ARTICLE

Mahalingam V, Kulanthaivel B, Vellingiri G, Sivaprakasam AP, Eswaran S, Mani G, Chelladurai D, Thiruvalliparam B, Selvaraj V. Climate change linked spatio-temporal drought prediction over Tamil Nadu (India). Plant Science Today. 2025; 12 (2): 1-6. <https://doi.org/10.14719/pst.6584>

Abstract

Climate change projections suggest an increased risk of extreme events, potentially accelerating the occurrence of droughts in the future. Drought is a recurring climatic phenomenon that significantly affects the social and economic development of agricultural countries. It can occur in many parts of the world, though its characteristics differ from one region to another. The study aims to analyze future drought events in Tamil Nadu to gain insights into their frequency and spatial extent. Dynamical downscaling of the global circulation model (CCSM4) was done by employing the regional climate model (RCM) RegCM4.4. The downscaled future rainfall data was used to calculate standardized precipitation index and drought characterization in the future period such as the near century (2010-2039), mid-century (2040-2069) and end of the century (2071-2099) under RCP 4.5 and RCP 8.5 scenario of CMIP5 project. The results of calculated SPI values showed that the drought frequency was higher in the Southwest Monsoon (20%) compared to the Northeast Monsoon in the future. Based on SPI, drought occurrence is expected to increase in the 4.5 scenario (50%) as compared to the 8.5 scenario, drought occurrence is projected to decrease due to an increase in rainfall projected in mid-century. This article paves the way for more accurate drought management strategies, contributing to improved resilience and sustainable development in drought-prone regions.

Keywords

climate change; drought; future; frequency; prediction

Introduction

Drought is one of the most dangerous natural disasters, occurring in almost all geographical areas and it is a complex phenomenon that is one of the least understood natural hazards because of the diverse causal factors or contributing elements that operate at various temporal and spatial scales (1). Drought had extensive impacts on regional and global scales in recent decades necessitating increased drought resilience (2). Increased CO₂ emissions from anthropogenic activities in recent decades triggered


limited by its inherent nature of dependence on the mean. The standardized precipitation index (SPI) expresses the actual rainfall as a standardized departure from the rainfall probability distribution function. As a result, this index has become important in recent years as a potential drought indicator permitting space and time comparisons (11) and assigned as a suitable tool for analysing drought for different hydrological purposes over different temporal and spatial variations (12).

The study could utilize climate models to project future drought patterns under various greenhouse gas emission scenarios, providing insights into the potential impacts of different climate change mitigation and adaptation strategies, tailored to the specific needs and vulnerabilities of Tamil Nadu.

Description of the study area

Climate projections

Drought indices are calculated as a single numerical value to characterize drought viz., severity and extent of drought and the values of drought indices are used for evaluating the drought hazards and making decisions on declarations of drought-affected areas (9). A drought index calculated from known values of selected parameters enables the description of drought to be expressed quantitatively (10). Rainfall deviation from the long-term mean continues to be a widely adopted indicator for drought intensity assessment because of its simplicity. However, the application of this indicator is strongly



<https://plantsciencetoday.online>

National Command Language (NCL)

After the simulation completion of the WRF model, National Command Language (NCL) version 6.3.1 was used to extract the rainfall data required for this study. A separate shell script was written for arranging the data into columns and to respective number of days in text format for further analysis.

Standardized precipitation index (SPI)

SPI was developed in Colorado to quantify the scarcity of precipitation over multiple time scales highlighting the impact of drought on the availability of the various water resources (14). The calculation of the SPI is based on the long-term record of precipitation. Data for SPI use are recommended for at least 30 consecutive years. Using the SPI, the study showed the relationship between drought frequency, drought duration and drought time scale.

$$SPI = \frac{X - \bar{X}}{\sigma} \quad \text{Eqn. 1}$$

where,

X = mean precipitation for the i^{th} station

\bar{X} = precipitation for the i^{th} station and k^{th} observation

σ = standardized deviation for the i^{th} station

SPI on a seasonal and annual basis was calculated at the district level for future. In the interpretation of SPI, the positive values indicate greater values than the median precipitation while negative values indicate lesser values than the median precipitation. The classification of SPI values as shown in Table 1 describes drought intensities.

Table 1. Classification of SPI values

SPI range	Drought intensity class
>2	Extremely wet
1.5 to 1.99	Very wet
1.0 to 1.49	Moderately wet
0.5 to 0.99	Mild wet
0.5 to -0.5	Near normal
-0.5 to 0.99	Mild drought
-1.0 to -1.49	Moderately dry
-1.5 to -1.99	Severe dry
-2 and less	Extremely dry

Results

Projected changes in drought frequency

Drought frequency varies significantly across Tamil Nadu due to a complex interplay of geographic and climatic factors (15). Rainfall patterns, particularly the erratic Northeast Monsoon, are a primary driver, with regions dependent on it experiencing more frequent droughts (16). The Western Ghats' rain shadow effect exacerbates drought in leeward districts (17). Topography, soil type and land use also play crucial roles in influencing water availability and retention (18).

SPI-based annual drought frequency projection revealed that the drought frequency is expected to have slight changes toward the increasing side (up to 3 years) in the near century for the RCP 4.5 scenario (Fig. 2). In contrast, the decline in drought frequency is projected up to 6 counts in the mid-century. The increase in drought frequency of 6 to 7 counts is projected for most regions by the end of the century. It is seen from the results that a few changes would happen in the drought occurrence in all the time scales under the RCP 8.5 scenario. In the future, the drought occurrence is projected to decrease in most places in Tamil Nadu. This might be due to the projected increase in the monsoon rainfall due to enhanced hydrological activity as a result of the expected increase in temperature. These results are in line with the findings of (19, 20). Seasonal SPI indicates that the changes in drought frequency patterns during SWM and NEM are expected to be almost similar to the changes in annual drought frequency in the future climatic conditions of the RCP 4.5 scenario (Fig. 3 & 4). The projected change in SWM and NEM drought frequency follows the same pattern of annual drought frequency in all the time scales except near century during SWM and mid-century of NEM with the RCP 8.5 scenario.

The percentage of drought frequency for moderate (RCP 4.5) and high (RCP 8.5) emission scenarios for the Southwest Monsoon (SWM) and Northeast Monsoon (NEM) at the near, mid and end of the century is presented in Table 2. It was found that the drought frequency is projected to be higher in the SWM (20%) compared to the NEM in the future.




Fig. 2. Projected changes in annual drought frequency in future based on SPI.




Fig. 3. Projected changes in SWM drought frequency in future based on SPI.




Fig. 4. Projected changes in NEM drought frequency in future based on SPI.

Table 2. Percentage of the number of drought years in future

No of drought years (%)	RCP 4.5						RCP 8.5					
	Near		Mid		End		Near		Mid		End	
	SWM	NEM	SWM	NEM	SWM	NEM	SWM	NEM	SWM	NEM	SWM	NEM
No drought		45	35	0	0	0	0	0	0	0	6	26
1	0	32	32	0	0	0	0	6	29	0	13	26
2	0	13	23	19	0	0	0	32	35	0	23	45
3	0	10	6	10	6	0	13	32	26	6	29	3
4	29	0	0	35	6	3	3	16	6	16	19	0
5	23	0	0	23	10	10	26	13	3	23	6	0
6	29	0	3	6	16	6	23	0	0	23	3	0
7	13	0	0	3	29	16	13	0	0	16	0	0
8	6	0	0	3	13	13	10	0	0	6	0	0
9	0	0	0	0	13	19	6	0	0	3	0	0
10	0	0	0	0	3	19	6	0	0	3	0	0
11	0	0	0	0	3	10	0	0	0	3	0	0
12	0	0	0	0	0	3	0	0	0	0	0	0

SPI index-based drought occurrence reveals that there would be more drought in the RCP 4.5 scenario (50%) compared to the RCP 8.5 scenario. These results align with previous research showing increased drought frequency under the RCP 4.5 scenario, suggesting more water scarcity compared to RCP 8.5 (19, 21). Related studies observed higher drought frequency under the moderate emission scenario RCP 4.5 compared to the high emission scenario RCP 8.5 (22). This can be attributed to complex regional rainfall shifts, where RCP 4.5 may lead to more pronounced drying in crucial agricultural areas (23). Although evapotranspiration is higher under RCP 8.5, increased precipitation in some regions might offset this effect (24). Changes in atmospheric circulation patterns could favor drought-prone conditions in specific locations under RCP 4.5. Additionally, drier initial soil moisture conditions under RCP 4.5 could contribute to more frequent droughts (25). During the mid-century (2041-2070), change in severe drought frequency based on SPI is projected to decrease over the majority of the regions in Tamil Nadu under the

RCP 4.5 and RCP 8.5 due to high rainfall projected in mid-century (26).

Conclusion

Drought prediction in Tamil Nadu has become important as climate change brings unpredictable weather conditions. In future, the frequency of drought will be higher in the Southwest monsoon compared to the Northeast monsoon. Based on SPI, drought occurrence is expected to increase in the 4.5 scenario as compared to the 8.5 scenario and both under RCP 4.5 and RCP 8.5, drought occurrence is projected to decrease due to an increase in rainfall projected in mid-century. This suggests that the future climate conditions, as predicted by these scenarios, may reduce drought events in Tamil Nadu. A decrease in drought occurrence projected in the future may be due to the rainfall patterns predicted by climatic models.

This research provides essential insights that strengthen early warning systems, facilitate proactive mitigation strategies and sustainable water management, prompting adaptation strategies like emission reduction, increased water storage and drought-resistant practices. These insights equip policymakers to make informed decisions, implement effective drought mitigation measures and build a more resilient towards the drought risks.

Acknowledgements

The first author would also like to express sincere thanks to Jagannathan R., vice-chancellor, Periyar University, Salem, Tamil Nadu for his constant support during the research period.

Authors' contributions

VG and MV conceptualized the study. MV and KB carried out the methodology. SE and SA were responsible for the software. VG and MV carried out the investigation. VG and CD handled data curation. MV and TB prepared the original draft. MV, KB, AS and TB were involved in the writing-review and editing. All authors have read and approved the final manuscript.

Compliance with ethical standards

Conflict of interest: The Authors declare that there is no conflict of interest.

Ethical issues: None

References

1. Radadiya Kiem AS, Johnson F, Westra S, van Dijk A, Evans JP, Donnell A, et al. Natural hazards in Australia: droughts. *Clim Change*. 2016;139:37–54. <https://doi.org/10.1007/s10584-016-1798-7>
2. Smith AB, Katz RW. US billion-dollar weather and climate disasters: data sources, trends, accuracy and biases. *Nat Hazards*. 2013;67(2):387–410. <https://doi.org/10.1007/s11069-013-0566-5>
3. Reichstein M, Bahn M, Ciais P, Frank D, Mahecha MD, Seneviratne SI, et al. Climate extremes and the carbon cycle. *Nature*. 2013;500(7462):287–95. <https://doi.org/10.1038/nature12350>
4. Roxy MK, Ritika K, Terray P, Murtugudde R, Ashok K, Goswami BN. Drying of Indian subcontinent by rapid Indian Ocean warming and a weakening land-sea thermal gradient. *Nat Commun*. 2015;6(1):7423. <https://doi.org/10.1038/ncomms8423>
5. Shah RD, Mishra V. Development of an experimental near-real-time drought monitor for India. *J Hydrometeorol*. 2015;16(1):327–45. <https://doi.org/10.1175/JHM-D-14-0041.1>
6. Prasanna V. Impact of monsoon rainfall on the total foodgrain yield over India. *J Earth Syst Sci*. 2014;123(5):1129–45. <https://doi.org/10.1007/s12040-014-0444-x>
7. Gadgil S, Rupa KK. The Asian monsoon – agriculture and economy. In: *The Asian monsoon*. Springer Praxis Books. Springer, Berlin, Heidelberg; 2006. p. 651–83. Available from: https://doi.org/10.1007/3-540-37722-0_18
8. Wetterhall F, Winsemius HC, Dutra E, Werner M, Pappenberger E. Seasonal predictions of agro-meteorological drought indicators for the Limpopo basin. *Hydrol Earth Syst Sci*. 2015;19(6):2577–86. <https://doi.org/10.5194/hess-19-2577-2015>
9. Tsakiris G, Pangalou D, Vangelis H. Regional drought assessment based on the Reconnaissance Drought Index (RDI). *Water Resour Manag*. 2007;21:821–33. <https://doi.org/10.1007/s11269-006-9105-4>
10. Bhalme HN. Droughts: Historical perspectives agrometeorology (Agricultural droughts: Data base management). Lecture Notes of the second SERC School; 1995. p. 54
11. Naresh KM, Murthy CS, Sessa SMV, Roy PS. On the use of Standardized Precipitation Index (SPI) for drought intensity assessment. *Meteorol Appl*. 2009;16(3):381–89. <https://doi.org/10.48550/arXiv.1509.04808>
12. Nosrati K, Zareiee AR. Assessment of meteorological drought using SPI in West Azarbaijan Province, Iran. *J Appl Sci Environ Manag*. 2011;15(4):563–69. <https://www.ajol.info/index.php/jasem/article/view/88628>
13. Meinshausen M, Smith SJ, Calvin K, Daniel JS, Kainuma ML, Lamarque JF, et al. The RCP greenhouse gas concentrations and their extensions from 1765 to 2300. *Clim Change*. 2011;109:213–41. <https://doi.org/10.1007/s10584-011-0156-z>
14. McKee TB, Doesken NJ, Kleist J. The relationship of drought frequency and duration to time scales. In: *Proceedings of the 8th Conference on Applied Climatology*; 1993 Jan 17–22; Anaheim, California; 1993. 17(22):179–83. Available from: https://www.droughtmanagement.info/literature/AMS_Relationship_Drought_Frequency_Duration_Time_Scales_1993.pdf
15. Rajkumar R, Shaijumon CS, Gopakumar B, Gopalakrishnan D. Extreme rainfall and drought events in Tamil Nadu, India. *Clim Res*. 2020;80(3):175–88. <https://doi.org/10.3354/cr01600>
16. Bharambe KP, Shimizu Y, Kantoush SA, Sumi T, Saber M. Impacts of climate change on drought and its consequences on the agricultural crop under worst-case scenario over the Godavari River Basin, India. *Clim Serv*. 2023;32:100415. <https://doi.org/10.1016/j.cliser.2023.100415>
17. Masroor M, Rehman S, Avtar R, Sahana M, Ahmed R, Sajjad H. Exploring climate variability and its impact on drought occurrence: Evidence from Godavari Middle sub-basin, India. *Weather Clim Extrem*. 2020;30:100277. <https://doi.org/10.1016/j.wace.2020.100277>
18. Zhang Z, Zhang Y, Henderson M, Wang G, Chen M, Fu Y, et al. Effect of land use type on soil moisture dynamics in the sloping lands of the black soil (Mollisols) region of Northeast China. *Agri*. 2024;14(8):1261. <https://doi.org/10.3390/agriculture14081261>
19. Mukherjee S, Aadhar S, Stone D, Mishra V. Increase in extreme precipitation events under anthropogenic warming in India. *Weather Clim Extrem*. 2018;20:45–53. <https://doi.org/10.1016/j.wace.2018.03.005>
20. Jayasankar CB, Surendran S, Rajendran K. Robust signals of future projections of Indian summer monsoon rainfall by IPCC AR5 climate models: Role of seasonal cycle and interannual variability. *Geophys Res Lett*. 2015;42(9):3513–20. <https://doi.org/10.1002/2015GL063659>
21. Turner AG, Annamalai H. Climate change and the South Asian summer monsoon. *Nat Clim Chang*. 2012;2(8):587–95. <https://doi.org/10.1038/nclimate1495>
22. Alemu MG, Wubneh MA, Worku TA, Womber ZR, Chanie KM. Comparison of CMIP5 models for drought predictions and trend analysis over Mojo catchment, Awash Basin, Ethiopia. *Sci Afr*. 2023;22:e01891. <https://doi.org/10.1016/j.sciaf.2023.e01891>
23. Seneviratne SI, Zhang X, Adnan M, Badi W, Dereczynski C, Luca AD, et al. Weather and climate extreme events in a changing climate; 2021 Available from: <https://www.ipcc.ch/report/ar6/wg1/chapter/chapter-11/>

24. Gurara MA, Jilo NB, Tolche AD. Impact of climate change on potential evapotranspiration and crop water requirement in Upper Wabe Bridge watershed, Wabe Shebele River Basin, Ethiopia. *J Afr Earth Sci.* 2021;180:104223. <https://doi.org/10.1016/j.jafrearsci.2021.104223>
25. Naumann G, Alfieri L, Wyser K, Mentaschi L, Betts RA, Carrao H, et al. Global changes in drought conditions under different levels of warming. *Geophys Res Lett.* 2018;45(7):3285–96. <https://doi.org/10.1002/2017GL076521>
26. Aadhar S, Mishra V. Impact of climate change on drought frequency over India. *Climate Change and Water Resources in India* Publisher: Ministry of Environment, Forest and Climate Change (MoEF and CC), Government of India; 2018 Available from: https://www.researchgate.net/publication/330161820_Impact_of_Climate_Change_on_Drought_Frequency_over_India