



RESEARCH ARTICLE

# Effect of irrigation regimes and nano-urea based nitrogen management on yield attributes, productivity, nutrient acquisition and use efficiency of wheat in Eastern Plateau region of India

Abhijit Mandal<sup>1\*</sup>, Teekam Singh<sup>1</sup>, Smruti Ranjan Padhan<sup>2</sup>, Ayan Sarkar<sup>1</sup>, Anchal Dass<sup>1</sup> & Manoj Chaudhary<sup>3</sup>

<sup>1</sup>Division of Agronomy, ICAR-Indian Agricultural Research Institute, New Delhi 110 012, India

<sup>2</sup>Division of Agronomy, KVK-East Sikkim, ICAR-Research Complex for NEH Region, Umiyam 737 135, Sikkim, India

<sup>3</sup>Division of Soil Science & Agricultural Chemistry, ICAR-Indian Agricultural Research Institute, 825 405, Jharkhand, India

\*Email - [abhijari9875@gmail.com](mailto:abhijari9875@gmail.com)



## ARTICLE HISTORY

Received: 11 January 2025

Accepted: 17 March 2025

Available online

Version 1.0 : 10 May 2025



## Additional information

**Peer review:** Publisher thanks Sectional Editor and the other anonymous reviewers for their contribution to the peer review of this work.

**Reprints & permissions information** is available at [https://horizonepublishing.com/journals/index.php/PST/open\\_access\\_policy](https://horizonepublishing.com/journals/index.php/PST/open_access_policy)

**Publisher's Note:** Horizon e-Publishing Group remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

**Indexing:** Plant Science Today, published by Horizon e-Publishing Group, is covered by Scopus, Web of Science, BIOSIS Previews, Clarivate Analytics, NAAS, UGC Care, etc See [https://horizonepublishing.com/journals/index.php/PST/indexing\\_abstracting](https://horizonepublishing.com/journals/index.php/PST/indexing_abstracting)

**Copyright:** © The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited (<https://creativecommons.org/licenses/by/4.0/>)

## CITE THIS ARTICLE

Abhijit M, Teekam S, Smruti RP, Ayan S, Anchal D, Manoj C. Effect of irrigation regimes and nano-urea based nitrogen management on yield attributes, productivity, nutrient acquisition and use efficiency of wheat in Eastern Plateau region of India. Plant Science Today (Early Access). <https://doi.org/10.14719/pst.7167>

## Abstract

Wheat (*Triticum aestivum* L.) production is vital for India's food security, with projections indicating a substantial 70-104 % increase in demand by 2050. However, Eastern Plateau regions face challenges of warmer climate leading to aberration in irrigation regimes (IRs) and nutrient, especially nitrogen (N) management resulting in lower productivity. Study revealed that IRs significantly influenced yield attributes viz., spikelets/spike and 1000-grain weight, with moisture stress during grain filling stages adversely impacting grain development, resulted in 29 % yield reduction. The 100 % recommended dose of nitrogen (RDN) resulted in significantly superior yield attributes; surpassing nano-urea based nitrogen application. The grain yield with 100 % RDN leading with the highest yield and grain N content of 7.0 and 14.5 % more compared to nano-urea based 50 % RDN+2 NUS (nano-urea spray) treatment respectively. Interaction effect of 3-irrigation regimes with 100 % RDN on grain yield was significantly superior over NUS nullifying the synergistic effect of NUS with IRs. However, 50 % RDN+2 NUS reported significantly ( $p=0.05$ ) superior N content (0.42 %) and N uptake (31.30 kg/ha) in straw. Additionally, five irrigations exhibited significantly higher grain and total N uptake by 30.9 and 25.78 % compared to two irrigations, respectively. Apparent nitrogen recovery and agronomic nitrogen use efficiency were the highest in 50 % RDN+2 NUS due to better N acquisition and less amount of N application through NUS. Thus 3-irrigation regimes and 100 % RDN can be recommended as an agronomic management practice for maximising wheat productivity in Eastern Plateau region of India.

## Keywords

irrigation regimes, moisture stress, nano-urea, nitrogen use efficiency, nutrient content, productivity

## Introduction

Wheat, scientifically known as *Triticum aestivum* L. is a high-energy consuming winter cereal that provides 112.74 million tons of grain annually from an area of 30.5 million hectares, accounting for 35 % of India's food grain production (1). According to projections, the total demand for wheat is expected to climb by 32-38 % by 2030 (74 kg/capita) and by 70-104 % by 2050 (94 kg/capita), highlighting the necessity of giving priority to expand the production and acreage to guarantee food and nutritional security (2). Even though wheat is highly adaptable in India but its production in Jharkhand (a state in India's eastern plateau region) makes

up less than 1 % of the country's total output. The crop is grown in an area of 2.21 lakh hectares, with an average productivity of 2.13 t/ha, which is lower than the national average of 3.5 t/ha. This is primarily because of the state's comparatively warmer climate (5-7 °C higher temperature), which affects the crop growth (3). This low productivity zone receives hot winds at grain filling stage which reduces growth duration and size of the grain. The Eastern States including Jharkhand have potential for higher wheat yields (4.5 t/ha) which was demonstrated by front-line demonstrations. Future improvements in production are expected from these low-productivity zones. The reduction of wheat cultivation in high-productivity areas could be driven by issues such as falling water tables, infestations of *Phalaris minor*, or the shift toward growing high-value crops. These changes could shift the focus of production to areas previously considered less productive. Irrigation during booting to heading phases improves spike and grain development and helps to increase productivity (4). Drought stress conditions negatively impact wheat yield and its components, highlighting the importance of proper irrigation management practices based on critical growth stages to minimize the yield gap (5). Nitrogen (N) deficiency in soils, is a significant factor contributing to the low productivity of wheat in Eastern India. Nitrogen serves as a fundamental structural component in various essential biological compounds such as proteins, enzymes, chlorophyll, Rubisco, nucleic acids and certain hormones. Consequently, nitrogen fertilization becomes a vital agronomic management practice to boost crop productivity, especially during the vegetative growth stages. Proper management of nitrogenous fertilizer is crucial for maximizing crop productivity. However, nitrogen losses such as nitrate leaching, denitrification and runoff to surface and groundwater reduce fertilizer efficiency to 50-60 %, causing economic losses and environmental pollution (6). To address these challenges, nano-fertilizers show promise in enhancing nutrient uptake and use efficiency, reducing losses through leaching and emissions and minimizing the risk of nutrient toxicity. However, information on the interaction between irrigation and different nitrogen management practices in the plateau region of India is limited. Keeping this in the context, a field experiment was set up to figure out the effect of various irrigation regimes and nitrogen management practices on the yield attributes, productivity, nutrient acquisition and use efficiency of wheat under eastern plateau region of India.

## Materials and Methods

The field experiment was conducted at the farm of ICAR-Indian Agricultural Research Institute, Jharkhand during 2021-2022. The farm is situated at 24°16' N latitude, 85°21' E longitude and has an elevation of 628 meters above MSL. The long-term weather data of the site of experimentation denotes it as a semi-arid and sub-tropical climate with hot, dry summers in May and June and moderately cold winters from late November to January. The soil in the experimental field is sandy clay loam, offering good drainage and low water-holding capacity. It has an acidic pH of 5.85, low electrical conductivity (EC) of 0.712dS/m, low cation exchange capacity of 7.8 c mol (P<sup>+</sup>)/kg, low organic carbon content of 0.25 % and low levels of available nitrogen and phosphorus (150.6 kg N/ha and 8.27 kg P<sub>2</sub>O<sub>5</sub>/ha)

and medium availability of potassium (132.16 kg K<sub>2</sub>O/ha). During the growing period, crop received a total rainfall of 100.5 mm, although it was unevenly distributed throughout the season. The experiment was set up in a split-plot design with the main plots assigned to three irrigation regimes (IRs) viz., I<sub>1</sub> (5-irrigations on a priority basis), I<sub>2</sub> (3-irrigations at CRI, flowering and milking stage) and I<sub>3</sub> (2-irrigations at CRI and flowering stage) and five nano urea-based nitrogen management practices (NMPs) in the subplots viz., N<sub>0</sub> (control without nitrogen application), N<sub>1</sub> (100 % RDN-120 kg N/ha, split as 1/3rd basal, 1/3rd at CRI and 1/3rd at the 2nd irrigation), N<sub>2</sub> (50 % RDN, half basal and half at CRI, with nano-urea spray at 60 DAS), N<sub>3</sub> (50 % RDN, half basal and half at CRI, with two nano-urea spray at 45 DAS & 70 DAS) and N<sub>4</sub> (75 % RDN, half basal and half at CRI, with nano-urea spray at 60 DAS) and replicated thrice. The main plot size was 22m x 5m, while the subplot size was 5m x 4m. N in the form of prilled urea was applied according to the treatments and full doses of phosphorus (P) and potassium (K) were applied as a basal application at a rate of N: P<sub>2</sub>O<sub>5</sub>: K<sub>2</sub>O - 120:60:40 kg/ha. Nano-urea was applied @ 4 ml per litre of water in the evening time, as per the treatment specifications. The first irrigation was applied at the crown root initiation stage and subsequent irrigations were given based on recommended guidelines, taking local weather conditions into consideration. Irrigation was withheld 15 days before harvesting of the crop. The wheat variety DBW-187 (Karan Vandana) was sown with row spacing of 22.5 cm and a seed rate of 100 kg/ha. The pests and weed control measures were undertaken across the treatments throughout the crop growth period as and when required. At maturity, the number of effective tillers per square meter was determined by counting tillers with more than 50 % of the ear filled. The length of the main spikes of the plants, measured from the base to the tip of the floret (excluding awns), was recorded as ear length (in cm). The grain quantity within each ear head was assessed and their weight was recorded. The average value of the number of grains per spike was calculated. Before threshing, biological yield was measured and after threshing, the total grain weight and straw weight from the net plot area were recorded and converted to a metric ton per hectare basis at a constant moisture content of 12 %. The Harvest Index (HI) was calculated by using following expression (7).

$$HI = \frac{\text{Grain yield}}{\text{Biological yield}}$$

The modified alkaline potassium permanganate method was used to estimate the amount of available N (8), for available P Bray and Kurtz method was used (9). The available K (kg/ha) was extracted using neutral normal ammonium acetate solution and measured using a flame photometer. The N concentration in dried grain and straw samples was determined by Kjeldahl digestion method (8). The P content was quantified using the Vanado-molybdate phosphoric acid yellow colour method (11) with a Spectronic-20 colorimeter with blue filter and the K content of grain and straw was assessed using a Flame photometer (11). The grain protein content was determined by multiplying N content in percent with a factor 5.83 (12).

Partial Factor Productivity (PFP) is calculated by following formulas:

$$PFP = \frac{Y_f}{N_a}$$

Where,

$Y_f$  = Yield obtained from fertilized plot

$N_a$  = Nutrient applied (kg/ha)

Efficiency indices were computed using the following formulas-

Agronomic N use efficiency (ANUE) =

$$\frac{Y_t - Y_o}{A_t} \text{ Kg grain/kg N applied}$$

$$\text{Physiological N use efficiency (PE}_N\text{)} = \frac{Y_t - Y_o}{U_t - U_o}$$

$$\text{Apparent N recovery (ANR)} = \frac{U_t - U_o}{N_a} \times 100$$

N efficiency ratio (NER) =

$$\frac{\text{Dry matter yield (kg/ha)}}{\text{N accumulated at harvest (kg/ha)}}$$

Physiological efficiency index of N (PEN) =

$$\frac{\text{Grain yield (kg/ha)}}{\text{N absorbed by biomass (kg)}}$$

Where,

$Y_t$  = Yield in the test treatment (kg/ha)

$Y_o$  = Yield in the control (kg/ha)

$A_t$  = Units of N applied in the test treatment (kg/ha)

$U_t$  = Uptake of N in the test treatment (kg/ha)

$U_o$  = Uptake of N in the control plot (kg/ha)

$N_a$  = N applied to the test treatment (kg/ha)

Data analysis for each character was conducted using Analysis of Variance (ANOVA) and the significance of a split plot design was assessed using the 'F' test (13). The analysis for the split plot design was carried out using an online data analysis module, specifically the strengthening statistical

computing for NARS portal (<https://sscnars.icar.gov.in/>). The treatment comparisons were made at a significance level of 5 % and Critical Difference (CD) and standard error of means (SEM $\pm$ ) were calculated for each character. Graphical representations of the data were included as needed.

## Results

### Yield attributes

Irrigation regimes (IRs) did not have any significant difference in ear-bearing tillers/m<sup>2</sup>, whereas among NMPs the highest no. of ear-bearing tillers (EBT)/m<sup>2</sup> was recorded with both  $N_1$  and  $N_3$ , which were almost 102 % higher than control (Table 1 and Fig. 1). The highest percentage of EBT/m<sup>2</sup> was observed with  $N_3$  (71.36 %) which was significantly superior to the  $N_1$  (66.50 %) and  $N_4$  (67.04 %). The NMPs settled a significant difference in ear length possessing the maximum ear length (10.64 cm) with  $N_1$ , similar with  $N_3$ . The  $N_2$ ,  $N_4$  and  $N_1$  produced 26.1, 32.5 and 42.8 % more ear length over  $N_0$ . Among NMPs,  $N_3$  resulted in the higher no. of spikelets per spike (21.44), statistically comparable to  $N_1$ , greater than other treatments (Fig. 2). Both  $I_1$  and  $I_2$  exhibited the highest no. of grains per ear head, surpassing  $I_3$  treatments. The  $N_1$  yielded the greatest grains per ear head, slightly outperforming  $N_3$  and  $N_4$ . Remarkably  $N_1$ ,  $N_2$ ,  $N_3$  and  $N_4$  exhibited 95.15 %, 54.35 %, 86.41 % and 67.93 % higher grain per ear head, respectively, in comparison to the  $N_0$ .  $I_1$  exhibited highest 1000-grain weight, significantly surpassing  $I_2$  and  $I_3$ . On the other hand,  $N_1$  achieved the maximum 1000-grain weight of 40.53 g, notably higher than other NMPs. Differences in 1000-grain weight, ranging from 8.4 % to 17.27 %, were observed due to various NMPs compared to the control.

### Yield

Irrigation regimes has a significant effect (P value = 0.0038), whereas NMPs has a highly significant effect (P-value = 0), suggesting differences between at least two treatments and the interaction between them is also significant (P value = 0.003), suggesting that the effect of NMPs depends on the IRs treatments. Adequate irrigation ( $I_1$ ) produced the higher yield of 3.71 t/ha, comparable to  $I_2$ , which significantly surpass  $I_3$  which resulted 29 % yield reduction compared to  $I_2$ . NMPs significantly impacted grain yield,  $N_3$  followed as the second-

**Table 1.** Yield attributes and yield of wheat influenced by irrigation regime and nitrogen management practices

Treatment	EBT/m <sup>2</sup>	Spike length (cm)	No. of spikelets/spike	No. of grains/spike	1000-grain weight (g)	Grain yield (t/ha)	Straw yield (t/ha)	Biological yield (t/ha)	HI
<b>Irrigation regime</b>									
$I_1$	227.13	9.39	19.93 <sup>a</sup>	38.53 <sup>a</sup>	40.09 <sup>a</sup>	3.71 <sup>a</sup>	5.67	9.37 <sup>a</sup>	0.41
$I_2$	230.60	9.12	19.20 <sup>ab</sup>	38.13 <sup>a</sup>	39.27 <sup>b</sup>	3.64 <sup>a</sup>	5.63	9.26 <sup>a</sup>	0.40
$I_3$	217.27	9.30	18.13 <sup>b</sup>	33.73 <sup>b</sup>	35.41 <sup>c</sup>	2.82 <sup>b</sup>	5.05	7.87 <sup>b</sup>	0.37
SEm $\pm$	6.97	0.18	0.28	0.46	0.06	0.09	0.16	0.19	0.01
LSD (P $\leq$ 0.05)	NS	NS	1.09	1.80	0.24	0.35	NS	0.76	NS
<b>Nitrogen management</b>									
$N_0$	129.78 <sup>d</sup>	6.36 <sup>d</sup>	15.44 <sup>d</sup>	22.89 <sup>e</sup>	34.56 <sup>d</sup>	1.23 <sup>e</sup>	1.52 <sup>d</sup>	2.75 <sup>d</sup>	0.45 <sup>a</sup>
$N_1$	263.11 <sup>a</sup>	10.64 <sup>a</sup>	21.11 <sup>a</sup>	44.67 <sup>a</sup>	40.53 <sup>a</sup>	4.62 <sup>a</sup>	7.41 <sup>a</sup>	12.03 <sup>a</sup>	0.38 <sup>bc</sup>
$N_2$	219.56 <sup>c</sup>	8.97 <sup>c</sup>	18.11 <sup>c</sup>	35.33 <sup>d</sup>	37.46 <sup>c</sup>	2.94 <sup>d</sup>	4.67 <sup>c</sup>	7.61 <sup>c</sup>	0.39 <sup>b</sup>
$N_3$	262.78 <sup>a</sup>	10.50 <sup>a</sup>	21.44 <sup>a</sup>	42.67 <sup>b</sup>	39.48 <sup>b</sup>	4.32 <sup>b</sup>	7.42 <sup>a</sup>	11.74 <sup>a</sup>	0.37 <sup>c</sup>
$N_4$	249.78 <sup>b</sup>	9.61 <sup>b</sup>	19.33 <sup>b</sup>	38.44 <sup>c</sup>	39.24 <sup>b</sup>	3.82 <sup>c</sup>	6.22 <sup>b</sup>	10.05 <sup>b</sup>	0.38 <sup>bc</sup>
SEm $\pm$	3.02	0.06	0.28	0.62	0.23	0.06	0.16	0.20	0.01
LSD (P $\leq$ 0.05)	8.82	0.21	0.83	1.81	0.58	0.17	0.47	0.57	0.02
IRs $\times$ NMPs	NS	NS	NS	NS	NS	0.29	NS	NS	NS
NMPs $\times$ IRs	NS	NS	NS	NS	NS	0.43	NS	NS	NS

( $I_1$ : 5 irrigations;  $I_2$ : 3 irrigations;  $I_3$ : 2 irrigations;  $N_0$ : control;  $N_1$ : 100 % RDN;  $N_2$ : 50 % RDN+1 NUS;  $N_3$ : 50 % RDN+2 NUS;  $N_4$ : 75 % RDN+1 NUS). Treatments with same letter are not significantly different (p=0.05)





**Fig. 1.** Picture of the plant samples from experiment at flowering stage.

(From Left to Right:- N<sub>0</sub>, N<sub>1</sub>, N<sub>2</sub>, N<sub>3</sub> and N<sub>4</sub>)



**Fig. 2.** Effect of irrigation regimes and nitrogen management on spikes of wheat.

best treatment with 4.32 t/ha, whereas N<sub>1</sub> leading with the highest yield, 7 % more yield compared to N<sub>3</sub>, significantly outperforming other NMPs. Notably, yield under N<sub>1</sub> was 21 % more than N<sub>4</sub>. N<sub>3</sub> recorded 45 % and 13 % more yield compared to N<sub>2</sub> and N<sub>4</sub> respectively. All NMPs - N<sub>1</sub>, N<sub>2</sub>, N<sub>3</sub> and N<sub>4</sub> yielded significantly higher results, increasing by 275.6 %, 139 %, 251 % and 210.5 % respectively, compared to the control (Table 1). Significant interactions emerged between IRs and NMPs where the higher yield was obtained with I<sub>2</sub>N<sub>1</sub> and was statistically like I<sub>1</sub>N<sub>1</sub>, while the lowest yield was in I<sub>3</sub>N<sub>0</sub>. The study revealed no statistically significant differences between IRs in terms of straw yield. However, NMPs strongly influenced wheat straw output. N<sub>1</sub> (7.41 t/ha) and N<sub>3</sub> (7.42 t/ha) significantly increased straw yield compared to others. In the realm of IRs, elevated biological yield was evident with I<sub>1</sub> and I<sub>2</sub> showcased 19.05 and 17.66 % increase in productivity over I<sub>3</sub>. The higher biological yield was observed in both N<sub>1</sub> and N<sub>3</sub> with yields of 337.4 % and 326.9 % higher than the control (N<sub>0</sub>) respectively. In comparison, N<sub>2</sub> and N<sub>4</sub> yielded 176.7 % and 265.4 % higher than the control. Notably, NMPs had a substantial impact on the HI, with the control group (N<sub>0</sub>) demonstrating a significantly higher index than other practices. Among managed nitrogen levels, N<sub>3</sub> had a lower harvest index, statistically like N<sub>1</sub> and N<sub>4</sub>, but significantly different from N<sub>2</sub>, which was also found to be similar to N<sub>1</sub> and N<sub>4</sub>. The highest grain yield was obtained with I<sub>2</sub>N<sub>1</sub> which was statistically like I<sub>1</sub>N<sub>1</sub> whereas the lowest was observed in I<sub>3</sub>N<sub>0</sub> (Fig. 3).

### Protein content

Protein content demonstrated differences, reduced moisture correlated with increased protein with the lowest under I<sub>1</sub> and the highest (10.21 %) under I<sub>3</sub> irrigation regimes. N<sub>1</sub> had the highest protein (10.99 %), contrasting with the lowest in the control (Table 2). Application of higher doses of prilled urea in soil resulted more protein content compared to nano-urea application. Protein content of grains under N<sub>4</sub> was 8.1 % and 4 % more compared to N<sub>3</sub> and N<sub>2</sub> respectively. There is no significant interaction between IRs and NMPs.

### N, P, K content and uptake by crop


Irrigation regimes did not significantly affect the nitrogen content in grain; however, higher irrigation levels resulted in a reduced nitrogen content, ranging from 1.67 % to 1.75 %, as shown in Table 2. Nitrogen content in straw decreased with fewer irrigations, with I<sub>1</sub> having the highest (0.37 %) and I<sub>3</sub> having the lowest (0.28 %) nitrogen content. N<sub>1</sub> recorded the highest grain nitrogen content (1.89 %) among practices, while control had the lowest. 1.48 %). N<sub>2</sub>, N<sub>3</sub> and N<sub>4</sub> showed 1.72 %, 1.65 % and 1.79 % nitrogen content, respectively. N<sub>1</sub>, N<sub>2</sub>, N<sub>3</sub> and N<sub>4</sub> contained 27.70 %, 16.22 %, 11.48 % and 20.95 % respectively more nitrogen in grain compared to the control. N<sub>3</sub> had the highest straw nitrogen content (0.42 %), followed by N<sub>1</sub> (0.35 %), N<sub>4</sub> (0.34 %) and N<sub>2</sub> (0.32 %), with N<sub>4</sub> was statistically like both N<sub>1</sub> and N<sub>2</sub>. Higher nitrogen content (0.45 %) in straw was observed in I<sub>1</sub>N<sub>3</sub> treatment which was statistically like I<sub>1</sub>N<sub>1</sub>, I<sub>2</sub>N<sub>3</sub> and I<sub>1</sub>N<sub>4</sub> treatments. I<sub>1</sub> exhibited 30.9 % higher total N uptake compared to I<sub>3</sub> as grain N uptake of I<sub>1</sub> was 25.78 % greater than I<sub>3</sub>. I<sub>1</sub> resulted the highest total N uptake (86.25 kg/ha) and resulted in 17.2 % and 47.6 % higher straw N uptake compared to I<sub>2</sub> and I<sub>3</sub>. On the other hand, N<sub>1</sub> demonstrated the highest grain N uptake (86.88 kg/ha) and lowest under the control (18.09 kg/ha). N<sub>3</sub> (71.46 kg/ha) was like N<sub>4</sub> (67.92 kg/ha), both surpassing N<sub>2</sub> (50.80 kg/ha) in grain N uptake. Notably, in the interaction, the highest straw N uptake was observed in I<sub>1</sub>N<sub>3</sub> (34.06 kg/ha), like I<sub>1</sub>N<sub>1</sub> and I<sub>2</sub>N<sub>3</sub> (Fig. 4). For the total N uptake, the highest total nitrogen uptake was observed in I<sub>1</sub>N<sub>1</sub> (27.46 kg/ha) that was statistically similar with I<sub>2</sub>N<sub>1</sub> (116.80 kg/ha) (Fig. 5).

Phosphorus (P) content in grains (0.3 %) and straw (0.07 %) showed no statistical differences across irrigation regimes (IRs). P content varied from 0.27 % to 0.32 % due to NMPs, with N<sub>1</sub> (0.32 %) and N<sub>3</sub> (0.31 %) exhibiting the higher values. For potassium (K), I<sub>1</sub> recorded the higher grain K content (0.36 %), followed by I<sub>2</sub> and I<sub>3</sub>, with similar trends in straw. The highest K content (0.38 %) was observed in N<sub>3</sub>, comparable to N<sub>2</sub> and N<sub>4</sub>. Increased nano-urea doses enhanced grain K concentration but reduced straw K content. Grain P uptake was highest in I<sub>1</sub> (11.32 kg/ha), comparable to I<sub>2</sub> and exceeded I<sub>3</sub> by 32.55 %. Among NMPs, N<sub>3</sub> (13.55 kg/ha)

**Table 2.** Nitrogen concentration, uptake and grain protein of wheat as influenced by irrigation regimes and nitrogen management practices


Treatment	N concentration in grain (%)	N concentration in straw (%)	N uptake in grain (kg/ha)	N uptake in straw (kg/ha)	Total N uptake (kg/ha)	Crude protein concentration of grain (%)
<b>Irrigation regimes</b>						
I <sub>1</sub>	1.67	0.37 <sup>a</sup>	63.43 <sup>a</sup>	22.82 <sup>a</sup>	86.25 <sup>a</sup>	9.72
I <sub>2</sub>	1.70	0.31 <sup>b</sup>	63.23 <sup>a</sup>	19.47 <sup>b</sup>	82.70 <sup>a</sup>	9.90
I <sub>3</sub>	1.75	0.28 <sup>c</sup>	50.43 <sup>b</sup>	15.46 <sup>c</sup>	65.89 <sup>b</sup>	10.21
SEm±	0.06	0.01	2.09	0.55	2.54	0.37
LSD (P≤0.05)	NS	0.02	8.23	2.15	9.98	NS
<b>Nitrogen management practices</b>						
N <sub>0</sub>	1.48 <sup>d</sup>	0.17 <sup>d</sup>	18.09 <sup>d</sup>	2.58 <sup>e</sup>	20.67 <sup>e</sup>	8.64 <sup>d</sup>
N <sub>1</sub>	1.89 <sup>a</sup>	0.35 <sup>b</sup>	86.88 <sup>a</sup>	26.15 <sup>b</sup>	113.03 <sup>a</sup>	10.99 <sup>a</sup>
N <sub>2</sub>	1.72 <sup>bc</sup>	0.32 <sup>c</sup>	50.80 <sup>c</sup>	14.97 <sup>d</sup>	65.77 <sup>d</sup>	10.03 <sup>bc</sup>
N <sub>3</sub>	1.65 <sup>c</sup>	0.42 <sup>a</sup>	71.46 <sup>b</sup>	31.30 <sup>a</sup>	102.75 <sup>b</sup>	9.64 <sup>c</sup>
N <sub>4</sub>	1.79 <sup>b</sup>	0.34 <sup>bc</sup>	67.92 <sup>b</sup>	21.27 <sup>c</sup>	89.19 <sup>c</sup>	10.42 <sup>b</sup>
SEm±	0.03	0.01	1.71	0.82	2.10	0.18
LSD (P≤0.05)	0.09	0.03	4.99	2.41	6.13	0.52
IRs×NMPs	NS	0.05	NS	4.18	10.61	NS
NMP×IRs	NS	0.05	NS	4.28	12.61	NS

(I<sub>1</sub>: 5 irrigations; I<sub>2</sub>: 3 irrigations; I<sub>3</sub>: 2 irrigations; N<sub>0</sub>: control, N<sub>1</sub>: 100 % RDN, N<sub>2</sub>: 50 % RDN+1 NUS, N<sub>3</sub>: 50 % RDN+2 NUS, N<sub>4</sub>: 75 % RDN+1 NUS). Treatments with same letter are not significantly different (p=0.05)



**Fig. 3.** Interaction effect of irrigation regimes and nitrogen management practices on the grain yield of wheat [Similar letters above boxplots signifies non-significance among treatments ( $p=0.05$ ), (I<sub>1</sub>: 5 irrigations; I<sub>2</sub>: 3 irrigations; I<sub>3</sub>: 2 irrigations, N<sub>0</sub>: control, N<sub>1</sub>: 100 % RDN, N<sub>2</sub>: 50 % RDN+1 NUS, N<sub>3</sub>: 50 % RDN+2 NUS, N<sub>4</sub>: 75 % RDN+1 NUS)].


Upper bar is the maximum value, lower bar is minimum. Upper boundary of box is 3<sup>rd</sup> quartile, middle line is median and lower boundary of box is 1<sup>st</sup> quartile. The box size demonstrates the interquartile range. The dot in the middle of the box indicates mean value.



**Fig. 4.** Interaction effect of irrigation regimes and nitrogen management practices on the straw nitrogen uptake of wheat [Similar letters above boxplots signifies non-significance among treatments ( $p=0.05$ ), (I<sub>1</sub>: 5 irrigations; I<sub>2</sub>: 3 irrigations; I<sub>3</sub>: 2 irrigations, N<sub>0</sub>: control, N<sub>1</sub>: 100 % RDN, N<sub>2</sub>: 50 % RDN+1 NUS, N<sub>3</sub>: 50 % RDN+2 NUS, N<sub>4</sub>: 75 % RDN+1 NUS)].

Upper bar is the maximum value, lower bar is minimum. Upper boundary of box is 3<sup>rd</sup> quartile, middle line is median and lower boundary of box is 1<sup>st</sup> quartile. The box size demonstrates the interquartile range. The dot in the middle of the box indicates mean value.





**Fig. 5.** Interaction effect of irrigation regimes and nitrogen management practices on the total nitrogen uptake of wheat [Similar letters above boxplots signifies non-significance among treatments ( $p=0.05$ ), (I<sub>1</sub>: 5 irrigations; I<sub>2</sub>: 3 irrigations; I<sub>3</sub>: 2 irrigations, N<sub>0</sub>: control, N<sub>1</sub>: 100 % RDN, N<sub>2</sub>: 50 % RDN+1 NUS, N<sub>3</sub>: 50 % RDN+2 NUS, N<sub>4</sub>: 75 % RDN+1 NUS).

Upper bar is the maximum value, lower bar is minimum. Upper boundary of box is 3<sup>rd</sup> quartile, middle line is median and lower boundary of box is 1<sup>st</sup> quartile. The box size demonstrates the interquartile range. The dot in the middle of the box indicates mean value.

**Table 3.** Phosphorous and potassium concentration, uptake and grain protein of wheat as influenced by irrigation regimes and nitrogen management practices

Treatment	concentration in grain (%)		concentration in straw (%)		uptake in grain (kg/ha)		uptake in straw (kg/ha)		Total uptake (kg/ha)	
	P	K	P	K	P	K	P	K	P	K
<b>Irrigation regimes</b>										
I <sub>1</sub>	0.30	0.36	0.07	1.04	11.32 <sup>a</sup>	13.61 <sup>a</sup>	3.89	59.18 <sup>a</sup>	15.20 <sup>a</sup>	72.79 <sup>a</sup>
I <sub>2</sub>	0.30	0.35	0.07	1.00	10.99 <sup>a</sup>	13.00 <sup>a</sup>	3.91	56.03 <sup>a</sup>	14.90 <sup>a</sup>	69.03 <sup>a</sup>
I <sub>3</sub>	0.30	0.33	0.07	0.95	8.54 <sup>b</sup>	9.63 <sup>b</sup>	3.52	48.79 <sup>b</sup>	12.06 <sup>b</sup>	58.42 <sup>b</sup>
SEm±	0.01	0.01	0.01	0.04	0.19	0.16	0.16	1.83	0.26	1.93
LSD (P≤0.05)	NS	NS	NS	NS	0.74	0.64	NS	7.19	1.02	7.57
<b>Nitrogen management practices</b>										
N <sub>0</sub>	0.27 <sup>c</sup>	0.30 <sup>c</sup>	0.07	1.00 <sup>b</sup>	3.36 <sup>e</sup>	3.70 <sup>d</sup>	1.10 <sup>d</sup>	15.34 <sup>e</sup>	4.46 <sup>e</sup>	19.03 <sup>e</sup>
N <sub>1</sub>	0.32 <sup>a</sup>	0.32 <sup>b</sup>	0.07	1.12 <sup>a</sup>	14.76 <sup>a</sup>	15.03 <sup>b</sup>	5.14 <sup>a</sup>	83.41 <sup>a</sup>	19.9 <sup>a</sup>	98.43 <sup>a</sup>
N <sub>2</sub>	0.29 <sup>b</sup>	0.36 <sup>a</sup>	0.07	0.96 <sup>bc</sup>	8.62 <sup>d</sup>	10.71 <sup>c</sup>	3.33 <sup>c</sup>	45.14 <sup>d</sup>	11.94 <sup>d</sup>	55.85 <sup>d</sup>
N <sub>3</sub>	0.31 <sup>a</sup>	0.38 <sup>a</sup>	0.07	0.93 <sup>c</sup>	13.55 <sup>b</sup>	16.45 <sup>a</sup>	5.14 <sup>a</sup>	68.19 <sup>b</sup>	18.69 <sup>b</sup>	85.36 <sup>b</sup>
N <sub>4</sub>	0.29 <sup>b</sup>	0.38 <sup>a</sup>	0.07	0.97 <sup>bc</sup>	11.12 <sup>c</sup>	14.53 <sup>b</sup>	4.16 <sup>b</sup>	60.54 <sup>c</sup>	15.28 <sup>c</sup>	75.06 <sup>c</sup>
SEm±	0.01	0.01	0.00	0.01	0.37	0.33	0.21	1.73	0.28	1.93
LSD (P≤0.05)	0.02	0.02	NS	0.04	0.62	0.97	0.04	5.04	0.81	5.62
IRs×NMPs	NS	NS	NS	NS	1.07	1.68	NS	NS	NS	NS
NMPs×IRs	NS	NS	NS	NS	1.20	1.63	NS	NS	NS	NS


(I<sub>1</sub>: 5 irrigations; I<sub>2</sub>: 3 irrigations; I<sub>3</sub>: 2 irrigations, N<sub>0</sub>: control, N<sub>1</sub>: 100 % RDN, N<sub>2</sub>: 50 % RDN+1 NUS, N<sub>3</sub>: 50 % RDN+2 NUS, N<sub>4</sub>: 75 % RDN+1 NUS). Treatments with same letter are not significantly different ( $p=0.05$ )

surpassed N<sub>4</sub> (11.12 kg/ha) and N<sub>2</sub> (8.62 kg/ha). N<sub>1</sub> recorded the highest straw P uptake (5.14 kg/ha), like N<sub>3</sub>. Total P uptake was 346.2 % and 319.1 % higher in N<sub>1</sub> and N<sub>4</sub> compared to control, with N<sub>3</sub> exceeding N<sub>2</sub> and N<sub>4</sub> by 56.5 % and 22.1 %, respectively. The highest grain P uptake occurred in I<sub>1</sub>N<sub>1</sub> (16.12 kg/ha), while the lowest was in I<sub>3</sub>N<sub>0</sub>. Potassium uptake followed similar trends. I<sub>1</sub> achieved the highest grain (13.61 kg/ha) and straw (59.18 kg/ha) K uptake, leading to the highest total K uptake (72.79 kg/ha), comparable to I<sub>2</sub> but significantly exceeding I<sub>3</sub>. N<sub>3</sub> yielded the highest grain K uptake (16.45 kg/ha), surpassing N<sub>1</sub> (15.03 kg/ha) and N<sub>4</sub> (14.93 kg/ha). N<sub>1</sub> recorded the highest straw K uptake (83.41 kg/ha), resulting in 15.31 % more total K uptake compared to N<sub>3</sub>, while N<sub>3</sub> exceeded N<sub>2</sub> and N<sub>4</sub> by 52.8 % and 13.74 %, respectively. The highest grain K uptake was observed in I<sub>1</sub> N<sub>4</sub> (18.84 kg/ha), comparable to treatments I<sub>1</sub>N<sub>1</sub>, I<sub>1</sub>N<sub>2</sub>, I<sub>2</sub>N<sub>1</sub> and I<sub>2</sub>N<sub>2</sub> (Fig. 6) with the lowest in I<sub>3</sub>N<sub>0</sub> (2.92 kg/ha).

### Nitrogen use efficiency


The higher values of Agronomic Nitrogen Use Efficiency (ANUE) was achieved under irrigation regime I<sub>2</sub> (30.15 kg grain/kg N), like I<sub>1</sub> but significantly better than I<sub>3</sub> (Table 4). Among nitrogen levels, N<sub>3</sub> (60 kg N+2 nano-urea spray) showed the highest ANUE (51.48 kg grain/kg N), outperforming other practices. Interaction analysis indicated significant effects, with the highest ANUE in I<sub>2</sub>N<sub>3</sub>, like I<sub>1</sub>N<sub>3</sub> (Fig. 7).

The higher rate of Apparent Nitrogen Recovery (ANR) was achieved under I<sub>1</sub> (79.28 kg N uptake/kg N applied), like I<sub>2</sub> (79.02 kg N uptake/kg N applied), while I<sub>3</sub> had the lowest ANR (60.49 kg N uptake/kg N applied). N<sub>3</sub> exhibited the highest ANR (136.52 kg N uptake/kg N applied), notably surpassing other treatments.



**Fig. 6.** Interaction effect of irrigation regimes and nitrogen management practices on the grain potassium uptake of wheat [Similar letters above boxplots signifies non-significance among treatments ( $p=0.05$ ), (I<sub>1</sub>: 5 irrigations; I<sub>2</sub>: 3 irrigations, I<sub>3</sub>: 2 irrigations, N<sub>0</sub>: control, N<sub>1</sub>: 100 % RDN, N<sub>2</sub>: 50 % RDN+1 NUS, N<sub>3</sub>: 50 % RDN+2 NUS, N<sub>4</sub>: 75 % RDN+1 NUS).

Upper bar is the maximum value, lower bar is minimum. Upper boundary of box is 3<sup>rd</sup> quartile, middle line is median and lower boundary of box is 1<sup>st</sup> quartile. The box size demonstrates the interquartile range. The dot in the middle of the box indicates mean value.



**Fig. 7.** Interaction effect of irrigation regimes and nitrogen management practices on the agronomic nitrogen use efficiency of wheat [Similar letters above boxplots signifies non-significance among treatments ( $p=0.05$ ), (I<sub>1</sub>: 5 irrigations; I<sub>2</sub>: 3 irrigations, I<sub>3</sub>: 2 irrigations, N<sub>0</sub>: control, N<sub>1</sub>: 100 % RDN, N<sub>2</sub>: 50 % RDN+1 NUS, N<sub>3</sub>: 50 % RDN+2 NUS, N<sub>4</sub>: 75 % RDN+1 NUS).

Upper bar is the maximum value, lower bar is minimum. Upper boundary of box is 3<sup>rd</sup> quartile, middle line is median and lower boundary of box is 1<sup>st</sup> quartile. The box size demonstrates the interquartile range. The dot in the middle of the box indicates mean value.



**Table 4.** Effect of irrigation regimes and nitrogen management practices on various nitrogen use efficiencies

Treatment	ANUE (kg grain/ kg N)	ANR (kg N uptake/kg N applied)	PE <sub>N</sub> (kg grain/kg N uptake)	NER	PEN	PFP <sub>N</sub>
<b>Irrigation regimes</b>						
I <sub>1</sub>	29.28 <sup>a</sup>	79.28 <sup>a</sup>	29.41	112.81	46.17	44.02 <sup>a</sup>
I <sub>2</sub>	30.15 <sup>a</sup>	79.02 <sup>a</sup>	31.11	116.41	46.84	43.43 <sup>a</sup>
I <sub>3</sub>	22.84 <sup>b</sup>	60.49 <sup>b</sup>	30.62	122.08	45.45	33.66 <sup>b</sup>
SEm±	0.96	3.31	1.57	1.88	1.63	1.11
LSD (P≤0.05)	3.78	12.99	NS	NS	NS	4.34
<b>Nitrogen management practices</b>						
N <sub>0</sub>	-	-	-	133.04 <sup>a</sup>	59.3 <sup>a</sup>	-
N <sub>1</sub>	28.28 <sup>b</sup>	76.97 <sup>b</sup>	36.85	107.17 <sup>c</sup>	40.95 <sup>c</sup>	38.52 <sup>d</sup>
N <sub>2</sub>	28.53 <sup>b</sup>	75.09 <sup>b</sup>	38.98	116.48 <sup>b</sup>	45.28 <sup>b</sup>	48.97 <sup>b</sup>
N <sub>3</sub>	51.48 <sup>a</sup>	136.52 <sup>a</sup>	37.94	114.82 <sup>b</sup>	42.21 <sup>c</sup>	71.90 <sup>a</sup>
N <sub>4</sub>	28.83 <sup>b</sup>	76.08 <sup>b</sup>	38.13	113.99 <sup>b</sup>	43.02 <sup>bc</sup>	42.46 <sup>c</sup>
SEm±	0.75	3.04	1.06	1.83	1.78	0.80
LSD (P≤0.05)	2.18	8.87	NS	5.34	2.52	2.32
IRs×NMPs	3.78	NS	NS	NS	NS	NS
NMPs×IRs	5.01	NS	NS	NS	NS	NS

(NER: Nitrogen efficiency ratio, ANUE: Agronomic nitrogen use efficiency, PE<sub>N</sub>: Physiological nitrogen use efficiency, ANR: Apparent nitrogen recovery, PEN: Physiological efficiency index of nitrogen, PFP<sub>N</sub>: Partial factor productivity of N). Treatments with same letter are not significantly different (p=0.05)

(I<sub>1</sub>: 5 irrigations; I<sub>2</sub>: 3 irrigations, I<sub>3</sub>: 2 irrigations, N<sub>0</sub>: control, N<sub>1</sub>: 100 % RDN, N<sub>2</sub>: 50 % RDN+1 NUS, N<sub>3</sub>: 50 % RDN+2 NUS, N<sub>4</sub>: 75 % RDN+1 NUS)

N<sub>2</sub> (60 kg N + 50g N from nano-urea) displayed the highest physiological N use efficiency (38.98 kg grain/kg N uptake) followed by N<sub>4</sub>, N<sub>3</sub> and N<sub>1</sub> (38.13, 37.94 and 36.85 kg grain/kg N uptake respectively).

The control (N<sub>0</sub>) demonstrated the highest nitrogen efficiency ratio (133.04), while N<sub>1</sub> had the lowest ratio (107.17) attributed to higher N accumulation by the crop at harvest compared to other treatments.


The control (N<sub>0</sub>) exhibited the highest physiological efficiency index of nitrogen (59.3), while N<sub>1</sub> had the lowest value (40.95). Notably, N<sub>2</sub> demonstrated significantly higher physiological efficiency index (45.02).

The maximum value of partial factor productivity of nitrogen (PFP<sub>N</sub>) was achieved under I<sub>1</sub> (44.02 kg grain/kg N applied), like I<sub>2</sub> (43.43 kg grain/kg N applied), while I<sub>3</sub> had the

lowest (33.36 kg grain/kg N applied). N<sub>3</sub> demonstrated the highest value of PFP<sub>N</sub> (71.9 kg grain/kg N applied), outperforming other levels like N<sub>1</sub>, N<sub>2</sub> and N<sub>4</sub> had values of 38.52, 48.97 and 42.46 kg grain/kg N applied respectively.

### Stepwise regression analysis

Correlation panel graph between different parameters of wheat i.e., number (no.) of spikelet per spike, no. of grains per spike, grain yield, N use efficiency and others were analysed (Fig. 8). Grain yield was highly correlated with number (no.) of spikelet per spike ( $R^2=0.916$ ), no. of grains/spike ( $R^2=0.968$ ). Grain yield was also highly correlated with grain N uptake ( $R^2=0.976$ ); total N uptake ( $R^2=0.985$ ). Also, total N uptake was highly correlated with the grain N uptake ( $R^2=0.990$ ). Partial factor productivity of N was highly correlated with straw N concentration ( $R^2=0.875$ ); straw yield ( $R^2=0.811$ ) grain yield ( $R^2=0.794$ ). Agronomic nitrogen use efficiency was also



**Fig. 8.** Correlation panel graph between different parameters of wheat (N=45) [NSS: number (no.) of spikelet/ spike, NGS: no. of grains/spike, GY: Grain Yield, SY: Straw Yield, GNC: Grain Nitrogen Concentration, SNC: Straw N Concentration, GNU: Grain N Uptake, TNU: Total N Uptake, PFPN: Partial Factor Productivity of N, ANUE: Agronomic N Use Efficiency, Significance level: p=0.05\*, p=0.01\*\*, p=0.001\*\*\*].

correlated with grain yield ( $R^2=0.836$ ); straw yield ( $R^2=0.854$ ); and total N uptake ( $R^2=0.819$ ).

## Discussion

### Yield attributes and yield

Irrigation regimes had substantial significant influence on the yield attributing characters i.e., spikelets per spike, grains per spike, 1000 grain weight. Minute variation in spikelets per spike was observed (Table 1) due to availability of soil moisture up to flowering stage under different irrigation regimes (14, 15). Optimum moisture availability throughout all growth stages increased the number of fertile grains, which likely resulted in a higher number of grains per ear head under the  $I_1$  irrigation regimes. Whereas moisture stress as well as heat stress during the grain filling stage significantly reduced the number of grains in the spike (Table 1) in  $I_3$  (14, 16). Heat stress generally increases the no. of sterile spikelets in the ear (17). Reduction in 1000-grain weight mainly because of moisture stress at grain filling stage adversely affected the translocation of photosynthates to reproductive parts (18). Heat stress at this stage also had a negative effect on kernel weight (19, 20). Differences in ear length and effective tillers /  $m^2$  brought on by different irrigation regimes were not statistically significant since these two yield attributes were determined before flowering and frequent rainfall up to the flowering stage was unable to produce any variation in these parameters. So, water stress as well as terminal heat stress at later stages significantly affected translocation of carbohydrates from source to sink. Due to higher temperature increased respiration rates (21) and declined photosynthetic rate (22) resulted in decreased grain filling duration (23) and thereby reduction in grain yield (24).

The nitrogen-based treatments had substantial impact on all yield-attributing variables, including spike length, spikelets / spike, grains / spike, 1000 grain weight and effective bearing tillers /  $m^2$ , which resulted in significant effect on grain, straw and biological yield. Stimulatory effects of N on tillering through cytokinin synthesis resulted in an increase in the number of wheat plants with productive tillers (25). Greater competition for nutrients exists in the lower levels of N application resulted in lower effective tiller percentage (26). Nitrogen application at the stem elongation stage significantly increases productive tillers/ $m^2$  (27). In contrast, percentage effective tillers per square metre (table. 1) were the highest with  $N_3$  (50 % RDN+2 NUS at 45 and 70 DAS) compared to  $N_1$ . 100 % RDN, the better survival of tillers with  $N_3$  resulted in more ears per plant. This might be due to the application of nano-urea at the active tillering stage, which ensured better nitrogen availability when needed. An increase in N level from 80 to 150 kg/ha significantly improved the effective tillers per unit area (28). Ear length increased with increasing nitrogen levels from 0 to 120 kg N/ha (29). A high level of N nutrition during ear differentiation helped plants to produce more no. of spikelets in the spikes and spike length (14,30). A high amount of N nutrition during ear differentiation may prolong the activity of the apical dome, so the application of N fertilizer at the stem-elongation stage increases no. of grains per ear head (30). The greater availability of nitrogen at this stage is the key factor for better

yield (31, 32). Application of nano-urea at 45 and 70 DAS greatly helped florets to develop grain compared to other nano-urea treated plots. A 25.6 % reduction in grains/spike was observed in the control plot compared to sufficient nitrogen applied plots (33). Increase in photosynthetic activity and photosynthate translocation due to nitrogen fertilisation, improved photosynthate partitioning in yield attributes and generated more grains with larger sizes, which in turn increased yield. Nano-urea application at 45 DAS helped plants to develop more vegetative growth compared to other nano-urea treated plots where the application was done at 60 DAS. The higher grain yield is attributed to better yield components viz., grain number / ear, 1000 grain weight, HI and weight of grain/ear. Grain yield of wheat was closely related to the LAI, aboveground biomass accumulation (34) and its remobilization into reproductive parts (33). Nitrogen application helped to increase these yield attributes significantly (Table 1). Increasing post-anthesis biomass production is an effective approach to increase crop yield (29, 34). The interaction between the grain yield with nitrogen management and irrigation regimes was significant (Fig. 3) as availability of nitrogen is strongly correlated with irrigation regimes practices (15, 35). When nitrogen is limited, the plant's ability to produce leaves and store energy in the form of dry matter is restricted. This reduction in vegetative growth significantly impacted straw yield (34, 36). Nano-urea application at 45 DAS resulted in more vegetative growth compared to other treatments. The variation in straw yield may be attributed to different doses of nitrogen application during the vegetative growth stages. (37, 38). Adequate irrigation ensures that water stress is minimized, allowing plants to maintain their metabolic processes, enhances nutrient transport, reduces the no. of chaffy grains, all these contributed to higher HI (39).

### Protein content

Although non-significant variation was observed in the protein content of wheat due to different irrigation regimes. Protein levels generally increased when moisture availability decreased. This might be because of moisture stress circumstances lead to reduced carbohydrate production (40). As a result, more nitrogen was building up every grain of starch. The nitrogen content in the grain was diluted by the accumulated starch due to increased irrigation, resulting in a lower protein content (41). The amount of N applied had significant effect on the protein content of wheat grains. In the present study increased nitrogen application significantly increased the protein content of wheat grains by stimulating the accumulation of gliadins and glutenins (42). Nitrogen significantly increased the amounts of leucine, phenylalanine and TAA (Total Amino Acid) in wheat grains (43). In addition, higher nitrogen considerably increased the protein and amino acid content of wheat grain compared to low nitrogen application, as well as the amounts of TAA, EAA (Essential Amino Acids) and NAA (Non-essential Amino Acids) (44). Due to the high concentrations of glutamate (Glu) and alanine (Ala) in NAA, wheat grains accumulated more NAA than EAA when nitrogen was applied (44, 4).

### N, P, K content and uptake by crop

Nutrient uptake in crops is influenced by nutrient content and

productivity. The uptake of nitrogen, phosphorus and potassium by grain and straw were significantly affected by irrigation regimes due to variations in grain yield. 100 % RDN resulting in higher N, P and K content and uptake in grain and straw (Table 2 and 3). Probably the reason behind this is the numerous healthier roots and higher the density of roots, which may have contributed to better uptake of nitrogen, potassium and phosphorus from soil (45). The increase in total nitrogen uptake at the higher nitrogen rates may be explained by the greater nitrogen content in the grains and higher yield, which allowed them to accumulate more nitrogen (46). Higher N availability in the soil at high nitrogen levels enhanced N uptake by the crop which positively influenced N uptake through grain and straw. Probable reason behind higher K content in grain under  $N_3$  compared to  $N_1$  is application of nano-urea helps better translocation of K from straw to grain. That's N uptake of straw and grain K uptake was the highest in  $N_3$ . Higher P and K uptake due to higher N might be attributed to higher biological yield on higher doses of N application (46).

### Nitrogen use efficiency

Irrigation regimes didn't produce any effect on physiological nitrogen use efficiency ( $PE_N$ ) and physiological efficiency index of nitrogen (PEN) as both irrigation regimes possess the same effect on grain yield and uptake of nitrogen by wheat. Whereas apparent nitrogen recovery (ANR) and ANUE were higher with higher irrigation regimes because of higher yield with the same amount of nitrogen application (Table 4). Higher nitrogen use efficiency at higher irrigation levels, was attributed to better N mineralization and minimal nitrogen loss through leaching and volatilization at the optimal soil moisture condition, which ultimately led to better plant uptake of nitrogen and, consequently, growth and yield (47). Apparent recovery efficiency was significantly increased with increasing the irrigation levels (15). Among the NMPs, both the PEN and nitrogen efficiency ratio (NER) were higher in control as the yield as well as nitrogen uptake was less by the crop. Both ANR and ANUE were higher in  $N_3$  as the yield obtained from  $N_3$  was higher, but the application of N was comparatively less from  $N_1$ . Uptake of N was almost similar for both nano-urea and granular urea, but the amount of nitrogen was the main difference. In  $N_3$  amount of N was 60kg+100g/ha (60 kg from urea and 100g from nano urea) whereas in  $N_1$  it was 120 kg/ha but the uptake of N for these two treatments were 102.75 and 113.03 kg N/ha, respectively. As more uptake of N with less amount of nitrogen application so ANR was better for  $N_3$ . Long term nano-urea application ( $N_3$ ) is not sustainable as the plant uptake large amount of nitrogen from soil and make the soil more deficient in nitrogen. Higher doses of N reduce ANR and Physiological nitrogen use efficiency (34). Higher partial factor productivity (Table 4) was found with irrigation regime where higher number of irrigations were applied. Higher yield from higher irrigation regimes with the same amount of N, P and K fertilizer was the main reason for higher partial factor productivity of nutrients. Among the nitrogen management techniques, higher  $PPF_N$  was obtained with  $N_3$  due to low level of N (60 kg N through prilled urea+100g N from nano-urea) resulted in comparative higher grain yield. Higher doses of N

reduced  $PPF_N$  of the crop compared to lower doses (15).

### Stepwise regression analysis

Grain yield (Fig. 8) was highly correlated with no. of spikelet/spike, no. of grains/spike (48). The increment in grain yield per spike was in turn associated to a concomitant increase in grain numbers per spike 19 % increment, grain number (21 %) and grain yield per spikelet (25 %). As N uptake is the product of N concentration and dry matter yield, strong relationship always exists between them (49). Grain yield is also correlated with grain N and protein concentration (50).

### Conclusion

Based on the experiment, it was observed that applying irrigation at the milky stage of wheat in  $I_1$  and  $I_2$  irrigation regimes helped to prevent significant reduction in the yield of wheat variety DBW-187. Furthermore, nano-urea application at sub-optimum nitrogen doses did not adequately supply nitrogen compared to the standard 100 % RDN, resulting in significantly higher growth and grain yield with 100 % RDN. Increased nitrogen application enhanced both protein content and amino acid accumulation, underscoring the significance of nitrogen management for enhancing grain quality. Integrated approaches of combining increased irrigation and optimized nitrogen management led to higher partial factor productivity of nutrients, emphasizing the importance of sustainable practices for maximizing wheat productivity and nutrient efficiency. In conclusion, the most favourable combination for achieving higher wheat productivity in the eastern plateau region of India was 100 % RDN with three times irrigation.

### Acknowledgements

We extend our sincere gratitude to ICAR-IARI for providing the necessary facilities and resources. Special thanks go to the field staff and laboratory technicians for their invaluable assistance during the experimental setup and data collection.

### Authors' contributions

All authors have contributed to the paper on conceptualization, review & editing, supervision, investigation, data collection, formal analysis, writing - original draft: All authors reviewed the results and approved the final version of the manuscript.

### Compliance with ethical standards

**Conflict of interest:** The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

**Ethical issues:** None

### References

1. Anonymous. Economic Survey of India 2022-23. Directorate of Economics and Statistics, Government of India, New Delhi. 2023.
2. Mottaleb KA, Kruseman G, Frija A, Sonder K, Lopez-Ridaura S. Projecting wheat demand in China and India for 2030 and 2050:



- Implications for food security. *Frontiers in Nutrition*. 2023;9:1077443. <https://doi.org/10.3389/fnut.2022.1077443>
3. Anonymous. Economic Survey of India 2012-13. Directorate of Economics and Statistics, Government of India, New Delhi. 2013.
  4. Zhang-Xu Cheng, Zhu-Runshen, Xia-Fang Qin, Shangguan-Zhou Ping. Effects of irrigation at the critical growth stages of spring wheat plants and the period of severe drought in the hilly loess plateau of middle Gansu province, Shaanxi, China. *Journal of Triticeae Crops*. 2011;26(5):74-8. <https://doi.org/10.5555/20073191810>
  5. Fang WB, Sakih MK, Nudi YB. Effect of water stress with phytic development on yield of wheat grown in semi-arid environment. *Field Crops Research*. 2006;5(3):55-67. [https://doi.org/10.1016/0378-4290\(82\)90006-5](https://doi.org/10.1016/0378-4290(82)90006-5)
  6. Singh A, Prasad SM. Nanotechnology and its role in agro-ecosystem: A strategic perspective. *International Journal of Environmental Science and Technology*. 2016;14:2277-300. <https://doi.org/10.1007/s13762-016-1062-8>
  7. Donald CM. In search of yield. *J. Australian Institute of Agriculture Science*. 1962;28:171-78.
  8. Subbiah BV and Asija GL. A rapid procedure for the estimation of available nitrogen in soils. *Current Science*. 1956;25:259-60. <https://doi.org/10.5555/19571900070>
  9. Bray RH, Kurtz IT. Determination of total, organic and available forms of phosphorus in soils. *Soil Science*. 1945;59:39-46. <https://doi.org/10.1097/00010694-194501000-00006>
  10. Piper CS. Soil and plant analysis. Hans Publishers. Bombay. 1966.
  11. Jackson ML. Soil Chemical Analysis. Prentice Hall of India Pvt. Ltd. New Delhi. 1987:425-85.
  12. Jones DB. Factors for converting percentages of nitrogen in foods and feeds into percentages of protein. Washington, DC: US Department of Agriculture-circular. 1941:183.
  13. Cochran WG, Cox and GM. Experimental designs, Asia Publishing House, New Delhi. 1957.
  14. Si Z, Zain M, Mehmood F, Wang G, Gao Y, Duan A. Effects of nitrogen application rate and irrigation regime on growth, yield and water-nitrogen use efficiency of drip-irrigated winter wheat in the North China Plain. *Agricultural Water Management*. 2020;231:106002. <https://doi.org/10.1016/j.agwat.2020.106002>
  15. Pradhan S, Chopra UK, Bandyopadhyay KK, Singh R, Jain AK, Chand I. Effect of deficit irrigation and nitrogen levels on water productivity and nitrogen use efficiency of wheat (*Triticum aestivum*) in a semi-arid environment. *The Indian Journal of Agricultural Sciences*. 2014;87(7):887-91. <https://doi.org/10.56093/ijas.v84i7.42033>
  16. Dubey R, Pathak H, Chakrabarti B, Singh S, Gupta DK, Harit RC. Impact of terminal heat stress on wheat yield in India and options for adaptation. *Agricultural Systems*. 2020;181:102826. <https://doi.org/10.1016/j.agsy.2020.102826>
  17. Zhang C, Li G, Chen T, Feng B, Fu W, Yan J, et al. Heat stress induces spikelet sterility in rice at anthesis through inhibition of pollen tube elongation interfering with auxin homeostasis in pollinated pistils. *Rice*. 2018;11(1):14. <https://doi.org/10.1186/s12284-018-0206-5>
  18. Mondal S, Singh RP, Crossa J, Huerta-Espino J, Sharma I, Chatrath R, et al. Earliness in wheat: A key to adaptation under terminal and continual high temperature stress in South Asia. *Field Crops Research*. 2013;151:19-26. <https://doi.org/10.1016/j.fcr.2013.06.015>
  19. Barlow KM, Christy BP, O'Leary GJ, Riffkin PA, Nuttall JG. Simulating the impact of extreme heat and frost events on wheat crop production: A Review. *Field Crops Research*. 2015;171:109-19. <https://doi.org/10.1016/j.fcr.2014.11.010>
  20. Lobell DB, Sibley A, Ivan Ortiz-Monasterio J. Extreme heat effects on wheat senescence in India. *Nature Climate Change*. 2012;2:186-89. <https://doi.org/10.1038/nclimate1356>
  21. Chakrabarti B, Singh SD, Kumar V, Harit RC, Misra S. Growth and yield response of wheat and chickpea crops under high temperature. *Indian Journal of Plant Physiology*. 2013;18:7-14. <https://doi.org/10.1007/s40502-013-0002-6>
  22. Djanaguiraman M, Boyle DL, Welti R, Jagadish SV, Prasad PV. Decreased photosynthetic rate under high temperature in wheat is due to lipid desaturation, oxidation, acylation and damage of organelles. *BMC Plant Biology*. 2018;18(1): 01-17. <https://doi.org/10.1186/s12870-018-1263-z>
  23. Pathak H, Ladha JK, Aggarwal PK, Peng S, Das S, Singh Y, et al. Trends of climatic potential and on-farm yields of rice and wheat in the indo-gangetic plains. *Field Crops Research*. 2003;80:223-34. <https://doi.org/10.1016/s0378-429002.00194-6>
  24. Xie Y, Zhang H, Zhu Y, Li ZH, Yang JH, Cha FN, et al. Grain yield and water use of winter wheat as affected by water and sulfur supply in the North China Plain. *Journal of Integrative Agriculture*. 2017;16:614-25. <https://doi.org/10.1016/s2095-3119.16.61481-8>
  25. Kibe AM, Singh S, Kalra N. Water-nitrogen relationships for wheat growth and productivity in late sown conditions. *Agricultural Water Management*. 2006;84:221-28. <https://doi.org/10.1016/j.agwat.2006.02.010>
  26. Eftreuei A, Gooding M, White E, Spink J, Hackett R. Effect of nitrogen fertilizer application timing on nitrogen use efficiency and grain yield of winter wheat in Ireland. *Irish Journal of Agricultural and Food Research*. 2016;55:63-73. <https://doi.org/10.1515/ijafr-2016-0006>
  27. Kaur H, Ram H, Sikka R, Kaur H. Dry matter accumulation and nutrient uptake of wheat (*Triticum aestivum* L.) cultivars in relation to nitrogen levels. *Progressive Research*. 2013;8(1):24-28.
  28. Yousaf M, Fahad S, Shah AN, Shaaban M, Khan MJ, Sabiel SAI, et al. The effect of nitrogen application rates and timings of first irrigation on wheat growth and yield. *International Journal of Agriculture Innovations Research*. 2014;24:645-65.
  29. Singh SP and Singh HB. Effect of irrigation time and nitrogen level on wheat *Triticum aestivum* under late-sown condition of Western Uttar Pradesh. *Indian Journal of Agronomy*. 1991;36:41-42. <https://doi.org/10.5555/19920758747>
  30. Wang Y, Wang D, Tao Z, Yang Y, Gao Z, Zhao G, et al. Impacts of nitrogen deficiency on wheat (*Triticum aestivum* L.) grain during the medium filling stage: Transcriptomic and metabolomic comparisons. *Frontiers in Plant Science*. 2021;12:674433. <https://doi.org/10.3389/fpls.2021.674433>
  31. Wang Y, Liu CP, Chiu TS, Chen FT, Chang TH, Huang CN. Study on the differentiation and degeneration of florets in winter wheat and their relationship to nitrogen level. *Chung-kuo nung yeh k'o hsueh*.= *Scientia agricultura sinica*. 1979.
  32. Shi Y, Yu Z, Man J, Ma S, Gao Z, Zhang Y. Tillage practices affect dry matter accumulation and grain yield in winter wheat in the North China Plain. *Soil and Tillage Research*. 2016;160:73-81. <https://doi.org/10.1016/j.still.2016.02.009>
  33. Xu X, Zhang M, Li J, Liu Z, Zhao Z, Zhang Y, et al. Improving water use efficiency and grain yield of winter wheat by optimizing irrigations in the North China Plain. *Field Crops Research*. 2018;221:219-227. <https://doi.org/10.1016/j.fcr.2018.02.011>
  34. Liu W, Wang J, Wang C, Ma G, Wei Q, Lu H, et al. Root Growth, water and nitrogen use efficiencies in winter wheat under different irrigation and nitrogen regimes in North China Plain. *Frontiers in Plant Science*. 2018;9:1798. <https://doi.org/10.3389/fpls.2018.01798>



35. Shirazi SM, Yusop Z, Zardari NH, Ismail Z. Effect of irrigation regimes and nitrogen levels on the growth and yield of wheat. *Advances in Agriculture*. 2014;2014(1):1–6. <https://doi.org/10.1155/2014/250874>
36. Yang X, Lu Y, Ding Y, Yin X, Raza S, Tong YA. Optimising nitrogen fertilisation: A key to improving nitrogen-use efficiency and minimising nitrate leaching losses in an intensive wheat/maize rotation 2008–2014. *Field Crops Research*. 2017;206:1–10. <https://doi.org/10.1016/j.fcr.2017.02.016>
37. Guo J, Jia Y, Chen H, Zhang L, Yang J, Zhang J, et al. Growth, photosynthesis and nutrient uptake in wheat are affected by differences in nitrogen levels and forms and potassium supply. *Scientific Reports*. 2019;9(1):1248. <https://doi.org/10.1038/s41598-018-37838-3>
38. Luo C, Ma L, Zhu J, Guo Z, Dong K, Dong Y. Effects of nitrogen and intercropping on the occurrence of wheat powdery mildew and stripe rust and the relationship with crop yield. *Frontiers in Plant Science*. 2021;12:637393. <https://doi.org/10.3389/fpls.2021.637393>
39. Torrión JA, Stougaard RN. Impacts and limits of irrigation water management on wheat yield and quality. *Crop Science*. 2017;57:3239–51. <https://doi.org/10.2135/cropsci2016.12.1032>
40. Tayel MY, Mansour HA and Pibars SK. Effect of two sprinkler irrigation types on coefficient of variation CV and some quality properties of grain wheat. *Global Advanced Research Journal of Agricultural Science*. 2015;47:353–60.
41. Lollato RP, Figueiredo BM, Dhillon JS, Arnall DB, Raun WR. Wheat grain yield and grain-nitrogen relationships as affected by N, P and K fertilization: A synthesis of long-term experiments. *Field Crops Research*. 2019;236:42–57. <https://doi.org/10.1016/j.fcr.2019.03.005>
42. Zhang M, Ma D, Wang C, Zhao H, Zhu Y, Guo T. Responses of amino acid composition to nitrogen application in high and low protein wheat cultivars at two planting environments. *Crop Science*. 2016;56:1277–87. <https://doi.org/10.2135/cropsci2015.08.0504>
43. Ercoli L, Masoni A, Pampana S, Mariotti M, Arduini I. As durum wheat productivity is affected by nitrogen fertilisation management in Central Italy. *European Journal of Agronomy*. 2013;44:38–45. <https://doi.org/10.1016/j.eja.2012.08.005>
44. Fuertes-Mendizábal T, González-Torralba J, Arregui LM, González-Murua C, González-Moro MB, Estavillo JM. Ammonium as sole N source improves grain quality in wheat. *Journal of the Science of Food and Agriculture*. 2013;93:2162–71. <https://doi.org/10.1002/jsfa.6022>
45. Hassan A, Malik A, Ahmad S, Asif M, Mir SA, Bashir O, et al. Yield and nitrogen content of wheat (*Triticum aestivum* L.) as affected by FYM and urea in cold arid region of India. *International Journal of Current Microbiology and Applied Sciences*. 2018;7:328–32. <https://doi.org/10.20546/ijcmas.2018.702.043>
46. Sharma S, Kaur G, Singh P, Alamri S, Kumar R, Siddiqui MH. Nitrogen and potassium application effects on productivity, profitability and nutrient use efficiency of irrigated wheat (*Triticum aestivum* L.) *Plos One*. 2022;17(5):e0264210. <https://doi.org/10.1371/journal.pone.0264210>
47. Bandyopadhyay KK, Misra AK, Ghosh PK, Hati KM, Mandal KG, Moahnty M. Effect of irrigation and nitrogen application methods on input use efficiency of wheat under limited water supply in a vertisol of Central India. *Irrigation Science*. 2009;28:285–99. <https://doi.org/10.1007/s00271-009-0190-z>
48. Philipp N, Weichert H, Bohra U, Weschke W, Schulthess AW, Weber H. Grain number and grain yield distribution along the spike remain stable despite breeding for high yield in winter wheat. *Plos One*. 2018;13(10):e0205452. <https://doi.org/10.1371/journal.pone.0205452>
49. Ali AM, Ibrahim SM, Singh B. Wheat grain yield and nitrogen uptake prediction using ATLEAF and GreenSeeker portable optical sensors at jointing growth stage. *Information Processing in Agriculture*. 2019;7:375–83. <https://doi.org/10.1016/j.inpa.2019.09.008>
50. Bogard M, Allard V, Brancourt-Hulmel M, Machet JM, Jeuffroy MH, Gate P. Deviation from the grain protein concentration-grain yield negative relationship is highly correlated to post-anthesis N uptake in winter wheat. *Journal of Experimental Botany*. 2010;61:4303–12. <https://doi.org/10.1093/jxb/erq238>