



RESEARCH ARTICLE

Classification of different edible plants based on total phenolic and *in vitro* antioxidant activities using principal component and hierarchical cluster analysis

Khalid Hamid Musa^{1*}, Mochammad Amrun Hidayat², Bambang Kuswandi² & Aminah Abdullah³

¹Department of Food Science and Human Nutrition, College of Agriculture and Food, Qassim University, Buraydah 51452, Saudi Arabia

²Chemo and Biosensors Group, Faculty of Pharmacy, University of Jember, Jember 68121, Indonesia

³Department of Food Science, Faculty of Science and Technology, Universiti Kebangsaan Malaysia, Bangi 43600, Malaysia

*Correspondence email - k.musa@qu.edu.sa

Received: 01 February 2025; Accepted: 07 May 2025; Available online: Version 1.0: 18 May 2025

Cite this article: Khalid HM, Mochammad AH, Bambang K, Aminah A. Classification of different edible plants based on total phenolic and *in vitro* antioxidant activities using principal component and hierarchical cluster analysis. Plant Science Today (Early Access). <https://doi.org/10.14719/pst.7544>

Abstract

An acetone-aqueous extracts of 16 edible plants was prepared; their Total Phenolic Content (TPC), 2,2-diphenyl-1-picrylhydrazyl radical scavenging (DPPH), 2,2'-azino-bis (3-ethylbenzthiazoline-6-sulfonic acid) radical scavenging (ABTS), Oxygen Radical Absorbance Capacity (ORAC) and Ferric Reducing Antioxidant Power (FRAP) were measured. Principal Component Analysis (PCA) and Hierarchical Cluster Analysis (HCA) were used to clarify the correlations between antioxidant assays and plant extracts. The first (PC1) and the second (PC2) principal components accounted for 91.60 % of the overall variability, whereas HCA classified samples into three main groups based on measured parameters. The 16 plants were classified into high, medium and low antioxidant activity groups based on according to TPC and the antioxidant activities results. Those in the high-antioxidant activity group were *Punica granatum*, *Rheum officinale*, *Phyllanthus niruri* and *Melaleuca leucadendra*. The plants in the medium-antioxidant activity group were *Thymus vulgaris*, *Datura metel*, *Menthae arvensis*, *Parameria laevigata*, *Glycyrrhiza glabra*, *Curcuma xanthorrhiza* and *Zingiber officinale*. The plants in the low-antioxidant activity group were *Plantago major*, *Guazuma ulmifolia*, *Centella asiatica*, *Sericocalyx crispus* and *Piper nigrum*. Positive relationships existed between TPC and the antioxidant activity of the extracts, indicating that the principal contributors to the antioxidant characteristics of the studied plants are phenolic compounds. Some of these plants may serve as valuable sources of natural antioxidants.

Keywords: antioxidant activity; hierarchical cluster analysis; Principal Component Analysis; Total Phenolic Content

Introduction

Reactive Oxygen Species (ROS) are detrimental by-products generated during cellular respiration. Excess ROS in cells can damage proteins, lipids and DNA, cumulatively resulting in oxidative stress (1). Oxidative stress, characterised by an imbalance favouring oxidants over antioxidants, has been proposed as a contributing factor to ageing and numerous diseases in humans (2, 3). Antioxidant activity reflects the power of a compound to stop oxidative damage. Dietary antioxidants can enhance cellular defences and protect biological components from oxidative damage (4). Natural antioxidants have received much attention recently because of their scavenging properties (5) and because they are natural products, they are favoured by consumers (6). Like synthetic antioxidants, natural antioxidant containing higher phenolic compounds can scavenge free radicals effectively, chelate transition metals and absorb ultraviolet light, so stopping progressive oxidative damage and preventing undesirable odours and flavours (7). Recently, there have been considerable efforts to identify new natural sources of active antioxidant compounds (8). Phenolic-rich plant

extracts are utilised in the food industry to enhance food quality by inhibiting the oxidative deterioration of lipids. There are two types of *in vitro* antioxidant activity assays: those based on Hydrogen Atom Transfer (HAT) and those based on Electron Transfer (ET). In HAT assays, the thermally generated peroxy radicals will be a target to compete between the antioxidant and substrate and ORAC assay is an example. ET assays measure the capacity of a compound to reduce an oxidant and resulted in a color change. The antioxidant concentration in sample is correlated with extent of the change in color. Examples of ET-based assays include DPPH, ABTS and FRAP (9). Due to different mechanisms of antioxidant action; antioxidant activity for a sample should be evaluated using several methods (10). PCA is a statistical technique which basically reduces the dimensionality of a large datasets in an interpretable way. The aim of this study was to analyse extracts prepared from 16 edible plants commonly used in Indonesia and worldwide for their total phenolic content and antioxidant activity. PCA was used to report the details variations in the antioxidant activity among the studied plants and to classify them.

Material and Methods

Chemicals

Sodium carbonate was from RDH (Germany), glacial acetic acid from Mallinckrodt Baker (USA), Trolox (6-hydroxy-2,5,8-tetramethylchroman-2-carboxylic acid), fluorescein disodium salt (FL), sodium acetate trihydrate, 2,2-Diphenyl-1-picrylhydrazyl (DPPH), TPTZ (2,4,6-tris(2-pyridyl)-s-triazine), gallic acid, 2,2'-azo-bis(2-amidinopropane) dihydrochloride, (AAPH), 2,2'-azino-bis(3-ethylbenzothiazoline-6-sulphonic acid (ABTS) and potassium persulfate were purchased from Sigma (USA), Folin-Ciocalteu reagent, ferric chloride ($\text{FeCl}_3 \cdot 6\text{H}_2\text{O}$) and HCl were obtained from Merck, (Germany). All chemicals and reagents used in this study were of analytical grade.

Preparation of plant extracts

Plant samples were collected from Jember District, East Java; all plants were properly authenticated by the Indonesian Institute of Sciences at Purwodadi Botanical Garden, East Java, Indonesia. Prior to extraction, the materials were cleaned, diced, dried for three days and grinded into a fine powder. Table 1 presents the common names, scientific names, family classifications and the part used for extracts preparation of all plants. Following the methods described previously (11), 0.1 g of dried plant powder were extracted with 10 mL of 70 % aqueous acetone using a magnetic stirrer at 1000 rpm (IKA, Germany). Extract was centrifuged for 10 min at 4750 g using thermo-line, a mini centrifuge (China) and the collected supernatants were used.

Determination of Total Phenolic Content (TPC)

TPC was reported following the method of Slinkard and Singleton (12). Folin-Ciocalteu reagent (500 μL) was added to each plant extract, gallic acid standard, or blank (100- μL), followed by the addition of 1 mL sodium carbonate (7.5 %, w/v). Results were compared to gallic acid at absorbance of 765 nm after 120 min of incubation using microplate reader spectrophotometer (Biotek, USA).

Determination of Ferric Reducing Antioxidant Power (FRAP) assay

The FRAP assay method (13) was followed to determine reducing/antioxidant power. FRAP reagent was prepared as using 300 mM acetate buffer, 10mM TPTZ (2,4,6- tri(2-pyridyl)-s-triazine) in 40 mM HCl and 20mM $\text{FeCl}_3 \cdot 6\text{H}_2\text{O}$ in the ratio of 10:1:1. For assays, 1900 μL FRAP reagent was mixed with 100 μL of each plant extract, Trolox standard, or blank and incubated for 30 min before reading absorbance at 595 nm using microplate reader spectrophotometer (Biotek, USA).

Determination of DPPH radical scavenging activity assay

The DPPH assay method (14) was followed to determine radical scavenging activity. A 3 mL of DPPH solution (40 mg/L) was mixed with 100 μL of each plant extract, Trolox standard or blank and incubated for 30 min before reading absorbance at 517 nm using microplate reader spectrophotometer (Biotek, USA).

Determination of ABTS radical scavenging activity

The ABTS method (15) was followed to determine radical scavenging activity. ABTS radical cations were generated by oxidizing 2.45 mM potassium persulfate and 7 mM ABTS in the dark. Using distilled water; ABTS solution was adjusted to an absorbance of 1 ± 0.02 at 734 nm. A 100 μL of each plant extract, Trolox standard, or blank was added to 1 mL of ABTS solution and incubated for 5 min before reading absorbance using microplate reader spectrophotometer (Biotek, USA).

Determination of Oxygen Radical Absorbance Capacity (ORAC)

A 150 μL fluorescein (10 nM) was mixed with 25 μL Trolox (6.5 μM to 50 μM), buffer (blank), or sample in 96-well black plate (Thermo Scientific, USA). An automatic injector was used to add 25 μL AAPH (153 mM) to the solution. The measurement was conducted at 485 nm excitation wavelength and 520 nm emission wavelengths. ORAC values were calculated by subtracting the area under the curve (AUC) of the blank from the AUC of each sample and by comparison with the Trolox standard curve. The absorbance of solutions was measured using FLUO star Omega microplate fluorescence reader (BMG Labtech, Germany).

Table 1. Common name, scientific name, family name of the plants collected from Jember, East Java, Indonesia

No.	Common name	Scientific name	Family name	Part used
1	Stonebreaker	<i>Phyllanthus niruri</i>	Euphorbiaceae	Aerial part
2	Thyme	<i>Thymus vulgaris</i>	Lamiaceae	Aerial part
3	Pomegranate	<i>Punica granatum</i>	Punicaceae	Fruit rind
4	Centella	<i>Centella asiatica</i>	Umbelliferae	Aerial part
5	Bastard cedar	<i>Guazuma ulmifolia</i>	Sterculiaceae	Leaves
6	Yellow strobilanthus	<i>Sericocalyx crispus</i>	Acanthaceae	Leaves
7	Broadleaf plantain	<i>Plantago major</i>	Plantaginaceae	Leaves
8	Black pepper	<i>Piper nigrum</i>	Piperaceae	Fruit
9	Cajeput tree	<i>Melaleuca leucadendra</i>	Myrtaceae	Leaves
10	Devil's trumpet	<i>Datura metel</i>	Solanaceae	Leaves
11	Wild mint	<i>Mentha arvensis</i>	Lamiaceae	Aerial part
12	Ginger	<i>Zingiber officinale</i>	Zingiberaceae	Rhizome
13	Tight wood	<i>Parameria laevigata</i>	Apocynaceae	Wood
14	Liquorice	<i>Glycyrrhiza glabra</i>	Leguminosae	Root
15	Rhubarb	<i>Rheum officinale</i>	Polygonaceae	Root
16	Java Ginger	<i>Curcuma xanthorrhiza</i>	Zingiberaceae	Rhizome

Statistical analysis

Results are expressed as means with Standard Deviation (SD) of three measurements. The significance of differences of $p < 0.05$ among mean values was evaluated by Analysis of Variance (ANOVA) using MINITAB® (16.00) software. Correlation analyses were performed using Pearson's correlation coefficient (r). PCA and HCA were applied to the data for classification.

Results and Discussion

As can be seen in Table 2 the TPC varied clearly among plant extracts tested. The range of the TPC value is from 126 to 6833 mg GAE /100 g (DW), a dissimilarity of almost 54-fold between the TPC values of the samples. *R. officinale*, *P. granatum*, *M. leucadendra* and *P. niruri* showed the highest TPC values (>5000 mg GAE/100 g DW) and *P. nigrum* showed the lowest value. FRAP values of the studied samples ranged from 246 to 20969 mg TE /100 g DW, a dissimilarity of nearly 85-fold between the FRAP values. *P. granatum*, *R. officinale*, *M. leucadendra* and *P. niruri*, also showed the highest antioxidant activities in the FRAP assay. *P. nigrum* and *P. major* showed the lowest FRAP values.

Table 2 showed the DPPH radical scavenging activities of the samples. The DPPH value ranged from 4377 to 41156 mg TE/100 g DW, with a difference of about 9-fold between the DPPH scavenging activities. *P. granatum* showed the highest antioxidant activity, followed by *R. officinale*, *P. niruri* and *M. leucadendra*, while the lowest antioxidant activity was shown by *P. nigrum*. ABTS radical scavenging activities are shown in Table 3. The activities determined in this assay ranged from 24 to 1741 mg TE/100 g DW, representing the largest difference between the lowest and highest activities (approximately 73-fold) among the four assays. *P. niruri*

showed the highest antioxidant activity, followed by *M. arvensis*, *P. laevigata*, *M. leucadendra*, *P. granatum* and *R. officinale*. Consistent with the results of the TPC, FRAP and DPPH assays, the *Piper nigrum* extract showed the lowest ABTS value (24 mg TE/100 g DW). The ORAC values are shown in Table 3, values ranged from 9997 to 69035 mg TE /100 g DW, showing about 7-fold variation between the lowest and highest ORAC values. The extract from *R. officinale* showed the highest ORAC value, followed by those from *Z. officinale*, *P. granatum* and *P. niruri*. *C. asiatica* and *G. ulmifolia* extracts showed the lowest ORAC values.

Several studies have evaluated the antioxidant activity of *R. officinale* (16-18) and *P. niruri* (19-21). *P. granatum* is one of the oldest known edible fruits and its antioxidant activity has been determined in other studies (22, 23). *M. leucadendra* is considered one of the plants with the highest antioxidant activities (24). Other studies have analyzed the antioxidant activities of *T. vulgaris* (25), *D. metel* (26), *M. arvensis* (27), *C. xanthorrhiza* (28-30), *G. glabra* (31, 33), *P. laevigata* (34) and *Z. officinale* (35, 36). Other studies have investigated the antioxidant activities of *C. asiatica* (37,38); *P. major* (39, 40), *G. ulmifolia* (41) and *S. crispus* (42). High antioxidant and radical scavenging activities were reported for *P. nigrum* in other studies (43, 44), but in this study where the *P. nigrum* extract showed low antioxidant activity.

The correlation coefficient between total phenolic content and antioxidant activities

Table 3 displays the correlation coefficient (R) values between TPC and antioxidant activities. TPC was strongly correlated with FRAP and with DPPH. TPC values for all plant extracts showed a medium correlation with ABTS and a good correlation with ORAC. These results show the importance of phenolic compounds in the radical scavenging activity as well

Table 2. Total Phenolic Content (TPC), Ferric Reducing Antioxidant Power (FRAP), DPPH radical scavenging activity (DPPH), ABTS radical scavenging activity (ABTS) and Oxygen Radical Absorbance Capacity (ORAC) in different plant extracts

Plant name	TPC ^a	FRAP ^b	DPPH ^b	ABTS ^b	ORAC ^b
<i>P. niruri</i>	5270.85 ± 275.92 ^D	9617.19 ± 1296.49 ^D	29501.2 ± 0 ^D	1740.77 ± 1.99 ^A	39550 ± 1552 ^D
<i>T. vulgaris</i>	2195.74 ± 93.07 ^H	3204.12 ± 123.95 ^G	14923 ± 0 ^G	1707.57 ± 11.28 ^{AB}	19248 ± 862 ^{GH}
<i>P. granatum</i>	6477.69 ± 27.18 ^B	20968.8 ± 1060.88 ^A	41156.3 ± 106.18 ^A	1732.47 ± 1.99 ^A	47439 ± 3537 ^C
<i>C. asiatica</i>	826.39 ± 14.67 ^L	1101.27 ± 60.78 ^{HI}	8529.11 ± 270.4 ^J	1148.96 ± 75.58 ^D	9997 ± 943 ^L
<i>G. ulmifolia</i>	926.91 ± 23.79 ^{KL}	1339.65 ± 72.49 ^{HI}	7624.05 ± 43.85 ^K	1170.75 ± 50.9 ^D	11287 ± 759 ^L
<i>S. crispus</i>	794.77 ± 47.79 ^L	1253.39 ± 29.98 ^H	6782.28 ± 95.01 ^L	988.69 ± 57.53 ^E	18284 ± 1100 ^{HI}
<i>P. major</i>	973.39 ± 7.4 ^{KL}	831.97 ± 73.36 ^H	7212.66 ± 197.32 ^K	1004.77 ± 32.32 ^E	12195.5 ± 1200.5 ^{KL}
<i>P. nigrum</i>	126.01 ± 6.42 ^M	245.5 ± 14.22 ^I	4377.22 ± 7.31 ^M	24.48 ± 3.59 ^F	11590 ± 276 ^{KL}
<i>M. leucadendra</i>	5893.11 ± 95.2 ^C	13803.8 ± 411.1 ^C	29018.4 ± 291.99 ^C	1735.06 ± 4.15 ^A	31172 ± 1188.19 ^E
<i>D. metel</i>	1467.54 ± 104.88 ^J	1520.74 ± 164.41 ^H	9117.72 ± 14.62 ^{HI}	1448.24 ± 16.28 ^C	13515 ± 9 ^{JK}
<i>M. arvensis</i>	3425.18 ± 202.54 ^F	5069.72 ± 526.84 ^F	13342.5 ± 325.17 ^F	1738.69 ± 1.99 ^A	32280 ± 416 ^E
<i>Z. officinale</i>	1100.36 ± 40.84 ^K	2839.39 ± 88.99 ^G	8836.78 ± 165.91 ^I	1679.56 ± 22.18 ^B	50496 ± 2672 ^B
<i>P. laevigata</i>	4460.21 ± 151.55 ^E	6312.66 ± 68.17 ^E	18549.4 ± 537.53 ^D	1738.17 ± 5.49 ^A	21535 ± 979 ^G
<i>G. glabra</i>	3158.75 ± 152.42 ^G	1671.8 ± 63.38 ^H	10647.1 ± 212.36 ^G	1727.28 ± 3.54 ^{AB}	26045 ± 549 ^F
<i>R. officinale</i>	6833.73 ± 175.52 ^A	18686.9 ± 992.71 ^B	32593.1 ± 491.08 ^B	1731.95 ± 3.59 ^A	69034.67 ± 1415.66 ^A
<i>C. xanthorrhiza</i>	1817.5 ± 12.32 ^I	1540.85 ± 78.31 ^H	9560.92 ± 205.72 ^{HI}	1656.22 ± 55.85 ^B	15745 ± 1295 ^{IJ}
Minimum	126.010	245.500	4377.220	24.480	9997.000
Maximum	6833.730	20968.800	41156.300	1740.770	69034.670
Mean	2859.258	5625.484	15735.734	1435.852	26838.323
Std. deviation	2259.711	6644.706	11152.196	471.768	17140.787

There are no significant differences among values marked with the same superscript capital letters in individual columns

^a mg gallic acid equivalents (GAE) per 100 g DW of sample

^b mg Trolox equivalents (TE) per 100 g DW of sample

Table 3. Correlations (R^2) between Total Phenolic Content (TPC) and antioxidant values obtained using Ferric Reducing Antioxidant Power (FRAP), DPPH radical scavenging activity (DPPH), ABTS radical scavenging activity (ABTS) and Oxygen Radical Absorbance Capacity (ORAC)

	TPC	FRAP	DPPH	ABTS	ORAC
TPC	-	0.924	0.948	0.655	0.739
FRAP	-	-	0.971	0.483	0.785
DPPH	-	-	-	0.557	0.724
ABTS	-	-	-	-	0.543


as the reducing ability of the extracts. There were strong correlations among the values obtained using the various assay methods. In particular, the values obtained using the DPPH and FRAP were very strongly correlated. ORAC showed a good correlation with FRAP and DPPH. The weakest correlations were between the ABTS and those obtained using FRAP, DPPH and ORAC. These differences might result from the different antioxidant mechanisms detected in each assay and this should be studied further.

Positive relationships between TPC, free radical scavenging activity and ferric reducing power have been reported in other studies (45, 46). However, some researchers found no or only weak correlations between phenolic contents and antioxidant activities (47) and between different antioxidant assay methods (48). The link between antioxidant activities and TPC could be affected by the various testing methods and by other substances with antioxidant properties,

such as ascorbic acid, sugars and carotenoids, present in the extracts. (49).

Principal Component Analysis

PCA was applied to the standardized results of the 16 plants and 79.359 % of the variability in the dataset was explained by the first principal component (PC1). This significant proportion indicates that the PC1 has successfully captured most of the patterns in the data. PC2 (second principal component) adds another 12.237 % to the total explained variability, bringing the cumulative total to 91.60 %. Adding the third principal component (PC3) into the mix raises the cumulative percentage to 98.62 %. The last two components, PC4 and PC5, contribute very little to the explained variability (only about 1.383 % collectively). Fig. 1 shows a sample score plot for PC1 vs PC2. *P. granatum* and *R. officinale* are located on the righthalf of the plot and *P. nigrum* is located away from other samples. This indicates that certain antioxidant

**Fig. 1.** Principal Component Analysis (PCA) scores plot for different plants and loading plots for different variables on PC1 and PC2. The locations of the plants in the plot are based on their antioxidant activity.

TPC: Total Phenolic Content; FRAP: Ferric Reducing/Antioxidant Power; DPPH: DPPH radical scavenging activity; ABTS: ABTS radical scavenging activity; ORAC: Oxygen Radical Absorbance Capacity; PC: Principal component analysis factors

characteristics are considerably different from those of the other samples.

The plot clusters TPC, FRAP, DPPH and ORAC on the right side, while ABTS occupies the upper right-hand side. Fig. 1 illustrates how the antioxidant activity of the plants determines their placement. The positioning of *P. granatum* in the lower right quadrant of Fig. 1 may be attributed to their elevated antioxidant activities, which are situated in this area of the principal component space. In contrast, *P. nigrum*, characterised by low antioxidant levels, is positioned in the lower left quadrant of Fig. 1 diametrically opposite to *P. granatum*.

The relations between TPC, FRAP, DPPH and ORAC and PCs are shown as evidenced by their Pearson correlation coefficients (Table 4). PC1 positively correlates with TPC, FRAP, DPPH, ORAC and moderately with ABTS, suggesting it essentially represents the variance in these variables.

The second component (PC2) has a strong positive relationship with ABTS and a moderate negative relationship with FRAP and DPPH, meaning it reflects different information, especially related to ABTS. PC3 has a notable positive correlation with ORAC, capturing some unique variance not explained by PC1 or PC2.

The fourth and fifth components (PC4 and PC5) show low correlation with all variables, indicating their small contribution to the total variance. This result implies that most of the original variables are significantly associated with the first three principal components, corroborating their contribution to over 98 % of the total variability.

The squared cosines of the variables in PCA, presented in Table 5, show how well each principal component represents each variable. TPC, FRAP and DPPH have high squared cosines with PC1 (0.933, 0.906 and 0.917 respectively),

signifying that this component predominantly represents them. PC1 and PC2 approximately equally represent ABTS, with squared cosines of 0.492 and 0.503, respectively, indicating that these components split its variance. ORAC's variance is mainly accounted for by PC1 and PC3, with squared cosines of 0.720 and 0.278 respectively. Low squared cosines show that PC4 and PC5 contribute little to the total variance and have weak connections with the original variables.

The given factor scores (Table 6) are the coordinates of the original observations (different plants) in the space defined by the principal components (PC1-PC5), indicating how each plant corresponds to the principal components. Plants like *R. officinale*, *P. granatum*, *P. niruri* and *M. leucadendra*, show high positive scores on PC1, implying high values for the variables most strongly associated with this component, while *Piper nigrum* shows a notably negative score, indicating low values for the same variables. Similarly, *C. xanthorrhiza*, *G. glabra*, *Z. officinale* and *T. vulgaris* present positive scores on PC2, suggesting high values for variables associated with PC2. Despite PC3-PC5 accounting for a smaller proportion of the total variance, they also provide valuable insights. Factor scores are instrumental for comparing observations and identifying patterns, with similar scores implying similarities in original variable values.

Table 2 also summarizes the statistics representing five variables, all with 16 observations and no missing data. The variables TPC, FRAP, DPPH and ORAC all have high standard deviations, suggesting substantial variability in these measures among the samples. TPC and FRAP have lower average values compared to DPPH and ORAC, which have the highest mean values, indicating they generally have higher measures in the dataset. In contrast, ABTS has the lowest mean value and least variability, suggesting it has more consistent, lower measurements across the samples.

Table 4. Correlations between TPC, FRAP, DPPH, ABTS radical scavenging activity (ABTS), ORAC and the PCA factors

	PC1	PC2	PC3	PC4	PC5
TPC	0.966	-0.033	-0.175	-0.186	0.007
FRAP	0.952	-0.271	-0.065	0.087	0.093
DPPH	0.957	-0.179	-0.189	0.083	-0.092
ABTS	0.702	0.709	-0.050	0.045	0.012
ORAC	0.848	-0.042	0.527	-0.017	-0.018

Table 5. Squared cosines of the variables (TPC, FRAP, DPPH, ABTS and ORAC)

	PC1	PC2	PC3	PC4	PC5
TPC	0.933	0.001	0.031	0.035	0.000
FRAP	0.906	0.074	0.004	0.008	0.009
DPPH	0.917	0.032	0.036	0.007	0.008
ABTS	0.492	0.503	0.002	0.002	0.000
ORAC	0.720	0.002	0.278	0.000	0.000

TPC: Total Phenolic Content

FRAP: Ferric Reducing/Antioxidant Power

DPPH: DPPH radical scavenging activity

ABTS: ABTS radical scavenging activity

ORAC: Oxygen Radical Absorbance Capacity

PC: Principal Component Analysis Factors

Table 6. Factor scores for the sixteen plant extracts indicating how each plant extract corresponds to the principal components (PC1 to PC5)

	PC1	PC2	PC3	PC4	PC5
<i>Phyllanthus niruri</i>	2.005	0.010	-0.176	-0.123	-0.437
<i>Thymus vulgaris</i>	-0.348	0.725	-0.302	0.228	-0.113
<i>Punica granatum</i>	3.830	-0.915	-0.453	0.460	0.015
<i>Centella asiatica</i>	-1.761	-0.079	-0.285	0.199	0.004
<i>Guazuma ulmifolia</i>	-1.711	-0.035	-0.211	0.148	0.084
<i>Sericocalyx crispus</i>	-1.745	-0.394	0.242	0.054	0.033
<i>Plantago major</i>	-1.862	-0.332	-0.117	0.011	0.017
<i>Piper nigrum</i>	-2.991	-2.168	0.240	-0.223	-0.068
<i>Melaleuca leucadendra</i>	2.213	-0.202	-0.765	-0.087	0.133
<i>Datura metel</i>	-1.240	0.456	-0.263	0.117	0.056
<i>Menthae arvensis</i>	0.351	0.653	0.240	-0.221	0.120
<i>Zingiber officinale</i>	-0.109	0.739	1.712	0.257	-0.050
<i>Parameria laevigata</i>	0.628	0.489	-0.651	-0.309	0.034
<i>Glycyrrhiza glabra</i>	-0.249	0.897	0.081	-0.392	-0.035
<i>Rheum officinale</i>	3.913	-0.688	0.949	-0.198	0.146
<i>Curcuma xanthorrhiza</i>	-0.924	0.845	-0.242	0.081	0.060

Dendrogram of HCA of plants extract

Dendrogram of HCA of plants extract is shown in Fig. 2; three well-defined clusters are grouped in based on their similarity. *M. leucadendra*, *P. niruri*, *P. granatum* and *R. officinale* comprise a group of samples (A) associated with high antioxidant activity. A second cluster (B) consists of *T. vulgaris*, *D. metel*, *M. arvensis*, *Z. officinale*, *P. laevigata*, *G. glabra* and *C. xanthorrhiza* because of their medium

antioxidant activity. A third cluster (C) includes *C. asiatica*, *G. ulmifolia*, *S. crispus*, *P. major* and *Piper nigrum*.

HCA results agree with PCA results indicating that the samples are distinct from others. It was observed that the plants with high antioxidant activity had TPC > 5000 mg GAE/100 g DW, FRAP > 9000, DPPH > 25000, ABTS > 400 and ORAC > 40000 (mg TE/ 100g DW). Medium antioxidant activity group had TPC ranging from 5000 to 1000 mg GAE/100 g DW, FRAP from 9000 to 1000, DPPH from 9000 to 25000, ABTS from 400 to 1000 and ORAC from 4000 to 15000 (mg TE/ 100g DW). The plants with low antioxidant activity had TPC < 1000




Fig. 2. Dendrogram of Hierarchical Cluster Analysis (HCA) of plants extract (The sample code as in Table 1). Three well-defined clusters are grouped in based on their similarity; (A): a group of samples associated with high antioxidant activity, (B): a second cluster for medium antioxidant activity group, (C): a cluster includes low antioxidant activity group.

mg GAE/100 g DW, FRAP < 1000, DPPH < 9000, ABTS < 100 and ORAC < 15000 (mg TE/ 100g DW).

Conclusion

The total phenolic contents and antioxidant activities of 16 plant extracts were differed significantly. The studied plants showing strong antioxidant activities are widely distributed, making them a promising source of natural antioxidants for food industries. The extracts from *P. granatum*, *R. officinale*, *P. niruri* and *M. leucadendra* showed the strongest antioxidant activities among the tested plants. The antioxidant activities of these plants are linked to their phenolic compounds, as higher levels of phenolic content were associated with stronger antioxidant activities. PCA and HCA facilitated the visualisation of the dataset and the underlying observed grouping relationships. Easy interpretation of differences and similarities in antioxidant activity of plants could be achieved by using a combination of sample characterization and multivariate data analysis. PCA and HCA provide powerful insights into the antioxidant profiles among different plants. Further investigations are required to identify the active compounds in these plant extracts and to demonstrate their biological activities.

Acknowledgements

The authors would like to acknowledge Progene Link Sdn Bhd (Malaysia) for lending the FLUOstar Omega microplate fluorescence reader (BMG Labtech) to perform the ORAC assay.

Authors' contributions

BK contributed the concepts. AA and KHM provided the overall design. BK, KHM and MAH verified the methodology part of the study. KHM and MAH checked the data curation. The original draft of the work was prepared and wrote by KHM and AA. BK, KHM and MAH edited the review part of the manuscript. All authors read and approved the final manuscript.

Compliance with ethical standards

Conflict of interest: Authors do not have any conflict of interests to declare.

Ethical issues: None

Data Availability: The datasets used and analyzed during the current study would be available from the corresponding author on request.

References

1. Zandi P, Schnug E. Reactive oxygen species, antioxidant responses and implications from a microbial modulation perspective. *Biol.* 2022;11(2):155. <https://doi.org/10.3390/biology11020155>
2. Hajam YA, Rani R, Ganie SY, Sheikh TA, Javaid D, Qadri SS, et al. Oxidative stress in human pathology and aging: Molecular mechanisms and perspectives. *Cells.* 2022;11(3):552. <https://doi.org/10.3390/cells11030552>
3. Sharifi-Rad M, Anil Kumar N V., Zucca P, Varoni EM, Dini L, Panzarini E, et al. Lifestyle, oxidative stress and antioxidants: back and forth in the pathophysiology of chronic diseases. *Front Physiol.* 2020;11:1-21. <https://doi.org/10.3389/fphys.2020.00694>
4. Pisoschi AM, Pop A, Iordache F, Stanca L, Predoi G, Serban AI. Oxidative stress mitigation by antioxidants - An overview on their chemistry and influences on health status. *Eur J Med Chem.* 2021;209:112891. <https://doi.org/10.1016/j.ejmech.2020.112891>
5. Parcheta M, Świsłocka R, Orzechowska S, Akimowicz M, Chojńska R, Lewandowski W. Recent developments in effective antioxidants: The structure and antioxidant properties. *Materials (Basel).* 2021;14(8):1984. <https://doi.org/10.3390/ma14081984>
6. Ghosh N, Chatterjee S, Sil PC. Evolution of antioxidants over times (including current global market and trend). In: Seyed MN, Ana SS, editors. *Antioxidants effects in health.* Elsevier. 2022:3-32. <https://doi.org/10.1016/B978-0-12-819096-8.00011-2>
7. Zeb A. Concept, mechanism and applications of phenolic antioxidants in foods. *J Food Biochem.* 2020;44(9):e13394. <https://doi.org/10.1111/jfbc.13394>
8. Lourenço SC, Moldão-Martins M, Alves VD. Antioxidants of natural plant origins: from sources to food industry applications. *Molecules.* 2019;24(22):4132. <https://doi.org/10.3390/molecules24224132>
9. Munteanu IG, Apetrei C. Analytical methods used in determining antioxidant activity: A review. *Int J Mol Sci.* 2021;22(7):3380. <https://doi.org/10.3390/ijms22073380>
10. Gulcin İ. Antioxidants and antioxidant methods: An updated overview. *Arch Toxicol.* 2020;94(3):651-715. <https://doi.org/10.1007/s00204-020-02689-3>
11. Musa KH, Abdullah A, Jusoh K, Subramaniam V. Antioxidant activity of pink-flesh guava (*psidium guajava* L.): Effect of extraction techniques and solvents. *Food Anal Methods.* 2011;4(1):100-07. <https://doi.org/10.1007/s12161-010-9139-3>
12. Slinkard K, Singleton VL. Total phenol analysis: Automation and comparison with manual methods. *Am J Enol Vitic.* 1977;28(1):49-55. <https://doi.org/10.5344/ajev.1977.28.1.49>
13. Elshaafi IM, Musa KH, Abdullah Sani N. Effect of oven and freeze drying on antioxidant activity, total phenolic and total flavonoid contents of fig (*Ficus carica* L.) leaves. *Food Res.* 2020;4(6):2114-21. [https://doi.org/10.26656/fr.2017.4\(6\).072](https://doi.org/10.26656/fr.2017.4(6).072)
14. Ismail M, Abdallah EM, Musa KH, Abdellatif AAH. First report of some chemical and biological properties of two wild honey types collected from a remote area in Southern Darfur, Sudan. *Food Res.* 2023;7(3):203-13. [https://doi.org/10.26656/fr.2017.7\(3\).360](https://doi.org/10.26656/fr.2017.7(3).360)
15. Re R, Pellegrini N, Proteggente A, Pannala A, Yang M, Rice-Evans C. Antioxidant activity applying an improved ABTS radical cation decolorization assay. *Free Radic Biol Med* 1999;26(9-10):1231-37. [https://doi.org/10.1016/S0891-5849\(98\)00315-3](https://doi.org/10.1016/S0891-5849(98)00315-3)
16. Dai L-X, Miao X, Yang X-R, Zuo L-P, Lan Z-H, Li B, et al. High value-added application of two renewable sources as healthy food: The nutritional properties, chemical compositions, antioxidant and anti-inflammatory activities of the stalks of *Rheum officinale* Baill. and *Rheum tanguticum* Maxim. ex Regel. *Front Nutr.* 2022;8:770264. <https://doi.org/10.3389/fnut.2021.770264>
17. Yuan S, Jian T, Li W, Huang Y. Extraction process optimization and activity assays of antioxidative substances from *Rheum officinale*. *J Food Meas Charact.* 2020;14(1):176-84. <https://doi.org/10.1007/s11694-019-00279-1>
18. Dai L, Miao X, Ma Y, Yang X, Li B, He J, et al. Changes in the nutrients, phytochemical profile and antioxidant activity of *Rheum officinale* Baill. leaf blades during different growth periods. *Front Nutr.* 2024;11:1387947. <https://doi.org/10.3389/fnut.2024.1387947>
19. Rajamanickam G, Manju SL. Identification and comparative study of *invitro* antioxidant potential of fractionated hydroalcoholic extract of *Phyllanthus Niruri* Linn. *Eur J Adv Chem Res.* 2020;1(1):1-8. <https://doi.org/10.24018/ejchem.2020.1.1.2>

20. Araújo OF, Kristiani EBE, Kasmiyati S, Nugroho LH. The Phytochemical and biological activities of two phyllanthus species: Insights into metabolit, antioxidant and antibacterial activity. *Biosaintifika J Biol Biol Educ.* 2024;16(3):403–14. <https://doi.org/10.15294/biosaintifika.v16i3.13944>
21. Swargiary D, Kashyap B, Sarma P, Ahmed SA, Gurumayum S, Barge SR, et al. Free radical scavenging polyphenols isolated from *Phyllanthus niruri* L. ameliorates hyperglycemia via SIRT1 induction and GLUT4 translocation in *in vitro* and *in vivo* models. *Fitoterapia.* 2024;173:105803. <https://doi.org/10.1016/j.fitote.2023.105803>
22. Sihag S, Pal A, Ravikant, Saharan V. Antioxidant properties and free radicals scavenging activities of pomegranate (*Punica granatum* L.) peels: An *in-vitro* study. *Biocatal Agric Biotechnol.* 2022;42:102368. <https://doi.org/10.1016/j.bcab.2022.102368>
23. Zhao X, Yuan Z. Anthocyanins from Pomegranate (*Punica granatum* L.) and their role in antioxidant capacities *In vitro*. *Chem Biodivers.* 2021;18(10):e2100399. <https://doi.org/10.1002/cbdv.202100399>
24. Siddique S, Parveen Z, Firdaus-e-Bareen, Mazhar S. Chemical composition, antibacterial and antioxidant activities of essential oils from leaves of three Melaleuca species of Pakistani flora. *Arab J Chem.* 2020;13(1):67-74. <https://doi.org/10.1016/j.arabjc.2017.01.018>
25. Rahimi A, Mohammadi MM, Siavash Moghaddam S, Heydarzadeh S, Gitari H. Effects of stress modifier biostimulants on vegetative growth, nutrients and antioxidants contents of garden thyme (*Thymus vulgaris* L.) under water deficit conditions. *J Plant Growth Regul.* 2022;41(5):2059–72. <https://doi.org/10.1007/s00344-022-10604-6>
26. Prasathkumar M, Anisha S, Khusro A, Mohamed Essa M, Babu Chidambaram S, Walid Qoronfleh M, et al. Anti-pathogenic, anti-diabetic, anti-inflammatory, antioxidant and wound healing efficacy of *Datura metel* L. leaves. *Arab J Chem.* 2022;15(9):104112. <https://doi.org/10.1016/j.arabjc.2022>
27. Saha A, Basak BB, Banerjee A. *In vitro* antioxidant evaluation and production of biochar from distillation waste biomass of *Mentha arvensis*. *J Appl Res Med Aromat Plants.* 2022;31:100428. <https://doi.org/10.1016/j.jarmap.2022.100428>
28. Widyastuti I, Luthfah HZ, Hartono YI, Islamadina R, Can AT, Rohman A. Antioxidant activity of temulawak (*Curcuma xanthorrhiza* Roxb.) and its classification with chemometrics. *Indonesian J Chemom Pharm Anal.* 2020;1(1):28-41. <https://doi.org/10.22146/ijcpa.507>
29. Asyhar R, Minarni M, Arista RA, Nurcholis W. Total phenolic and flavonoid contents and their antioxidant capacity of *Curcuma xanthorrhiza* accessions from Jambi. *Biodiversitas J Biol Divers.* 2023;24(9). <https://doi.org/10.13057/biodiv/d240944>
30. Kusumadewi AP, Martien R, Pramono S, Setyawan AA, Windarsih A, Rohman A. Application of FTIR spectroscopy and chemometrics for correlation of antioxidant activities, phenolics and flavonoid contents of Indonesian *Curcuma xanthorrhiza*. *Int J Food Prop.* 2022;25(1):2364–72. <https://doi.org/10.1080/10942912.2022.2134418>
31. Petrosyan HR, Nigaryan AA, Hovhannisyan HA, Soloyan AM, Vardapetyan VV, Martiryan AI. Evaluation of antioxidant activity and heavy metals content in licorice (*Glycyrrhiza glabra* L.) growing wild in Armenia. *Heliyon* 2023;9(11):e22442. <https://doi.org/10.1016/j.heliyon.2023.e22442>
32. Mutaillifu P, Bobakulov K, Abuduwalli A, Huojiaihemaiti H, Nuerxiati R, Aisa HA, et al. Structural characterization and antioxidant activities of a water soluble polysaccharide isolated from *Glycyrrhiza glabra*. *Int J Biol Macromol.* 2020;144:751–59. <https://doi.org/10.1016/j.ijbiomac.2019.11.245>
33. Semenescu I, Avram S, Similie D, Minda D, Diaconeasa Z, Muntean D, et al. Phytochemical, antioxidant, antimicrobial and safety profile of *Glycyrrhiza glabra* L. extract obtained from Romania. *Plants.* 2024;13(23):3265. <https://doi.org/10.3390/plants13233265>
34. Satake T, Kamiya K, Ohno A, Horii Y, Endang H, Umar M. A-Type Proanthocyanidins from the bark of *Parameria laevigata*. *Heterocycles.* 2003;60(7):1697. <https://doi.org/10.3987/COM-03-9793>
35. Mustafa I, Chin NL. Antioxidant properties of dried ginger (*Zingiber officinale* Roscoe) var. Bentong. *Foods.* 2023;12(1):178. <https://doi.org/10.3390/foods12010178>
36. Alfuraydi AA, Aziz IM, Almajhdi FN. Assessment of antioxidant, anticancer and antibacterial activities of the rhizome of ginger (*Zingiber officinale*). *J King Saud Univ - Sci.* 2024;36(3):103112. <https://doi.org/10.1016/j.jksus.2024.103112>
37. Kandasamy A, Aruchamy K, Rangasamy P, Varadhaiyan D, Gowri C, Oh TH, et al. Phytochemical analysis and antioxidant activity of centella asiatica extracts: an experimental and theoretical investigation of flavonoids. *Plants.* 2023;12(20):3547. <https://doi.org/10.3390/plants12203547>
38. Sabaragamuwa R, Perera CO. Total triterpenes, polyphenols, flavonoids and antioxidant activity of bioactive phytochemicals of *Centella asiatica* by different extraction techniques. *Foods.* 2023;12(21):3972. <https://doi.org/10.3390/foods12213972>
39. Moradi-Ozarlou M, Javanmardi S, Tayefi-Nasrabadi H. Antioxidant property of *Plantago major* leaf extracts reduces testicular torsion/detorsion-induced ischemia/reperfusion injury in rats. *Vet Res forum an Int Q J.* 2020;11(1):27-33. <https://doi.org/10.30466/vrf.2019.102182.2432>
40. Şahin B, Savcı A. Bioactive component analysis and *in vitro* antioxidant activities of *Plantago major* L. *Türk Doğa ve Fen Derg.* 2024;13(4):129-34. <https://doi.org/10.46810/tdfd.1529274>
41. Araujo L, Nunes K, Mello J, Nakamura C, Gomes R, Bergamasco R. Evaluation of the antioxidant, photoprotective and wound healing capacity of *Guazuma ulmifolia* lam. extracts in L-929 cells exposed to UV-A and UV-B irradiation. *J Braz Chem Soc.* 2024;35(3):e-20230149. <https://doi.org/10.21577/0103-5053.20230149>
42. Marda M, Naid T, Rahmawati R. Antioxidant activity of dechlorophyllation of kejobeling leaf (*Strobilanthes crispus* L.) extract using the free radical reduction method. *Pharm Reports.* 2024;3(1):6-11. <https://doi.org/10.33096/pharmrep.v3i1.285>
43. Deng L, Huang G. Ultrasound-assisted extraction, optimization, characteristics and antioxidant activity of *Piper nigrum* L. polysaccharides. *Ultrason Sonochem.* 2025;116:107309. <https://doi.org/10.1016/j.ultsonch.2025.107309>
44. Mollik M, Rahman MH, Al-Shaeri M, Ashraf GM, Alexiou A, Gafur MA. Isolation, characterization and *in vitro* antioxidant activity screening of pure compound from black pepper (*Piper nigrum*). *Environ Sci Pollut Res.* 2022;29(34):52220–32. <https://doi.org/10.1007/s11356-022-19403-8>
45. Khiya Z, Oualcadi Y, Gamar A, Berrekhis F, Zair T, Hilali F EL. Azaroual M, editor. Correlation of total polyphenolic content with antioxidant activity of hydromethanolic extract and their fractions of the *Salvia officinalis* leaves from different regions of Morocco. *J Chem.* 2021:1–11. <https://doi.org/10.1155/2021/8585313>
46. Bibi N, Shah MH, Khan N, Al-Hashimi A, Elshikh MS, Iqbal A, et al. Variations in total phenolic, total flavonoid contents and free radicals' scavenging potential of onion varieties planted under diverse environmental conditions. *Plants.* 2022;11(7):950. <https://doi.org/10.3390/plants11070950>
47. Muflihah YM, Gollavelli G, Ling Y-C. Correlation study of antioxidant activity with phenolic and flavonoid compounds in 12 Indonesian indigenous herbs. *Antioxidants.* 2021;10(10):1530. <https://doi.org/10.3390/antiox10101530>
48. Rumpf J, Burger R, Schulze M. Statistical evaluation of DPPH, ABTS, FRAP and Folin-Ciocalteu assays to assess the antioxidant capacity of lignins. *Int J Biol Macromol.* 2023;233:123470. <https://doi.org/10.1016/j.ijbiomac.2023.123470>
49. Kourouma V, Mu T, Zhang M, Sun H. Comparative study on chemical composition, polyphenols, flavonoids, carotenoids and antioxidant

activities of various cultivars of sweet potato. Int J Food Sci Technol. 2020;55(1):369–78. <https://doi.org/10.1111/ijfs.14336>

Additional information

Peer review: Publisher thanks Sectional Editor and the other anonymous reviewers for their contribution to the peer review of this work.

Reprints & permissions information is available at https://horizonpublishing.com/journals/index.php/PST/open_access_policy

Publisher's Note: Horizon e-Publishing Group remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Indexing: Plant Science Today, published by Horizon e-Publishing Group, is covered by Scopus, Web of Science, BIOSIS Previews, Clarivate Analytics, NAAS, UGC Care, etc
See https://horizonpublishing.com/journals/index.php/PST/indexing_abstracting

Copyright: © The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits

unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited (<https://creativecommons.org/licenses/by/4.0/>)

Publisher information: Plant Science Today is published by HORIZON e-Publishing Group with support from Empirion Publishers Private Limited, Thiruvananthapuram, India.