



RESEARCH ARTICLE

# Balanced fertilization for yield maximization, profitability and quality of Indian mustard (*Brassica juncea* L.) in North Western Himalayas

Rajeev Bharat<sup>1\*</sup>, Sarabdeep Kour<sup>2</sup>, Meenakshi Gupta<sup>3</sup>, Vivek Bhagat<sup>3</sup>, Divya Chadha<sup>2</sup>, Bupesh Kumar<sup>1</sup>, Sunil Kumar Rai<sup>1</sup> & Narinder Panotra<sup>4</sup>

<sup>1</sup>Division of Plant Breeding & Genetics, Sher-e-Kashmir University of Agricultural Sciences & Technology of Jammu, Chatha 180 009, India

<sup>2</sup>Division of Soil Science and Agricultural Chemistry, Sher-e-Kashmir University of Agricultural Sciences & Technology of Jammu, Chatha 180 009, India

<sup>3</sup>Division of Agronomy, Sher-e-Kashmir University of Agricultural Sciences & Technology of Jammu, Chatha 180 009, India

<sup>4</sup>School of Biotechnology, Sher-e-Kashmir University of Agricultural Sciences & Technology of Jammu, Main Campus, Chatha 180 009, India

\*Correspondence email - [sangra.rajeev@gmail.com](mailto:sangra.rajeev@gmail.com)

Received: 23 February 2025; Accepted: 27 May 2025; Available online: Version 1.0: 16 July 2025

**Cite this article:** Rajeev B, Sarabdeep K, Meenakshi G, Vivek B, Divya C, Bupesh K, Sunil KR, Narinder P. Balanced fertilization for yield maximization, profitability and quality of Indian mustard (*Brassica juncea* L.) in North Western Himalayas. Plant Science Today (Early Access).

<https://doi.org/10.14719/pst.7893>

## Abstract

Indian mustard is one of the major oilseed crops of the country. The productivity of this crop is quite low compared to the rapeseed mustard crops in some other countries, such as Canada and Germany. One of the major reasons for lower productivity is the lack of balanced nutrition in many parts of the country. Sulphur (S), boron (B) and zinc (Zn) play an important role in the metabolism of Rapeseed mustard crops. A field experiment was conducted at Research Farm, SKUAST-Jammu in 2022 and 2023, to investigate the influence of different fertility levels with macro and micro nutrients on productivity, profitability and quality of Indian mustard. The treatment consisted of four fertility levels containing N:P<sub>2</sub>O<sub>5</sub>:K<sub>2</sub>O doses viz., F<sub>1</sub>-control, F<sub>2</sub>- 80:40:20 kg ha<sup>-1</sup>, F<sub>3</sub>- 100:50:25 kg ha<sup>-1</sup> and F<sub>4</sub>- 120:60:30 kg ha<sup>-1</sup> respectively in main plots and four macro and micronutrient treatments namely, N<sub>1</sub>- 20 kg S+2.5 kg Zn + 0.5 kg B, N<sub>2</sub>- 40 kg S+ 5.0 kg Zn +1.0 kg B, N<sub>3</sub>- N<sub>1</sub> + 500 kg FYM ha<sup>-1</sup>, N<sub>4</sub>- N<sub>2</sub> + 500 kg FYM ha<sup>-1</sup> in sub plots, which were laid out in split-plot design and replicated thrice. The results revealed that the application of N:P<sub>2</sub>O<sub>5</sub>:K<sub>2</sub>O at 120:60:30 kg ha<sup>-1</sup> in combination with 40 kg S+ 5.0 kg Zn +1.0 kg B enriched with 500 kg FYM ha<sup>-1</sup> recorded a significant increase in seed and oil yield along with yield attributes, nutrient uptake, soil microbial population, net returns and B:C ratio of Indian mustard than other treatments in comparison. A significant positive correlation was noted between seed yield and nutrient uptake.

**Keywords:** boron; farmyard manure; nutrient uptake; quality; rapeseed mustard; sulphur; yield

## Introduction

In India, oilseeds are the second utmost vital crop after cereals. They are major sources of crucial fatty acids and vitamin E. After USA, China and Argentina, India is the fourth largest oilseeds producer with 10 % contribution to total global production. Rapeseed mustard crops are cultivated on an area of 41.95 million hectares worldwide, with the total production and productivity of 91.0 million tonnes and 21.10 tonnes per hectare respectively (1). In 2024, India imports 15 % of the world's edible oil and becomes the largest global purchaser of edible oils, with \$16.5 billion in imports (2). In India, amongst all oilseed crops, rapeseed-mustard is one of the most important oilseed crops with maximum acreage in the country (3). The total oilseed production in the country is 39.59 million tonnes, out of which rapeseed mustard crops are the single largest contributor with 33.24 % share in the total oilseed production. However, the productivity in the country is quite low when

compared to the leading rapeseed mustard producing countries, such as China, Germany, Canada besides other European countries where the per hectare productivity is more than 20 quintals per hectare average yield in comparison to 15 quintal per hectare average yield realized in the country so far.

In North western Himalayan region, the use of old and obsolete low yielding varieties, agronomic practices, water stress, delayed sowing, climate variability, early crop stage or terminal heat stress besides the imbalanced fertilization and non-application of secondary macronutrient like sulphur (S) and other micronutrients like boron (B) and zinc (Zn) on regular basis in the fields are the major causes of lower productivity of Indian mustard. The exclusive application of chemical fertilizers like Urea, Diammonium phosphate (DAP) and Muriate of Potash (MOP) results in depletion of soil quality and drastic reduction in crop yields owing to decreased factor productivity. Mustard crop is highly sensitive to the deficiency of sulphur,


zinc and boron which is common in mustard growing areas of North western region and is the major constraint in achieving higher productivity. Thus, potential productivity of Indian mustard can only be achieved by balanced use of nutrients including secondary macro and micro-nutrients. Further, integration of different nutrients like nitrogen, phosphorus and potassium with macro and micro nutrients and farm yard manure (FYM) is pivotal for increasing the productivity in the country as it has a beneficial effect on the growth, yield and quality of Indian mustard. The enrichment of synthetic fertilizers with FYM have shown promising results in sustaining productivity. This act of bio-energizing helps in enhancing the activities of microbial population and soil physico-chemical properties through the increasing the mineralization process through direct release of macro and micro nutrients which resulted in increased crop yield (4). Also, sulphur is considered as the most important nutrient after nitrogen, phosphorus and potassium (5). It plays an indispensable role in formation of three important amino acids *viz.*, cysteine, cystine and methionine which are essential for protein synthesis and various metabolic activities in plant (6). Sulphur also participates in the activities of sulphhydryl (SH-) linkages that are responsible for formation of Sinigrin, which is the root cause of pungency in oilseeds (7). Similarly, zinc is the first micronutrient to be recognized as crucial for plant. It plays a crucial role in stability of cytoplasmic ribosomes, cell division, dehydrogenase, proteinase and peptide enzymes and helps in the formation of protein and carotene (8). Better zinc nutrition of crop encouraged both primary and secondary branches, which results in higher seed and stover yield (9). Likewise, boron is the essential micronutrient which is needed by the crop for normal growth. It maintains proper water equilibrium within the plant and facilitates the fruiting processes. It is involved in the synthesis of both oil and protein. The other important functions of boron in plants are formation and

stability of cell wall, maintenance of structural and functional integrity of the biological membranes and transportation of sugar products from source to sink (10). All these factors have necessitated research to determine FYM enriched, sulphur, zinc and boron levels in conjunction with different fertilizer levels for the supply of vital micro and macronutrients in order to provide meaningful recommendations to farmers in the region as it is expected to help in increasing the yield of Indian mustard in Jammu region significantly in the years to come in view of increasing area under cultivation of mustard crop under Jammu conditions.

To ensure sustainable production of Indian mustard, it is vital to optimize nutrient management in this crop. In North western region, many agricultural soils face the problem of nutrient imbalance or deficiency specifically sulphur, zinc and boron. Therefore, the present investigation aimed to rectify this problem and provides the region-specific fertilizer recommendation which helps to empowering the farmer community to make informed decisions for their crop. This research also supports food security and contributes to the economic well being of farming communities in the North Western region.

## Materials and Methods

The experiment was performed at the Oilseed Experimental Area, Research Farm, Sher-e-Kashmir University of Agricultural Sciences and Technology, Jammu (Fig. 1), during the *rabi* season of 2022 and 2023 to study the effect of different fertilizer levels with macro and micronutrients on growth, productivity and quality of Indian mustard. Split Plot Design (SPD) was used with three replications. The net plot size was 5 m × 4.5 m. The crop was sown in lines 30 cm apart (row-to-row). The chemical fertilizers were applied as per treatments *viz.*, 80:40:20, 100:50:25 and 120:60:30 kg NPK ha<sup>-1</sup>



**Fig. 1.** Location map of the experimental site at Research Farm, SKUAST-Jammu (indicated as red location sign).

for 100, 125 and 150 % fertility levels, respectively. The specific levels of nitrogen (N), phosphorus (P), potassium (K) and sulphur (S) were chosen based on their initial levels of these nutrients in experimental field. The soil in the experimental field was low in organic carbon, available nitrogen and zinc but medium in available phosphorous, potassium, sulphur and boron, respectively. Therefore, the recommended dose of fertilizers were increased up to 150 % to examine whether the higher nutrients levels would contribute to an increase in the seed yield of mustard crop. Also, the mustard crop variety RH 725 used in the present study has a higher yield potential and require a higher dose of nutrient to achieve potential yield (11). Further, the levels of available zinc (Zn) and boron (B), their selection was based on low availability of these micronutrients in the experimental field, as well as some previous studies in the region that utilized almost similar levels (12, 13). The nutrient sources were urea, diammonium phosphate, muriate of potash, sulphur bentonite -90 % S, zinc sulphate monohydrate and borax, respectively. The required secondary macro and micronutrients as per the specified treatments namely sulphur, zinc and boron were enriched using farm yard manure at 500 kg ha<sup>-1</sup> by mixing the fertilizers for 15 days in advance at the site for bioactivation. Frequent stirring of the mixture after every 3 days period was ensured for uniform mixing and microbial activation. The experimental treatments were as follow:

### Treatments

#### Main plot treatments (Fertility levels)

F<sub>1</sub>: Control

F<sub>2</sub>: N:P<sub>2</sub>O<sub>5</sub>:K<sub>2</sub>O at 80:40:20 kg ha<sup>-1</sup>

F<sub>3</sub>: N:P<sub>2</sub>O<sub>5</sub>:K<sub>2</sub>O at 100:50:25 kg ha<sup>-1</sup>

F<sub>4</sub>: N:P<sub>2</sub>O<sub>5</sub>:K<sub>2</sub>O at 120:60:30 kg ha<sup>-1</sup>

#### Sub plot treatments (Macro and micro nutrients)

N<sub>1</sub>: 20 kg S+2.5 kg Zinc+0.5 kg Boron

N<sub>2</sub>: 40 kg S+ 5.0 kg Zinc+1.0 kg Boron

N<sub>3</sub>: 20 kg S+2.5 kg Zinc+0.5 kg Boron enriched with 500 kg FYM ha<sup>-1</sup>

N<sub>4</sub>: 40 kg S+ 5.0 kg Zinc+1.0 kg Boron enriched with 500 kg FYM ha<sup>-1</sup>

### Soil analysis

The soil analysis was done before applying the experimental treatments and after harvesting of mustard crop. Soil samples from each plot were randomly collected from a depth of 0-15 cm at five different spots within the plot. These samples were combined to form a composite sample for the experimental area, which was then air dried, ground into a fine powder and passed through a 2 mm sieve. The physico-chemical analysis of the experimental area is presented in Table 1.

**Table 1.** Physico-chemical properties of soil

Parameter	Sampling depth	Units	Values	
			Pre-sowing	
			2022-23	2023-24
pH			7.48	7.41
EC		dS m <sup>-1</sup>	0.26	0.25
Organic carbon		g kg <sup>-1</sup>	4.09	4.11
Nitrogen		kg ha <sup>-1</sup>	147.03	147.36
Phosphorus		kg ha <sup>-1</sup>	15.10	15.11
Potassium		kg ha <sup>-1</sup>	118.04	119.56
Sulphur		kg ha <sup>-1</sup>	28.01	27.98
Zinc		mg kg <sup>-1</sup>	0.56	0.57
Boron		mg kg <sup>-1</sup>	0.52	0.53
Soil Texture			Sandy clay loam	

### Meteorological data

Weather data was acquired from the meteorological observatory located close to the experimental site at Research Farm, Sher-e-Kashmir University of Agricultural Sciences & Technology of Jammu, Jammu & Kashmir, India. The total amount of rainfall observed was 44.5 and 31.3 mm coinciding with the critical crop growth stages viz. flowering and siliquae formation and seed filling stage during both the crop growth period, respectively. Generally, lower temperature and higher relative humidity and rainfall were recorded in 2023 than 2022. The climatic conditions that existed during the growing period of crop as shown in Table 2.

### Crop husbandry

The mustard variety RH-725 was sown manually on 22 October 2022 and 20 October 2023 using seed rate of 5 kg ha<sup>-1</sup>. The crop was planted in rows spaced 30 cm apart. Only one irrigation was given to the crop at the initial stage in both years as the water received from rainfall at different growth stages was sufficient for the crop during both years of experimentation. The crop was harvested on 23 March 2023 and 22 March 2024, respectively, when 80 % of siliqua color turned yellowish. Harvesting was done manually using sickles. Immediately after harvesting of the crop, bundles were tied and tagged bundles were kept in plots for sun drying. After proper sun drying, threshing was done with the help to sticks and seed yield was noted plot wise using electronic balance.

### Observations recorded and procedure for recording observations

**Number of siliqua plant<sup>-1</sup>:** Siliquae were counted from five tagged plants and their mean value represented the Siliqua plant<sup>-1</sup>.

**Seeds siliqua<sup>-1</sup>:** Similarly, it was calculated by mean counts from five siliquae of each tagged plant.

**1000-seed weight:** A random sample of 1000 seeds from each net plot was weighed using a digital balance and noted in grams.

**Table 2.** Meteorological data during the crop season

Month	2022-23					2023-24				
	Temperature (°C)		Relative Humidity (%)		Rainfall (mm)	Temperature (°C)		Relative Humidity (%)		Rainfall (mm)
	Max	Min	8 am	5 pm		Max	Min	8 am	5 pm	
October	30.3	13.65	85.9	43.15	0.0	29.8	14.3	89.7	48.5	0.0
November	25.05	9.3	91.12	48.45	6.2	25.6	10.4	91.4	52.2	18.2
December	20.92	5.55	92.95	51.9	2.4	21.2	5.7	95.9	53.7	0.8
January	15.7	4.8	95.1	71.8	4.9	12.5	4.7	95.8	77.1	0.0
February	23.6	7.7	89.5	53.0	8.2	20.7	5.6	92.9	52.7	10.5
March	27.0	12.3	90.1	50.3	9.6	25.1	9.3	89.3	45.0	15.0

**Seed yield:** The seed were threshed and cleaned from each net plot area. The value was attained in kg plot<sup>-1</sup> then converted in kg ha<sup>-1</sup> by given formula

Seed yield (kg ha<sup>-1</sup>) =

$$\frac{\text{Yield obtained from net plot (kg)} \times 10000}{\text{Area of net plot}}$$

**Oil content:** First, seed samples were oven dried at 70 °C to eliminate moisture then crushed in a pestle-mortar and analyzed for oil content using the Soxhlet apparatus (14).

Oil content =

$$\frac{\text{Weight of flask + petroleum ether extract} - \text{Weight of flask with oil}}{\text{Weight of sample (g)}} \times 100$$

**Oil yield:** It was calculated by multiplying oil content with seed yield of mustard using the formula:

Oil yield (kg ha<sup>-1</sup>) =

$$\frac{\text{Oil content (\%)} \times \text{Seed yield (kg ha}^{-1})}{100}$$

**Nutrient uptake:** Harvested samples were collected for nutrient analysis of N, P, K, S, Zn and B in seed and stover. After oven-drying, the plant samples were grinded and analyzed for N, P, K, S, Zn and B concentration. Nutrient uptake was calculated by multiplying nutrient concentration and dry matter accumulation (seed and stover).

Nutrient uptake (kg ha<sup>-1</sup>) =

$$\frac{\text{Nutrient content (\%)} \times \text{dry matter accumulation (kg ha}^{-1})}{100}$$

#### Available nutrients in soil

**Nitrogen:** It was determined using alkaline permanganate method (15). Processed soil is added to distillation flask along with 0.32 % KMnO<sub>4</sub> and 2.5 % NaOH. The resulting ammonia is captured in boric acid, which contains a mixed indicator of methyl red and bromocresol green adjusted to pH of 8.5. After that the collected solution was titrated with standard acid. It was expressed in kg ha<sup>-1</sup>.

**Phosphorus:** Determined using Olsen's method with spectrophotometer at 660 nm and expressed in kg ha<sup>-1</sup> (16).

**Potassium:** It was measured using neutral normal ammonium acetate solution and determined by flame photometer and expressed in kg ha<sup>-1</sup> (17).

**Sulphur:** Assessed using calcium chloride (0.15 %) (18). It was determined by spectrophotometer at 420 nm and expressed in kg ha<sup>-1</sup>.

**Zinc:** Determined using DTPA extract. It was analyzed by atomic absorption spectrophotometer and represented as mg kg<sup>-1</sup>.

**Boron:** Quantified by the colorimetric Azomethine-H method (19). It was expressed as mg kg<sup>-1</sup>.

**Microbial population:** Soil microbial population viz., bacteria, fungi and actinomycetes were determined using the serial dilution agar plating method. The diluted samples were

plated on respective media and incubated at 25±2 °C. For bacterial growth, incubation was carried out for 24 hr on Nutrient Agar Medium; for fungal growth, 7 days on Potato Dextrose Agar (PDA) whereas for actinomycetes it was kept for 7 days on Yeast Malt Agar (YMA). Colonies were counted, averaged and multiplied by the dilution factor to estimate microbial density per gram of soil (20).

Number of cells mL<sup>-1</sup> or g<sup>-1</sup> = Number of colonies (average of 3 replications) × Dilution factor

Dilution factor = Reciprocal of the dilution (e.g. 10<sup>9</sup> = 10)

#### Economics

**Net returns:** It was determined by subtracting total cost of cultivation from gross return and expressed as ₹ ha<sup>-1</sup>.

**B: C ratio:** The benefit-cost ratio was calculated by dividing net returns by total cost of cultivation to identify the most cost-effective treatment.

#### Statistical analysis

The recorded data of both years were statistically evaluated using a two-way analysis of variance (ANOVA). Fisher's test of significance was used to compare significant treatment means at 5 % probability level. All statistical analyses were performed using OP STAT software and graphs were made by using MS Excel and R software version 4.2.2.

## Results and Discussion

#### Yield attributes

Siliqua plant<sup>-1</sup> was significantly affected by fertility levels and macro and micro nutrients during 2022 and 2023 cropping seasons. During 2022, the highest siliqua plant<sup>-1</sup>, seeds siliqua<sup>-1</sup> and 1000-seed weight was recorded in F<sub>4</sub> (348; 18.58; 4.62), followed by F<sub>3</sub> (328.33; 17.23; 4.38) and F<sub>2</sub> (245.50; 16.17; 3.97), respectively (Table 3). Whereas the lowest value was recorded under the control (F<sub>1</sub>) treatment. Similarly, the maximum siliqua plant<sup>-1</sup>, seeds siliqua<sup>-1</sup> and 1000-seed weight was recorded under treatment F<sub>4</sub> (317.38; 17.80; 5.05), closely followed by F<sub>3</sub> (309.75; 17.65; 4.85) and F<sub>2</sub> (281.22; 15.37; 4.59), respectively and the least under the control (F<sub>1</sub>) treatment during 2023. Higher nutrient availability was the major cause of improvement in yield attributes. Among the macro and micro nutrients, N<sub>4</sub> achieved significantly highest siliqua plant<sup>-1</sup>, seeds siliqua<sup>-1</sup> and 1000-seed weight, (285; 16.80; 4.14), respectively, than the rest of the treatments. Though the control was the least performing treatment in the year of 2022. During the second growing season, the maximum number of siliqua plant<sup>-1</sup>, seeds siliqua<sup>-1</sup> and 1000-seed weight were observed under the treatment N<sub>4</sub> (298.68; 16.57; 4.83) followed by N<sub>3</sub> (289.22; 15.98; 4.63), N<sub>2</sub> (267.92; 14.50; 4.43) and N<sub>1</sub> (258.93; 14.15; 4.32), respectively. Better availability of sulphur directly increased chlorophyll production, protein synthesis, carbohydrate metabolism and the translocation of photosynthates. The combined application of B and Zn also have beneficial effects on the reproductive parts of mustard (21-24). The interaction effect between the fertility levels and macro and micro nutrients in yield attributes was non-significant.

**Table 3.** Response of fertility levels and integrated nutrient management on yield attributes and yield of Indian mustard

Treatments	2022-23				2023-24			
	Number of siliqua plant <sup>-1</sup>	Seeds siliqua <sup>-1</sup>	1000-seed weight (g)	Seed yield (kg ha <sup>-1</sup> )	Number of siliqua plant <sup>-1</sup>	Seeds siliqua <sup>-1</sup>	1000-seed weight (g)	Seed yield (kg ha <sup>-1</sup> )
<b>Fertility Levels</b>								
F <sub>1</sub>	184.25	12.73	2.76	980	206.40	10.38	3.71	1011
F <sub>2</sub>	245.50	16.17	3.97	1466	281.22	15.37	4.59	1488
F <sub>3</sub>	328.33	17.23	4.38	1608	309.75	17.65	4.85	1675
F <sub>4</sub>	348.25	18.58	4.62	1698	317.38	17.80	5.05	1699
SEm(±)	6.75	0.43	0.10	28.23	5.84	0.37	0.10	34.22
LSD (0.05)	23	1.49	0.35	97	20.63	1.32	0.35	118
<b>Macro and micro nutrients</b>								
N <sub>1</sub>	262.91	15.37	3.65	1242	258.93	14.15	4.32	1362
N <sub>2</sub>	277.08	16.00	3.87	1325	267.92	14.50	4.43	1408
N <sub>3</sub>	282.00	16.55	4.06	1569	289.22	15.98	4.63	1525
N <sub>4</sub>	285.33	16.80	4.14	1617	298.68	16.57	4.83	1578
SEm(±)	3.06	0.14	0.03	32.97	3.86	0.22	0.09	18.62
LSD (0.05)	9	0.41	0.12	96	11.36	0.67	0.27	54
F × N	NS	NS	NS	NS	NS	NS	NS	NS

### Seed yield

The different fertility levels and macro and micro nutrients have an essential role in increasing the seed yield of Indian mustard. Consequently, a higher seed yield (1698) was documented during the cropping season of 2022 in F<sub>4</sub>, followed by F<sub>3</sub> (1608) and F<sub>2</sub> (1466) (Table 3), while a noticeable decrease in seed yield (980) was recorded in F<sub>1</sub> treatment as compared to other three treatments. Similarly, for the 2023 growing season, maximum seed yield was achieved in F<sub>4</sub> (1699) than in all other three treatments. Amongst macro and micro nutrients highest seed yield (1617) in 2022 was statistically achieved by treatment N<sub>4</sub> and treatment N<sub>1</sub> provided the lowest seed yield (1242) as compared to other treatments during 2022. In the growing season of 2023, maximum seed yield was observed under the treatment N<sub>4</sub> (1578), followed by N<sub>3</sub> (1525), N<sub>2</sub> (1408) and N<sub>1</sub> (1362). The significant increase in seed yield of Indian mustard was due to better growth and yield attributes through higher supply of major nutrients, which helped to compensate for the restricted nutrient availability. Also, micronutrients enriched with FYM may have affected the plant growth and development of Indian mustard during both the crop growing seasons. Moreover, the favourable rainfall and temperature during the flowering and seed formation stage in second crop growing season played a key role in higher crop growth and yield (25-28). Interaction between fertility levels and macro and micro nutrients in seed yield was non-significant.

### Oil content

During 2022, oil content in treatment F<sub>4</sub> was increased to 38.67, whereas statistically minimum oil content was observed under the control treatment, in contrary to all other three treatments (Table 6). Consequently, maximum oil content was prominently obtained in treatment F<sub>4</sub> (38.79), followed by F<sub>3</sub> (38.19) and F<sub>2</sub> (37.46), while least oil content was recorded in F<sub>1</sub> (34.02) treatment during 2023. Application of NPK fertilizers may have activated the enzymes responsible for producing higher oil content in Indian mustard. A major enhancement in oil content (38.63) was attained under N<sub>4</sub> treatment as compared to rest of the treatments in the growing season of 2022. During 2023, higher oil content was obtained in treatment N<sub>4</sub> (38.74), followed by N<sub>3</sub> (38.14), N<sub>2</sub> (36.20) and N<sub>1</sub> (35.38). Sulphur has a significant role in increasing oil content and is vital for synthesis of fatty acid and oil synthesis. Similarly, zinc may have a role in


enzymatic activation, which aids in more oil synthesis in the crop (16, 29). However, interaction between both the factors was non-significant.

### Oil yield

Significantly maximum oil yield was noticed in treatment F<sub>4</sub> (657.56), followed by F<sub>3</sub> (615.89) and F<sub>2</sub> (551.66), while minimum oil yield (333.40) was attained from the control (F<sub>1</sub>) treatment during the growing season of 2022. During 2023, again highest oil yield was recorded under F<sub>4</sub> (660.02) treatment. However, a significant decline in oil yield was noticed in F<sub>1</sub> (345.89) (Fig. 2) as compared to rest of the treatments. Among the macro and micro nutrients, the highest oil yield was attained in N<sub>4</sub> (630.27) in contrary to all other three treatments during 2022. The maximum oil yield (617.07) was statistically obtained in N<sub>4</sub>, followed by N<sub>3</sub> (586.77), N<sub>2</sub> (515.45) and N<sub>1</sub> (488.14) during 2023. Oil yield in crops is influenced not just by seed yield but also by the biochemical processes governing oil synthesis, which are highly sensitive to nutrient balance and environmental stress. An adequate and balanced supply of macronutrients creates optimal conditions for seed formation. However, micronutrients also played a critical role in enhancing oil yield. Sulfur is directly involved in the synthesis of certain amino acids and enzymes essential for oil formation. Zinc and boron influence hormonal balance and cell division, which are vital during the flowering and seed development stages. Furthermore, boron facilitates the translocation of sugars and oils to the developing seeds. The enrichment of these micronutrients with FYM ensured a slow and steady release of nutrients, helping the plant maintain physiological functions. This not only supported higher seed formation but also improved oil content in those seeds, thus increasing total oil yield (30-32).

### Nutrient uptake

The fertility levels, as well as macro and micro nutrients, have a major effect on nutrient uptake (N, P, K, S, Zn and B) of Indian mustard and bio-energizing farmyard manure with sulphur, zinc and boron at variable rates, which significantly increased the uptake of nutrients in the crop. However, maximum macronutrient uptake in respect of N, P, K and S (86.44; 19.01; 96.44; 20.46 kg ha<sup>-1</sup>) and micronutrient uptake in respect of Zn and B (410.90; 173.52 g ha<sup>-1</sup>) was recorded in treatment F<sub>4</sub> (Table 4) during 2022. A significant decrease in



**Fig. 2.** Response of fertility levels and integrated nutrient management on oil yield and grain yield of Indian mustard.

**Table 4.** Response of fertility levels and integrated nutrient management on nutrient uptake of Indian mustard

Treatments	2022-23						2023-24					
	N	P	K	S	Zn	B	N	P	K	S	Zn	B
<b>Fertility Levels</b>												
F <sub>1</sub>	39.72	8.74	49.72	12.43	259.17	113.23	43.47	9.30	51.22	13.12	261.97	114.70
F <sub>2</sub>	79.02	17.38	89.02	18.89	387.75	161.71	84.02	17.98	90.52	19.69	390.55	163.31
F <sub>3</sub>	81.39	17.90	91.39	19.39	395.14	165.48	86.39	18.50	92.89	20.19	397.94	167.08
F <sub>4</sub>	86.44	19.01	96.44	20.46	410.90	173.52	91.44	19.78	98.10	21.38	413.95	175.28
SEm(±)	1.73	0.38	1.73	0.37	5.39	2.84	1.80	0.39	1.74	0.43	5.41	2.84
LSD (0.05)	6.00	1.31	6.00	1.30	18.62	9.80	6.23	1.37	6.03	1.28	18.67	9.82
<b>Macro and micro nutrients</b>												
N <sub>1</sub>	67.90	14.93	77.90	16.98	351.63	147.35	71.65	15.50	79.40	17.66	354.43	148.81
N <sub>2</sub>	70.02	15.40	80.02	17.44	358.23	150.80	75.02	16.00	81.52	18.24	361.03	152.40
N <sub>3</sub>	73.50	16.17	83.50	18.21	368.97	156.57	78.50	16.77	85.00	19.00	317.77	158.17
N <sub>4</sub>	75.15	16.53	85.15	18.56	374.13	159.57	80.15	17.30	86.82	19.47	377.18	160.99
SEm(±)	1.00	0.22	1.00	0.23	3.05	1.76	0.87	0.23	1.00	0.24	3.05	1.73
LSD (0.05)	2.92	0.64	2.92	0.68	8.90	5.16	2.56	0.67	2.93	0.77	8.92	5.07
F × N	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS

N: Nitrogen (kg ha⁻¹); P: Phosphorus (kg ha⁻¹); K: Potassium (kg ha⁻¹); S: Sulphur (kg ha⁻¹); Zn: Zinc (g ha⁻¹); B: Boron (g ha⁻¹)

nutrient uptake was noticed under (F<sub>1</sub>), respectively, from all the rest of the treatments. During 2023, the highest macronutrient uptake in respect of N, P, K and S (91.44; 19.78; 98.10; 21.38 kg ha⁻¹) and micronutrient uptake in respect of Zn and B (413.95; 175.28 g ha⁻¹) was observed in F<sub>4</sub> followed by F<sub>3</sub> and F<sub>2</sub> and lower nutrient uptake was recorded in F<sub>1</sub>. Among the macro and micro nutrients, maximum N, P, K and S uptake (75.15; 16.53; 85.15; 18.56 kg ha⁻¹) and micronutrient uptake (374.13; 159.57 g ha⁻¹) was observed under the treatment N<sub>4</sub> whereas a minimum nutrient uptake was achieved in N<sub>1</sub> during the crop season of 2022. During 2023, again significantly highest uptake of N, P, K and S (80.15; 17.30; 86.82; 19.47 kg ha⁻¹) and Zn and B (377.18; 160.99 g ha⁻¹) was recorded in N<sub>4</sub>, followed by N<sub>3</sub>, N<sub>2</sub> and N<sub>1</sub>. The essential macronutrients improved nutrient uptake by increasing their concentration in the soil, making them more accessible even in a smaller active root zone. In addition, application of FYM enhanced nutrient uptake efficiency. FYM contributes organic

acids that help in the chelation and availability of micronutrients like zinc and boron. It also supports microbial activity, which facilitates the breakdown of organic matter and the release of nutrients. Moreover, FYM improves soil texture and water-holding capacity, creating a more favorable environment for root growth and nutrient absorption. These combined effects led to improved nutrient availability and uptake, which in turn supported better plant metabolism, higher yield and improved seed and oil quality (33-35).

#### Available nutrients in soil

After the harvest of the mustard crop, available macronutrient (N, P, K, S) and micronutrient (Zn and B) content of soil significantly increased with the combined application of fertilizers as per treatments, as well as macro and micro nutrients. The highest amount of available macronutrients with respect to N, P, K and S (151.25; 15.28; 123.34; 24.44) and

micronutrients with respect to Zn and B (0.57; 0.56) were recorded under the treatment F<sub>4</sub> during 2022 (Table 5). Again, a similar trend was observed during 2023. Amongst macro and micro nutrients, maximum availability of macronutrients (150.47; 15.22; 123.04; 24.92 kg ha<sup>-1</sup> N, P, K and S, respectively) and micronutrients (0.56; 0.55 g ha<sup>-1</sup> Zn and B, respectively) was recorded under N<sub>4</sub>, followed by N<sub>3</sub>, N<sub>2</sub> and N<sub>1</sub> during 2022. Similarly, trend was observed in 2023. However, the interaction effect was found to be non-significant. The direct addition of N, P, K and micro nutrients in soil enhanced the availability of nutrients in soil pool (36-39).

### Microbial population

The microbial population in the soil was significantly affected with fertility levels and macro and micro nutrients. The highest numerical value of bacteria, fungi and actinomycetes, respectively (5.88; 3.81; 2.63) were observed under fertility levels F<sub>4</sub> followed by F<sub>3</sub> (5.74; 3.57; 2.45) and F<sub>2</sub> (5.60; 3.47; 2.38) while lowest value were recorded under F<sub>1</sub> (5.33; 3.44; 2.30) during both the growing seasons. A significant increase in mean microbial population (bacteria, fungi and actinomycetes) was observed under N<sub>4</sub> (5.93; 3.83; 2.65), respectively, than the rest of the treatments during 2022 and 2023 (Fig. 3).

### Economics

During 2022, F<sub>4</sub> gave maximum net returns and B: C ratio, the values being 68175 and 2.80, respectively followed by F<sub>3</sub> (64244; 2.74) and F<sub>2</sub> (57365; 2.55) whereas minimum net returns and B: C ratio (34542; 1.83) were obtained under control (F<sub>1</sub>) treatment. Similarly, the highest net return and B: C ratio was recorded in F<sub>4</sub> (71664; 2.95) and the lowest were observed in F<sub>1</sub> (38222; 2.03), respectively, during 2023.


**Table 5.** Response of fertility levels and integrated nutrient management on available nutrients in soil

Treatments	2022-23						2023-24					
	N	P	K	S	Zn	B	N	P	K	S	Zn	B
<b>F<sub>1</sub></b>	145.67	14.87	120.24	23.45	0.52	0.51	145.91	15.18	120.46	22.63	0.54	0.52
<b>F<sub>2</sub></b>	146.39	15.12	122.56	23.75	0.53	0.52	147.83	15.48	123.80	23.72	0.56	0.53
<b>F<sub>3</sub></b>	148.85	15.17	122.70	24.37	0.56	0.54	150.33	15.49	123.94	24.67	0.58	0.55
<b>F<sub>4</sub></b>	151.25	15.28	123.34	24.44	0.57	0.56	152.74	15.57	124.59	24.80	0.59	0.57
<b>SEm(±)</b>	1.17	0.08	0.63	0.20	0.01	0.004	1.15	0.07	0.61	0.25	0.01	0.004
<b>LSD (0.05)</b>	NS	NS	NS	0.70	0.02	0.01	NS	NS	NS	0.79	0.03	0.01
<b>Fertility Levels</b>												
<b>N<sub>1</sub></b>	144.96	15.00	121.41	23.10	0.53	0.51	146.38	15.33	122.64	23.41	0.53	0.53
<b>N<sub>2</sub></b>	147.61	15.09	122.18	23.54	0.54	0.53	149.07	15.43	123.42	23.86	0.57	0.54
<b>N<sub>3</sub></b>	149.11	15.13	122.20	24.45	0.55	0.54	150.58	15.43	123.44	24.81	0.58	0.57
<b>N<sub>4</sub></b>	150.47	15.22	123.04	24.92	0.56	0.55	151.97	15.54	124.29	25.34	0.59	0.58
<b>SEm(±)</b>	1.48	0.04	0.37	0.17	0.01	0.006	1.47	0.03	0.36	0.16	0.01	0.005
<b>LSD (0.05)</b>	NS	NS	NS	0.50	0.02	0.01	NS	NS	NS	0.52	0.03	0.01
<b>F × N</b>	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
<b>Macro and micro nutrients</b>												
<b>N<sub>1</sub></b>	144.96	15.00	121.41	23.10	0.53	0.51	146.38	15.33	122.64	23.41	0.53	0.53
<b>N<sub>2</sub></b>	147.61	15.09	122.18	23.54	0.54	0.53	149.07	15.43	123.42	23.86	0.57	0.54
<b>N<sub>3</sub></b>	149.11	15.13	122.20	24.45	0.55	0.54	150.58	15.43	123.44	24.81	0.58	0.57
<b>N<sub>4</sub></b>	150.47	15.22	123.04	24.92	0.56	0.55	151.97	15.54	124.29	25.34	0.59	0.58
<b>SEm(±)</b>	1.48	0.04	0.37	0.17	0.01	0.006	1.47	0.03	0.36	0.16	0.01	0.005
<b>LSD (0.05)</b>	NS	NS	NS	0.50	0.02	0.01	NS	NS	NS	0.52	0.03	0.01
<b>F × N</b>	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS


N: Nitrogen (kg ha<sup>-1</sup>); P: Phosphorus (kg ha<sup>-1</sup>); K: Potassium (kg ha<sup>-1</sup>); S: Sulphur (kg ha<sup>-1</sup>); Zinc (mg kg<sup>-1</sup>); B: Boron (mg kg<sup>-1</sup>)

**Table 6.** Response of fertility levels and integrated nutrient management on oil content (%), net return (₹ ha<sup>-1</sup>) and B:C ratio

Treatment	2022-23			2023-24		
	Oil content	Net return	B: C ratio	Oil content	Net return	B: C ratio
<b>Fertility Levels</b>						
<b>F<sub>1</sub></b>	33.90	34542	1.83	34.02	38222	2.03
<b>F<sub>2</sub></b>	37.35	57365	2.55	37.46	61558	2.74
<b>F<sub>3</sub></b>	38.11	64244	2.74	38.19	71227	3.05
<b>F<sub>4</sub></b>	38.67	68175	2.80	38.79	71664	2.95
<b>SEm(±)</b>	0.53	1539	0.07	0.53	1934	0.09
<b>LSD (0.05)</b>	1.85	5311	0.25	1.84	6674	0.31
<b>Macro and micro nutrients</b>						
<b>N<sub>1</sub></b>	35.27	47038	2.25	35.38	56301	2.69
<b>N<sub>2</sub></b>	36.08	49265	2.12	36.20	56612	2.43
<b>N<sub>3</sub></b>	38.05	63864	2.91	38.14	64530	2.95
<b>N<sub>4</sub></b>	38.63	64160	2.64	38.74	65228	2.69
<b>SEm(±)</b>	0.32	1797	0.08	0.31	1052	0.05
<b>LSD (0.05)</b>	0.94	5247	0.23	0.95	3072	0.15
<b>F × N</b>	NS	NS	NS	NS	NS	NS



**Fig. 3.** Response of fertility levels and integrated nutrient management on total mean microbial population of Indian mustard.



**Fig. 4.** Principal component analysis (PCA) among yield attributes and nutrient uptake of Indian mustard. SY: Seed yield; SPP: Siliqua plant<sup>-1</sup>; SPS: Seed siliqua<sup>-1</sup>; TW: Test weight; NU: Nitrogen uptake; PU: Phosphorus uptake; KU: Potassium uptake; SU: Sulphur uptake; ZnU: Zinc uptake and BU: boron uptake

In 2022 and 2023 (Fig. 5 & 6), Pearson's correlation analysis demonstrated a significant positive correlation among seed yield, siliqua plant<sup>-1</sup>, seeds siliqua<sup>-1</sup>, test weight, N, P, K, S, Zn and B uptake. The linear regression graph containing mean values for both the years of experimentation pertaining to nutrient uptake presented in Fig. 6 revealed that a mean positive correlation ( $r = 0.94, 0.94, 0.94, 0.93, 0.95$  and  $0.95$ ) between seed yield and N, P, K, S, Zn and B uptake respectively was observed.

## Conclusion

Based on the above results after two years of experimentation, it may be concluded that F3 (N:P2O5:K2O at 100:50:25 kg ha<sup>-1</sup>) when supplemented with application of N3 (20 kg S+ 2.5 kg Zinc+ 0.5 kg Boron enriched with 500 kg FYM ha<sup>-1</sup>) was found to be the most economical treatment which resulted in significant increase in the nutrient uptake of nitrogen, phosphorus, potassium, sulphur, boron and zinc and increased the seed


yield significantly due to higher yield attributing characters than other treatments and hence may be recommended for getting higher and sustainable seed yield of Indian mustard in North Western Himalayan region of India.

## Acknowledgements


The authors extend their thanks to the ICAR-Directorate of Rapeseed Mustard Research (Indian Council of Agricultural Research), Bharatpur, Rajasthan, India, for funding the research through the All India Coordinated Research Project on Rapeseed Mustard Research Project.

## Authors' contributions

RB and VB wrote the original manuscript and prepared tables and figures. SK, MG and DC reviewed the manuscript. BK, SKR and NP participated in the sequence alignment. All authors read and approved the final manuscript.



**Fig. 5.** Pearson's correlation between yield attributes and nutrient uptake of Indian mustard. SY: Seed yield; SPP: Siliqua plant<sup>-1</sup>; SPS: Seed siliqua<sup>-1</sup>; TW: Test weight; NU: Nitrogen uptake; PU: Phosphorus uptake; KU: Potassium uptake; SU: Sulphur uptake; ZnU: Zinc uptake and BU: Boron uptake.



**Fig. 6.** The correlation between mean grain yield and mean nutrient uptake ( $\text{kg ha}^{-1}$ ) of Indian mustard under different fertility levels and integrated nutrient management.

## Compliance with ethical standards

**Conflict of interest:** Authors do not have any conflict of interest to declare.

**Ethical issues:** None

## References

1. AICRP (Rapeseed-Mustard). Annual Progress Report. ICAR-Directorate of Rapeseed-Mustard Research, Sewar, Bharatpur, Rajasthan; 2023
2. NITI Aayog. NITI Aayog unveils report on Pathways and Strategies for Accelerating Growth in Edible Oils Towards Atmanirbharta [Internet] [Accessed on 02 Feb 2025]. <https://pib.gov.in/PressReleasePage.aspx?PRID=2049737>

3. Bharat R, Gupta V, Gupta M, Rai SK. Effect of different sowing schedules and planting geometry on yield and productivity of *Brassica juncea* L. *Bangladesh J Bot.* 2022;51(3):631–35. <https://doi.org/10.3329/bjb.v51i3.62011>
4. Singh AK, Kumar P. Nutrient management in rainfed-dry land agro-eco systems in the impending climate change scenario. *Agricultural Situation in India.* 2009;66(5):265–69.
5. Das SK, Biswas B, Jana K. Effect of farm yard manure, phosphorous and sulphur on yield parameters, yield, nodulation, nutrient uptake and quality of chickpea (*Cicer arietinum* L.). *J Appl Nat Sci.* 2016;8(2):545–49. <https://doi.org/10.31018/jans.v8i2.833>
6. Dubey SK, Tripathi SK, Singh B. Effect of sulphur and zinc levels on growth, yield and quality of mustard (*Brassica juncea* L. Czern & Coss). *J Crop Sci Tech.* 2013;2(1):1–11.
7. Kumar V, Tyagi S, Paul SC, Dubey SK, Suman S. Effect of sources and doses of sulphur on S uptake and yield of mustard (*Brassica juncea* L.). *Int J Curr Microbiol Appl Sci.* 2018;7:5042–47.
8. Das K, Dang R, Shivananda TN, Sur P. Interaction between phosphorous and zinc on the biomass yield and yield attributes of the medical plant stevia (*Stevia rebaudiana*). *Sci World J.* 2005;5:390–95. <https://doi.org/10.1100/tsw.2005.49>
9. Maqsood MA, Rahmatullah KS, Aziz T, Ashraf M. Evaluation of Zn distribution among grain and straw of twelve indigenous wheat (*Triticum aestivum* L.) genotypes. *Pak J Bot.* 2009;41(1):225–31.
10. Dordas C, Brown PH. Boron deficiency affects cell viability, phenolic leakage and oxidative burst in rose cell cultures. *Plant Cell Physiol.* 2005;46:293–301. <https://doi.org/10.1007/s11104-004-0309-1>
11. Kour S, Arora S, Jalali VK, Bali AS, Gupta M. Direct and residual effect of sulphur fertilization on yield, uptake and use efficiency in Indian mustard and succeeding rice crop. *J Plant Nutr.* 2014;37(14):2291–301. <https://doi.org/10.1080/01904167.2014.920389>
12. Kour S, Gupta M, Bharat R, Sharma V. Effect of zinc and boron on yield, nutrient uptake and economics of mustard (*Brassica Juncea* L.) in mustard-maize cropping sequence. *Bangladesh J Bot.* 2017;46(2):817–21. <https://doi.org/10.5958/0974-0228.2017.00039.1>
13. Kour S, Gupta M, Kachroo D, Bharat R. Direct and residual effect of zinc and boron on productivity, nutrient uptake and quality on mustard (*Brassica juncea*) and succeeding maize (*Zea mays*) in subtropical inceptisols of Jammu. *J Indian Soc Soil Sci.* 2017;65(3):334–40. <https://doi.org/10.5958/0974-0228.2017.00039.1>
14. Sadasivam S, Manickam A. Biochemical methods for agricultural sciences. New Delhi: Wiley Eastern Limited; 1992
15. Subbaiah BV. A rapid procedure for estimation of available nitrogen in soil. *Curr Sci.* 1956;25:259–60.
16. Olsen SR. Estimation of available phosphorous in soils by extraction with sodium bicarbonate. US Department of Agriculture Press Inc. 1954;939.
17. Hanway JJ, Heidel H. Soil analysis methods as used in Iowa State College Soil Testing Laboratory. *Iowa Agric.* 1952;57:1–13.
18. Chesnin L, Yien CH. Turbid metric method for determination of sulphur. *Soil Sci Soc Am J.* 1950;4(1):149–51. <https://doi.org/10.2136/sssaj1951.036159950015000C0032x>
19. John KK, Chuah HH, Neufeld JH. Application of improved azomethine-H method to determination of boron in soils and plants. *Anal Lett.* 1975;8(8):559–68. <https://doi.org/10.1080/00032717508058240>
20. Rao SNS. Soil microorganism and plant growth. New Delhi: Oxford and IBH Publishing Co; 1999
21. Bhari NR, Siaz RK, Mann PS. Response of Indian mustard (*Brassica juncea* L.) to nitrogen and phosphorous on torripsamments of North Western Rajasthan. *Indian J Agron.* 2000;45(4):746–51. <https://doi.org/10.5979/ija.v45i4.3440>
22. Premi OP, Kumar M. Response of Indian mustard to different levels of nitrogen and phosphorous under rainfed condition. *Indian J Agron.* 2004;38(1):47–48.
23. Alloway BJ. Zinc in soils and crop nutrition (2nd Eds). IZA and IFA Publication; 2008
24. Shireen F, Nawaz MA, Chen C, Zhang Q, Zheng Z, Sohail H, et al. Boron: Functions and approaches to enhance its availability in plants for sustainable agriculture. *Int J Mol Sci.* 2018;19:1856. <https://doi.org/10.3390/ijms19071856>
25. Jain NK, Sharma PP. Integrated nutrient management in mustard. *Indian J Agron.* 2000;45(1):127–29.
26. Roul PK, Sarawgi SK, Shrivastava GK, Kumar D. Effect of integrated nutrient management techniques on productivity, nitrogen uptake, N use efficiency, economics and energetics of rice (*Oryza sativa*)-Indian mustard (*Brassica juncea*) sequence. *Indian J Agron.* 2006;51(3):170–73. <https://doi.org/10.5979/ija.v51i3.5000>
27. Rao KT, Rao AU, Sekhar D. Effect of sources and levels of sulphur on groundnut. *J Acad Ind Res.* 2013;2(5):268–70.
28. Sipai AH, Patel JJ, Patel NI. Effect of sulphur and zinc with and without FYM on yield and yield attributes of mustard (*Brassica juncea* L.) grown on light textured soil of Kachchh. *Asian J Soil Sci.* 2015;10(2):191–200. <https://doi.org/10.15740/has/ajss/10.2/191-200>
29. Singh SP, Pal MS. Effect of integrated nutrient management on productivity, quality, nutrient uptake and economics of mustard (*Brassica juncea* L.). *Indian J Agron.* 2011;56(4):381–87. <https://doi.org/10.5979/ija.v56i4.4704>
30. Tripathi MK, Chaturvedi S, Shukla DK, Mahapatra BS. Yield performance and quality in Indian mustard (*Brassica juncea*) as affected by integrated nutrient management. *Indian J Agron.* 2010;55(2):138–42. <https://doi.org/10.5979/ija.v55i2.4731>
31. Sah D, Sewak R, Singh AK, Swami S. Growth, yield and profitability of Indian mustard *Brassica juncea* (L.) czern. & coss.) with different weed control measures and sulphur levels. *Agric Sci Digest.* 2013;33(1):15–20.
32. Bonde AS, Gawande SN. Effect of integrated nutrient management of growth, yield and nutrient uptake by soybean (*Glycine max*). *Ann Plant Soil Res.* 2017;19(2):154–58.
33. Jain NK, Vyas AK, Singh AK. Effect of phosphorous and sulphur fertilization on growth and nutrient uptake by mustard (*Brassica juncea* L. Czern and Coss). *Ann Agric Sci.* 1995;16:389–90.
34. Chakraborty AK, Das DK. Interaction between B and S on different quality parameters of rape (*Brassica campestris*). *Res Crops.* 2000;1:326–29.
35. Balaguravaiah D, Adinorayana G, Prathap S, Yalla MRT. Influence of long-term use of inorganic and organic manure on soil fertility and sustainable productivity of rainfed groundnut in alfisols. *J Ind Soc Soil Sci.* 2005;53:608–11.
36. Arya RL, Varshney JG, Kumar L. Effect of integrated nutrient application in Chickpea + Mustard intercropping system in the semi-arid tropics of North India. *Commun Soil Sci Plant Anal.* 2007;38(1-2):229–40. <https://doi.org/10.1080/00103620601094189>
37. Singh S, Singh V, Chandel BS. Effect of integrated nutrient management on Indian mustard and soil fertility. *Ann Agric Res New Series.* 2013;34(3):231–35.
38. Meena MK, Yadav BL, Dotaniya ML, Meena MD, Jat RS, Choudhary RL, et al. Can plant nutrient dynamics and Indian mustard yield be influenced by addition of organic amendments under saline sodic soils. *Commun Soil Sci Plant Anal.* 2024;55(6):875–87. <https://doi.org/10.1080/00103624.2023.2283000>
39. Choudhary P, Jhajharia A, Kumar R. Influence of sulphur and zinc fertilization on yield, yield components and quality traits of soybean (*Glycine max* L.). *Bioscan.* 2014;9(1):137–42.

**Additional information**

**Peer review:** Publisher thanks Sectional Editor and the other anonymous reviewers for their contribution to the peer review of this work.

**Reprints & permissions information** is available at [https://horizonpublishing.com/journals/index.php/PST/open\\_access\\_policy](https://horizonpublishing.com/journals/index.php/PST/open_access_policy)

**Publisher's Note:** Horizon e-Publishing Group remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

**Indexing:** Plant Science Today, published by Horizon e-Publishing Group, is covered by Scopus, Web of Science, BIOSIS Previews, Clarivate Analytics, NAAS, UGC Care, etc

See [https://horizonpublishing.com/journals/index.php/PST/indexing\\_abstracting](https://horizonpublishing.com/journals/index.php/PST/indexing_abstracting)

**Copyright:** © The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited (<https://creativecommons.org/licenses/by/4.0/>)

**Publisher information:** Plant Science Today is published by HORIZON e-Publishing Group with support from Empirion Publishers Private Limited, Thiruvananthapuram, India.